

5

## **THERAPEUTIC EPITOPES AND USES THEREOF**

The invention relates to epitopes useful in the diagnosis and therapy of coeliac disease, including diagnostics, therapeutics, kits, and methods of using the foregoing.

10 An immune reaction to gliadin (a component of gluten) in the diet causes coeliac disease. It is known that immune responses in the intestinal tissue preferentially respond to gliadin which has been modified by an intestinal transglutaminase. Coeliac disease is diagnosed by detection of anti-endomysial antibodies, but this requires confirmation by the finding of a lymphocytic inflammation in intestinal biopsies. The taking of such a biopsy is inconvenient  
15 for the patient.

Investigators have previously assumed that only intestinal T cell responses provide an accurate indication of the immune response against gliadins. Therefore they have concentrated on the investigation of T cell responses in intestinal tissue<sup>1</sup>. Gliadin epitopes which require transglutaminase modification  
20 (before they are recognised by the immune system) are known<sup>2</sup>.

The inventors have found the immunodominant T cell A-gliadin epitope recognised by the immune system in coeliac disease, and have shown that this is recognised by T cells in the peripheral blood of individuals with coeliac disease (see WO 01/25793). Such T cells were found to be present at high enough  
25 frequencies to be detectable without restimulation (i.e. a 'fresh response' detection system could be used). The epitope was identified using a non-T cell cloning based method which provided a more accurate reflection of the epitopes being recognised. The immunodominant epitope requires transglutaminase modification (causing substitution of a particular glutamine to glutamate) before  
30 immune system recognition.

Based on this work the inventors have developed a test which can be used to diagnose coeliac disease at an early stage. The test may be carried out on a sample from peripheral blood and therefore an intestinal biopsy is not required. The test is more sensitive than the antibody tests which are currently being used.

5           The invention thus provides a method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual comprising:

          (a) contacting a sample from the host with an agent selected from (i) the epitope comprising sequence which is: SEQ ID NO:1 (PQPELPY) or SEQ ID NO:2 (QLQFPQPPELPYPQPQS), or an equivalent sequence from a naturally occurring homologue of the gliadin represented by SEQ ID NO:3, (ii) an epitope comprising sequence comprising: SEQ ID NO:1, or an equivalent sequence from a naturally occurring homologue of the gliadin represented by SEQ ID NO:3 (shown in Table 1), which epitope is an isolated oligopeptide derived from a gliadin protein, (iii) an analogue of (i) or (ii) which is capable of being recognised by a T cell receptor that recognises (i) or (ii), which in the case of a peptide analogue is not more than 50 amino acids in length, or (iv) a product comprising two or more agents as defined in (i), (ii) or (iii), and (b) determining *in vitro* whether T cells in the sample recognise the agent, recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease.

          Through comprehensive mapping of wheat gliadin T cell epitopes (see Example 13), the inventors have also found epitopes bioactive in coeliac disease in HLA-DQ2+ patients in other wheat gliadins, having similar core sequences (e.g., SEQ ID NOS:18-22) and similar full length sequences (e.g., SEQ ID NOS:31-36), as well as in rye secalins and barley hordeins (e.g., SEQ ID NOS:39-41); see also Tables 20 and 21. Additionally, several epitopes bioactive in coeliac disease in HLA-DQ8+ patients have been identified (e.g., SEQ ID NOS:42-44, 46). This comprehensive mapping thus provides the dominant epitopes recognized by T cells in coeliac patients. Thus, the above-described method and other methods of the invention described herein may be performed using any of these additional identified epitopes, and analogues and equivalents thereof; (i) and (ii) herein include these additional epitopes. That is, the agents of the invention also include these novel epitopes.

          The invention also provides use of the agent for the preparation of a diagnostic means for use in a method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual, said method comprising

5 determining whether T cells of the individual recognise the agent, recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease.

The finding of an immunodominant epitope which is modified by transglutaminase (as well as the additional other epitopes defined herein) also allows diagnosis of coeliac disease based on determining whether other types of immune response to this epitope are present. Thus the invention also provides a method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual comprising determining the presence of an antibody that binds to the epitope in a sample from the individual, the presence of the antibody indicating that the individual has, or is susceptible to, coeliac disease.

15 The invention additionally provides the agent, optionally in association with a carrier, for use in a method of treating or preventing coeliac disease by tolerising T cells which recognise the agent. Also provided is an antagonist of a T cell which has a T cell receptor that recognises (i) or (ii), optionally in association with a carrier, for use in a method of treating or preventing coeliac disease by antagonising such T cells. Additionally provided is the agent or an analogue that binds an antibody (that binds the agent) for use in a method of treating or preventing coeliac disease in an individual by tolerising the individual to prevent the production of such an antibody.

25 The invention provides a method of determining whether a composition is capable of causing coeliac disease comprising determining whether a protein capable of being modified by a transglutaminase to an oligopeptide sequence as defined above is present in the composition, the presence of the protein indicating that the composition is capable of causing coeliac disease.

30 The invention also provides a mutant gliadin protein whose wild-type sequence can be modified by a transglutaminase to a sequence that comprises an epitope comprising sequence as defined above, but which mutant gliadin protein has been modified in such a way that it does not contain sequence which can be modified by a transglutaminase to a sequence that comprises such an epitope comprising sequence; or a fragment of such a mutant gliadin protein which is at

5 least 15 amino acids long and which comprises sequence which has been  
modified in said way.

The invention also provides a protein that comprises a sequence which is  
able to bind to a T cell receptor, which T cell receptor recognises the agent, and  
which sequence is able to cause antagonism of a T cell that carries such a T cell  
10 receptor.

Additionally the invention provides a food that comprises the proteins  
defined above.

### **SUMMARY OF THE INVENTION**

15 The present invention provides methods of preventing or treating coeliac disease  
comprising administering to an individual at least one agent selected from: a) a  
peptide comprising at least one epitope comprising a sequence selected from the  
group consisting of SEQ ID NOs:18-22, 31-36, 39-44, and 46, and equivalents  
thereof; and b) an analogue of a) which is capable of being recognised by a T cell  
20 receptor that recognises the peptide of a) and which is not more than 50 amino  
acids in length; and c) optionally, in addition to the agent selected from a) and b),  
a peptide comprising at least one epitope comprising a sequence selected from  
SEQ ID NO:1 and SEQ ID NO:2. In some embodiments, the agent is HLA-DQ2-  
restricted, HLA-DQ8-restricted or one agent is HLA-DQ2-restricted and a second  
25 agent is HLA-DQ8-restricted. In some embodiments, the agent comprises a wheat  
epitope, a rye epitope, a barley epitope or any combination thereof either as a  
single agent or as multiple agents.

The present invention also provides methods of preventing or treating  
coeliac disease comprising administering to an individual a pharmaceutical  
30 composition comprising an agent above and pharmaceutically acceptable carrier  
or diluent.

The present invention also provides methods of preventing or treating  
coeliac disease comprising administering to an individual a pharmaceutical  
composition comprising an antagonist of a T cell which has a T cell receptor as  
35 defined above, and a pharmaceutically acceptable carrier or diluent.

5           The present invention also provides methods of preventing or treating coeliac disease comprising administering to an individual a composition for tolerising an individual to a gliadin protein to suppress the production of a T cell or antibody response to an agent as defined above, which composition comprises an agent as defined above.

10           The present invention also provides methods of preventing or treating coeliac disease by 1) diagnosing coeliac disease in an individual by either: a) contacting a sample from the host with at least one agent selected from: i) a peptide comprising at least one epitope comprising a sequence selected from the group consisting of: SEQ ID NOS:18-22, 31-36, 39-44, and 46, and equivalents  
15           thereof; and ii) an analogue of i) which is capable of being recognised by a T cell receptor that recognises i) and which is not more than 50 amino acids in length; and iii) optionally, in addition to the agent selected from i) and ii), a peptide comprising at least one epitope comprising a sequence selected from SEQ ID NOS:1 and 2; and determining *in vitro* whether T cells in the sample recognise  
20           the agent; recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease; or b) administering an agent as defined above and determining *in vivo* whether T cells in the individual recognise the agent, recognition of the agent indicating that the individual has or is susceptible to coeliac disease; and 2) administering to an individual diagnosed as having, or  
25           being susceptible to, coeliac disease a therapeutic agent for preventing or treating coeliac disease.

          The present invention also provides agents as defined above, optionally in association with a carrier, for use in a method of treating or preventing coeliac disease by tolerising T cells which recognise the agent.

30           The present invention also provides antagonists of a T cell which has a T cell receptor as defined above, optionally in association with a carrier, for use in a method of treating or preventing coeliac disease by antagonising such T cells.

          The present invention also provides proteins that comprises a sequence which is able to bind to a T cell receptor, which T cell receptor recognises an

5 agent as defined above, and which sequence is able to cause antagonism of a T cell that carries such a T cell receptor.

The present invention also provides pharmaceutical compositions comprising an agent or antagonist as defined and a pharmaceutically acceptable carrier or diluent.

10 The present invention also provides compositions for tolerising an individual to a gliadin protein to suppress the production of a T cell or antibody response to an agent as defined above, which composition comprises an agent as defined above.

The present invention also provides compositions for antagonising a T cell response to an agent as defined above, which composition comprises an antagonist as defined above.

The present invention also provides mutant gliadin proteins whose wild-type sequence can be modified by a transglutaminase to a sequence which is an agent as defined in claim 1, which mutant gliadin protein comprises a mutation  
20 which prevents its modification by a transglutaminase to a sequence which is an agent as defined above; or a fragment of such a mutant gliadin protein which is at least 15 amino acids long and which comprises the mutation.

The present invention also provides polynucleotides that comprises a coding sequence that encodes a protein or fragment as defined above.

25 The present invention also provides cells comprising a polynucleotide as defined above or which has been transformed with such a polynucleotide.

The present invention also provides mammals that expresses a T cell receptor as defined above.

The present invention also provides methods of diagnosing coeliac  
30 disease, or susceptibility to coeliac disease, in an individual comprising: a) contacting a sample from the host with at least one agent selected from i) a peptide comprising at least one epitope comprising a sequence selected from the group consisting of: SEQ ID NOS:18-22, 31-36, 39-44, and 46, and equivalents thereof; and ii) an analogue of i) which is capable of being recognised by a T cell  
35 receptor that recognises i) and which is not more than 50 amino acids in length;

5 and iii) optionally, in addition to the agent selected from i) and ii), a peptide comprising at least one epitope comprising a sequence selected from SEQ ID NOS:1 and 2; and b) determining *in vitro* whether T cells in the sample recognise the agent; recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease.

10 The present invention also provides methods of determining whether a composition is capable of causing coeliac disease comprising determining whether a protein capable of being modified by a transglutaminase to an oligopeptide sequence is present in the composition, the presence of the protein indicating that the composition is capable of causing coeliac disease.

15 The present invention also provides methods of identifying an antagonist of a T cell, which T cell recognises an agent as defined above, comprising contacting a candidate substance with the T cell and detecting whether the substance causes a decrease in the ability of the T cell to undergo an antigen specific response, the detecting of any such decrease in said ability indicating that  
20 the substance is an antagonist.

The present invention also provides kits for carrying out any of the method described above comprising an agent as defined above and a means to detect the recognition of the peptide by the T cell.

25 The present invention also provides methods of identifying a product which is therapeutic for coeliac disease comprising administering a candidate substance to a mammal as defined above which has, or which is susceptible to, coeliac disease and determining whether substance prevents or treats coeliac disease in the mammal, the prevention or treatment of coeliac disease indicating that the substance is a therapeutic product.

30 The present invention also provides processes for the production of a protein encoded by a coding sequence as defined above which process comprises:  
a) cultivating a cell described above under conditions that allow the expression of the protein; and optionally b) recovering the expressed protein.

5           The present invention also provides methods of obtaining a transgenic plant cell comprising transforming a plant cell with a vector as described above to give a transgenic plant cell.

          The present invention also provides methods of obtaining a first-generation transgenic plant comprising regenerating a transgenic plant cell  
10 transformed with a vector as described above to give a transgenic plant.

          The present invention also provides methods of obtaining a transgenic plant seed comprising obtaining a transgenic seed from a transgenic plant obtainable as described above.

          The present invention also provides methods of obtaining a transgenic  
15 progeny plant comprising obtaining a second-generation transgenic progeny plant from a first-generation transgenic plant obtainable by a method as described above, and optionally obtaining transgenic plants of one or more further generations from the second-generation progeny plant thus obtained.

          The present invention also provides transgenic plant cells, plants, plant  
20 seeds or progeny plants obtainable by any of the methods described above.

          The present invention also provides transgenic plants or plant seeds comprising plant cells as described above.

          The present invention also provides transgenic plant cell calluses  
25 comprising plant cells as described above obtainable from a transgenic plant cell, first-generation plant, plant seed or progeny as defined above.

          The present invention also provides methods of obtaining a crop product comprising harvesting a crop product from a plant according to any method described above and optionally further processing the harvested product.

30           The present invention also provides food that comprises a protein as defined above.

## **BRIEF DESCRIPTION OF THE DRAWINGS**

          The invention is illustrated by the accompanying drawings in which:



5           Figure 1 shows freshly isolated PBMC (peripheral blood mononuclear cell) IFN $\gamma$  ELISPOT responses (vertical axis shows spot forming cells per  $10^6$  PBMC) to transglutaminase (tTG)-treated and untreated peptide pool 3 (each peptide 10  $\mu$ g/ml) including five overlapping 15mers spanning A-gliadin 51-85 (see Table 1) and a-chymotrypsin-digested gliadin (40  $\mu$ g/ml) in coeliac disease  
 10   Subject 1, initially in remission following a gluten free diet then challenged with 200g bread daily for three days from day 1 (a). PBMC IFN $\gamma$  ELISPOT responses by Subject 2 to tTG-treated A-gliadin peptide pools 1-10 spanning the complete A-gliadin protein during ten day bread challenge (b). The horizontal axis shows days after commencing bread.

15           Figure 2 shows PBMC IFN $\gamma$  ELISPOT responses to tTG-treated peptide pool 3 (spanning A-gliadin 51-85) in 7 individual coeliac disease subjects (vertical axis shows spot forming cells per  $10^6$  PBMC), initially in remission on gluten free diet, challenged with bread for three days (days 1 to 3). The horizontal axis shows days after commencing bread. (a). PBMC IFN $\gamma$  ELISPOT responses to  
 20   tTG-treated overlapping 15mer peptides included in pool 3; bars represent the mean ( $\pm$  SEM) response to individual peptides (10  $\mu$ g/ml) in 6 Coeliac disease subjects on day 6 or 7(b). (In individual subjects, ELISPOT responses to peptides were calculated as a % of response elicited by peptide 12 - as shown by the vertical axis.)

25           Figure 3 shows PBMC IFN $\gamma$  ELISPOT responses to tTG-treated truncations of A-gliadin 56-75 (0.1  $\mu$ M). Bars represent the mean ( $\pm$  SEM) in 5 Coeliac disease subjects. (In individual subjects, responses were calculated as the % of the maximal response elicited by any of the peptides tested.)

30           Figure 4 shows how the minimal structure of the dominant A-gliadin epitope was mapped using tTG-treated 7-17mer A-gliadin peptides (0.1  $\mu$ M) including the sequence, PQPQLPY (SEQ ID NO:4) (A-gliadin 62-68) (a), and the same peptides without tTG treatment but with the substitution Q $\rightarrow$ E65 (b). Each line represents PBMC IFN $\gamma$  ELISPOT responses in each of three Coeliac disease subjects on day 6 or 7 after bread was ingested on days 1-3. (In individual

5 subjects, ELISPOT responses were calculated as a % of the response elicited by the 17mer, A-gliadin 57-73.)

Figure 5 shows the amino acids that were deamidated by tTG. A-gliadin 56-75 LQLQPFPPQQLPYPPQSF (SEQ ID NO:5) (0.1  $\mu$ M) was incubated with tTG (50  $\mu$ g/ml) at 37°C for 2 hours. A single product was identified and purified by reverse phase HPLC. Amino acid analysis allowed % deamidation (Q→E) of each Gln residue in A-gliadin 56-75 attributable to tTG to be calculated (vertical axis).

Figure 6 shows the effect of substituting Q→E in A-gliadin 57-73 at other positions in addition to Q65 using the 17mers: QLQPFPPQPELPYPQPES (SEQ ID NO:6) (E57,65), QLQPFPPQPELPYPQPES (SEQ ID NO:7) (E65,72), ELQPFPPQPELPYPQPES (SEQ ID NO:8) (E57, 65, 72), and QLQPFPPQPELPYPQPQS (SEQ ID NO:2) (E65) in three Coeliac disease subjects on day 6 or 7 after bread was ingested on days 1-3. Vertical axis shows % of the E65 response.

Figure 7 shows that tTG treated A-gliadin 56-75 (0.1  $\mu$ M) elicited IFN- $\gamma$  ELISPOT responses in (a) CD4 and CD8 magnetic bead depleted PBMC. (Bars represent CD4 depleted PBMC responses as a % of CD8 depleted PBMC responses; spot forming cells per million CD8 depleted PBMC were: Subject 4: 29, and Subject 6: 535). (b) PBMC IFN $\gamma$  ELISPOT responses (spot forming cells/million PBMC) after incubation with monoclonal antibodies to HLA-DR (L243), -DQ (L2) and -DP (B7.21) (10  $\mu$ g/ml) 1h prior to tTG-treated 56-75 (0.1  $\mu$ M) in two coeliac disease subjects homozygous for HLA-DQ a1\*0501, b1\*0201.

Figure 8 shows the effect of substituting Glu at position 65 for other amino acids in the immunodominant epitope. The vertical axis shows the response in the 3 subjects in relation to the immunodominant epitope.

Figure 9 shows the immunoreactivity of naturally occurring gliadin peptides (measuring responses from 3 subjects) which contain the sequence PQLPY (SEQ ID NO:12) with (shaded) and without (clear) transglutaminase treatment.

5           Figure 10 shows CD8, CD4,  $\beta_7$ , and  $\alpha^E$ -specific immunomagnetic bead depletion of peripheral blood mononuclear cells from two coeliac subjects 6 days after commencing gluten challenge followed by interferon gamma ELISpot. A-gliadin 57-73 QE65 (25mcg/ml), tTG-treated chymotrypsin-digested gliadin (100 mcg/ml) or PPD (10 mcg/ml) were used as antigen.

10           Figure 11 shows the optimal T cell epitope length.

          Figure 12 shows a comparison of A-gliadin 57-73 QE65 with other peptides in a dose response study. On Sheet 12 of 47, Fig. 12(a) discloses the amino acid sequence SEQ ID NO:2. On Sheet 12 of 47, Fig. 12(b) discloses the amino acid sequence SEQ ID NO:101. On Sheet 13 of 47, Fig.12(c) discloses the  
 15 amino acid sequence SEQ ID NO:10. On Sheet 13 of 47, Fig.12(d) discloses the amino acid sequence SEQ ID NO:72. On Sheet 14 of 47, Fig.12(e) discloses the amino acid sequence SEQ ID NO:13 (labelled E65) and amino acid sequence SEQ ID NO:53 (labelled Q65). On Sheet 15 of 47, Fig.12(f) discloses the amino acid sequence SEQ ID NO:47 (labelled E65) and amino acid sequence SEQ ID  
 20 NO:102 (labelled Q65). On Sheet 16 of 47, Fig.12(g) discloses the amino acid sequence SEQ ID NO:99 (labelled E) and amino acid sequence SEQ ID NO:44 (labelled Q). On Sheet 17 of 47, Fig.12(h) discloses the amino acid sequence SEQ ID NO:48 (labelled E) and amino acid sequence SEQ ID NO:103 (labelled Q).

25           Figure 13 shows a comparison of gliadin and A-gliadin 57-73 QE65 specific responses.

          Figure 14 shows the bioactivity of gliadin polymorphisms in coeliac subjects, On Sheets 20 through 23 of 47, sixteen amino acids are identified (A-P) in the legend of each figure. The amino acid sequences A through P correspond to  
 30 the following sequence identifiers:

A – SEQ ID NO: 10

B – SEQ ID NO: 26

C – SEQ ID NO: 51

D – SEQ ID NO: 104

35           E – SEQ ID NO: 68

- 5        F – SEQ ID NO:28  
        G – SEQ ID NO: 69  
        H – SEQ ID NO:70  
        I – SEQ ID NO:71  
        J – SEQ ID NO:105  
 10       K – SEQ ID NO:72  
        L – SEQ ID NO:73  
        M – SEQ ID NO:74  
        N – SEQ ID NO:75  
        O – SEQ ID NO:97  
 15       P – SEQ ID NO:77

Figures 15 and 16 show the defining of the core epitope sequence.

Figures 17 to 27 show the agonist activity of A-gliadin 57-73 QE65 variants. On Sheets 25 through 35 of 47, Figs. 17 through 27 disclose the amino acid sequence SEQ ID NO:2.

- 20       Figure 28 shows responses in different patient groups.  
        Figure 29 shows bioactivity of prolamins homologues of A-gliadin 57-73.  
        Figure 30 shows, for healthy HLA-DQ2 subjects, the change in IFN-  
        gamma ELISpot responses to tTG-deamidated gliadin peptide pools.  
        Figure 31 shows, for coeliac HLA-DQ2 subjects, the change in IFN-  
 25       gamma ELISpot responses to tTG-deamidated gliadin peptide pools.  
        Figure 32 shows individual peptide contributions to “summed” gliadin  
        peptide response.  
        Figure 33 shows, for coeliac HLA-DQ2/8 subject C08, gluten challenge  
        induced IFN $\gamma$  ELISpot responses to tTG-deamidated gliadin peptide pools.  
 30       Figure 34 shows, for coeliac HLA-DQ2/8 subject C07, gluten challenge  
        induced IFN $\gamma$  ELISpot responses to tTG-deamidated gliadin peptide pools.  
        Figure 35 shows, for coeliac HLA-DQ8/7 subject C12, gluten challenge  
        induced IFN $\gamma$  ELISpot responses to tTG-deamidated gliadin peptide pools.  
        Figure 36 shows, for coeliac HLA-DQ6/8 subject C11, gluten challenge  
 35       induced IFN $\gamma$  ELISpot responses to tTG-deamidated gliadin peptide pools.

5

### Detailed Description of the Invention

The term "coeliac disease" encompasses a spectrum of conditions caused by varying degrees of gluten sensitivity, including a severe form characterised by a flat small intestinal mucosa (hyperplastic villous atrophy) and other forms characterised by milder symptoms.

The individual mentioned above (in the context of diagnosis or therapy) is human. They may have coeliac disease (symptomatic or asymptomatic) or be suspected of having it. They may be on a gluten free diet. They may be in an acute phase response (for example they may have coeliac disease, but have only ingested gluten in the last 24 hours before which they had been on a gluten free diet for 14 to 28 days).

The individual may be susceptible to coeliac disease, such as a genetic susceptibility (determined for example by the individual having relatives with coeliac disease or possessing genes which cause predisposition to coeliac disease).

### The agent

The agent is typically a peptide, for example of length 7 to 50 amino acids, such as 10 to 40, or 15 to 30 amino acids in length.

SEQ ID NO:1 is PQPELPY. SEQ ID NO:2 is QLQPFQPELPYPQPQS. SEQ ID NO:3 is shown in Table 1 and is the sequence of a whole A-gliadin. The glutamate at position 4 of SEQ ID NO:1 (equivalent to position 9 of SEQ ID NO:2) is generated by transglutaminase treatment of A-gliadin.

The agent may be the peptide represented by SEQ ID NO:1 or 2 or an epitope comprising sequence that comprises SEQ ID NO:1 which is an isolated oligopeptide derived from a gliadin protein; or an equivalent of these sequences from a naturally occurring gliadin protein which is a homologue of SEQ ID NO:3. Thus the epitope may be a derivative of the protein represented by SEQ ID NO:3. Such a derivative is typically a fragment of the gliadin, or a mutated derivative of the whole protein or fragment. Therefore the epitope of the

5 invention does not include this naturally occurring whole gliadin protein, and does not include other whole naturally occurring gliadins.

The epitope may thus be a fragment of A-gliadin (e.g. SEQ ID NO:3), which comprises the sequence of SEQ ID NO:1, obtainable by treating (fully or partially) with transglutaminase, i.e. with 1, 2, 3 or more glutamines substituted to  
10 glutamates (including the substitution within SEQ ID NO:1).

Such fragments may be or may include the sequences represented by positions 55 to 70, 58 to 73, 61 to 77 of SEQ ID NO:3 shown in Table 1. Typically such fragments will be recognised by T cells to at least the same extent that the peptides represented by SEQ ID NO:1 or 2 are recognised in any of the  
15 assays described herein using samples from coeliac disease patients.

Additionally, the agent may be the peptide represented by any of SEQ ID NOS:18-22, 31-36, 39-44, and 46 or a protein comprising a sequence corresponding to any of SEQ ID NOS:18-22, 31-36, 39-44, and 46 (such as fragments of a gliadin comprising any of SEQ ID NOS:18-22, 31-36, 39-44, and  
20 46, for example after the gliadin has been treated with transglutaminase). Bioactive fragments of such sequences are also agents of the invention. Sequences equivalent to any of SEQ ID NOS:18-22, 31-36, 39-44, and 46 or analogues of these sequences are also agents of the invention.

In the case where the epitope comprises a sequence equivalent to the  
25 above epitopes (including fragments) from another gliadin protein (e.g. any of the gliadin proteins mentioned herein or any gliadins which cause coeliac disease), such equivalent sequences will correspond to a fragment of a gliadin protein typically treated (partially or fully) with transglutaminase. Such equivalent peptides can be determined by aligning the sequences of other gliadin proteins  
30 with the gliadin from which the original epitope derives, such as with SEQ ID NO:3 (for example using any of the programs mentioned herein). Transglutaminase is commercially available (e.g. Sigma T-5398). Table 4 provides a few examples of suitable equivalent sequences.

The agent which is an analogue is capable of being recognised by a TCR  
35 which recognises (i) or (ii). Therefore generally when the analogue is added to T

5 cells in the presence of (i) or (ii), typically also in the presence of an antigen presenting cell (APC) (such as any of the APCs mentioned herein), the analogue inhibits the recognition of (i) or (ii), i.e. the analogue is able to compete with (i) or (ii) in such a system.

The analogue may be one which is capable of binding the TCR which  
10 recognises (i) or (ii). Such binding can be tested by standard techniques. Such TCRs can be isolated from T cells which have been shown to recognise (i) or (ii) (e.g. using the method of the invention). Demonstration of the binding of the analogue to the TCRs can then shown by determining whether the TCRs inhibit the binding of the analogue to a substance that binds the analogue, e.g. an  
15 antibody to the analogue. Typically the analogue is bound to a class II MHC molecule (e.g. HLA-DQ2) in such an inhibition of binding assay.

Typically the analogue inhibits the binding of (i) or (ii) to a TCR. In this case the amount of (i) or (ii) which can bind the TCR in the presence of the analogue is decreased. This is because the analogue is able to bind the TCR and  
20 therefore competes with (i) or (ii) for binding to the TCR.

T cells for use in the above binding experiments can be isolated from patients with coeliac disease, for example with the aid of the method of the invention. Other binding characteristics of the analogue may also be the same as (i) or (ii), and thus typically the analogue binds to the same MHC class II  
25 molecule to which the peptide binds (HLA-DQ2 or -DQ8). The analogue typically binds to antibodies specific for (i) or (ii), and thus inhibits binding of (i) or (ii) to such antibodies.

The analogue is typically a peptide. It may have homology with (i) or (ii), typically at least 70% homology, preferably at least 80, 90%, 95%, 97% or 99%  
30 homology with (i) or (ii), for example over a region of at least 15 more (such as the entire length of the analogue and/or (i) or (ii), or across the region which contacts the TCR or binds the MHC molecule) contiguous amino acids. Methods of measuring protein homology are well known in the art and it will be understood by those of skill in the art that in the present context, homology is

calculated on the basis of amino acid identity (sometimes referred to as “hard homology”).

For example the UWGCG Package provides the BESTFIT program which can be used to calculate homology (for example used on its default settings) (Devereux *et al* (1984) *Nucleic Acids Research* **12**, p387-395). The PILEUP and BLAST algorithms can be used to calculate homology or line up sequences (typically on their default settings), for example as described in Altschul S. F. (1993) *J Mol Evol* 36:290-300; Altschul, S, F *et al* (1990) *J Mol Biol* 215:403-10.

Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information on the world wide web through the internet at, for example, “www.ncbi.nlm.nih.gov/”. This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al*, supra). These initial neighbourhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90: 5873-5787. One measure of similarity provided by the BLAST



5 algorithm is the smallest sum probability ( $P(N)$ ), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 10 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

The homologous peptide analogues typically differ from (i) or (ii) by 1, 2, 3, 4, 5, 6, 7, 8 or more mutations (which may be substitutions, deletions or insertions). These mutations may be measured across any of the regions 15 mentioned above in relation to calculating homology. The substitutions are preferably 'conservative'. These are defined according to the following Table. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

|           |                   |         |
|-----------|-------------------|---------|
| ALIPHATIC | Non-polar         | G A P   |
|           |                   | I L V   |
|           | Polar – uncharged | C S T M |
|           |                   | N Q     |
|           | Polar – charged   | D E     |
|           |                   | K R     |
| AROMATIC  |                   | H F W Y |

20

Typically the amino acids in the analogue at the equivalent positions to amino acids in (i) or (ii) that contribute to binding the MHC molecule or are responsible for the recognition by the TCR, are the same or are conserved.

Typically the analogue peptide comprises one or more modifications, 25 which may be natural post-translation modifications or artificial modifications. The modification may provide a chemical moiety (typically by substitution of a hydrogen, e.g. of a C-H bond), such as an amino, acetyl, hydroxy or halogen (e.g.

5 fluorine) group or carbohydrate group. Typically the modification is present on the N or C terminus.

The analogue may comprise one or more non-natural amino acids, for example amino acids with a side chain different from natural amino acids. Generally, the non-natural amino acid will have an N terminus and/or a C  
10 terminus. The non-natural amino acid may be an L- or a D- amino acid.

The analogue typically has a shape, size, flexibility or electronic configuration that is substantially similar to (i) or (ii). It is typically a derivative of (i) or (ii). In one embodiment the analogue is a fusion protein comprising the sequence of SEQ ID NO:1 or 2, or any of the other peptides mentioned herein;  
15 and non-gliadin sequence.

In one embodiment the analogue is or mimics (i) or (ii) bound to a MHC class II molecule. 2, 3, 4 or more of such complexes may be associated or bound to each other, for example using a biotin/streptavidin based system, in which typically 2, 3 or 4 biotin labelled MHC molecules bind to a streptavidin moiety.  
20 This analogue typically inhibits the binding of the (i) or (ii)/MHC Class II complex to a TCR or antibody which is specific for the complex.

The analogue is typically an antibody or a fragment of an antibody, such as a Fab or (Fab)<sub>2</sub> fragment. The analogue may be immobilised on a solid support, particularly an analogue that mimics peptide bound to a MHC molecule.

25 The analogue is typically designed by computational means and then synthesised using methods known in the art. Alternatively the analogue can be selected from a library of compounds. The library may be a combinatorial library or a display library, such as a phage display library. The library of compounds may be expressed in the display library in the form of being bound to a MHC  
30 class II molecule, such as HLA-DQ2 or -DQ8. Analogues are generally selected from the library based on their ability to mimic the binding characteristics (i) or (ii). Thus they may be selected based on ability to bind a TCR or antibody which recognises (i) or (ii).

Typically analogues will be recognised by T cells to at least the same  
35 extent as any of the agents (i) or (ii), for example at least to the same extent as the

5 equivalent epitope and preferably to the same extent as the peptide represented by  
SEQ ID NO:2, is recognised in any of the assays described herein, typically using  
T cells from coeliac disease patients. Analogues may be recognised to these  
extents *in vivo* and thus may be able to induce coeliac disease symptoms to at  
least the same extent as any of the agents mentioned herein (e.g. in a human  
10 patient or animal model).

Analogues may be identified in a method comprising determining whether  
a candidate substance is recognised by a T cell receptor that recognises an epitope  
of the invention, recognition of the substance indicating that the substance is an  
analogue. Such TCRs may be any of the TCRs mentioned herein, and may be  
15 present on T cells. Any suitable assay mentioned herein can be used to identify  
the analogue. In one embodiment this method is carried out *in vivo*. As  
mentioned above preferred analogues are recognised to at least the same extent as  
the peptide SEQ ID NO:2, and so the method may be used to identify analogues  
which are recognised to this extent.

20 In one embodiment the method comprises determining whether a  
candidate substance is able to inhibit the recognition of an epitope of the  
invention, inhibition of recognition indicating that the substance is an analogue.

The agent may be a product comprising at least 2, 5, 10 or 20 agents as  
defined by (i), (ii) or (iii). Typically the composition comprises epitopes of the  
25 invention (or equivalent analogues) from different gliadins, such as any of the  
species or variety of or types of gliadin mentioned herein. Preferred  
compositions comprise at least one epitope of the invention, or equivalent  
analogue, from all of the gliadins present in any of the species or variety  
mentioned herein, or from 2, 3, 4 or more of the species mentioned herein (such  
30 as from the panel of species consisting of wheat, rye, barley, oats and triticale).  
Thus, the agent may be monovalent or multivalent.

## Diagnosis

5 As mentioned above the method of diagnosis of the invention may be based on the detection of T cells that bind the agent or on the detection of antibodies that recognise the agent.

The T cells that recognise the agent in the method (which includes the use mentioned above) are generally T cells that have been pre-sensitised *in vivo* to  
10 gliadin. As mentioned above such antigen-experienced T cells have been found to be present in the peripheral blood.

In the method the T cells can be contacted with the agent *in vitro* or *in vivo*, and determining whether the T cells recognise the agent can be performed *in vitro* or *in vivo*. Thus the invention provides the agent for use in a method of  
15 diagnosis practiced on the human body. Different agents are provided for simultaneous, separate or sequential use in such a method.

The *in vitro* method is typically carried out in aqueous solution into which the agent is added. The solution will also comprise the T cells (and in certain embodiments the APCs discussed below). The term 'contacting' as used herein  
20 includes adding the particular substance to the solution.

Determination of whether the T cells recognise the agent is generally accomplished by detecting a change in the state of the T cells in the presence of the agent or determining whether the T cells bind the agent. The change in state is generally caused by antigen specific functional activity of the T cell after the  
25 TCR binds the agent. The change of state may be measured inside (e.g. change in intracellular expression of proteins) or outside (e.g. detection of secreted substances) the T cells.

The change in state of the T cell may be the start of or increase in secretion of a substance from the T cell, such as a cytokine, especially IFN- $\gamma$ , IL-  
30 2 or TNF- $\alpha$ . Determination of IFN- $\gamma$  secretion is particularly preferred. The substance can typically be detected by allowing it to bind to a specific binding agent and then measuring the presence of the specific binding agent/substance complex. The specific binding agent is typically an antibody, such as polyclonal or monoclonal antibodies. Antibodies to cytokines are commercially available, or  
35 can be made using standard techniques.

5           Typically the specific binding agent is immobilised on a solid support. After the substance is allowed to bind the solid support can optionally be washed to remove material which is not specifically bound to the agent. The agent/substance complex may be detected by using a second binding agent that will bind the complex. Typically the second agent binds the substance at a site  
10       which is different from the site which binds the first agent. The second agent is preferably an antibody and is labelled directly or indirectly by a detectable label.

          Thus the second agent may be detected by a third agent that is typically labelled directly or indirectly by a detectable label. For example the second agent may comprise a biotin moiety, allowing detection by a third agent which  
15       comprises a streptavidin moiety and typically alkaline phosphatase as a detectable label.

          In one embodiment the detection system which is used is the *ex-vivo* ELISPOT assay described in WO 98/23960. In that assay IFN- $\gamma$  secreted from the T cell is bound by a first IFN- $\gamma$  specific antibody that is immobilised on a  
20       solid support. The bound IFN- $\gamma$  is then detected using a second IFN- $\gamma$  specific antibody which is labelled with a detectable label. Such a labelled antibody can be obtained from MABTECH (Stockholm, Sweden). Other detectable labels which can be used are discussed below.

          The change in state of the T cell that can be measured may be the increase  
25       in the uptake of substances by the T cell, such as the uptake of thymidine. The change in state may be an increase in the size of the T cells, or proliferation of the T cells, or a change in cell surface markers on the T cell.

          In one embodiment the change of state is detected by measuring the change in the intracellular expression of proteins, for example the increase in  
30       intracellular expression of any of the cytokines mentioned above. Such intracellular changes may be detected by contacting the inside of the T cell with a moiety that binds the expressed proteins in a specific manner and which allows sorting of the T cells by flow cytometry.

          In one embodiment when binding the TCR the agent is bound to an MHC  
35       class II molecule (typically HLA-DQ2 or -DQ8), which is typically present on the

5 surface of an antigen presenting cell (APC). However as mentioned herein other agents can bind a TCR without the need to also bind an MHC molecule.

Generally the T cells which are contacted in the method are taken from the individual in a blood sample, although other types of samples which contain T cells can be used. The sample may be added directly to the assay or may be  
10 processed first. Typically the processing may comprise diluting of the sample, for example with water or buffer. Typically the sample is diluted from 1.5 to 100 fold, for example 2 to 50 or 5 to 10 fold.

The processing may comprise separation of components of the sample. Typically mononuclear cells (MCs) are separated from the samples. The MCs  
15 will comprise the T cells and APCs. Thus in the method the APCs present in the separated MCs can present the peptide to the T cells. In another embodiment only T cells, such as only CD4 T cells, can be purified from the sample. PBMCs, MCs and T cells can be separated from the sample using techniques known in the art, such as those described in Lavani *et al* (1997) *J. Exp. Med.* **186**, p859-865.

20 In one embodiment, the T cells used in the assay are in the form of unprocessed or diluted samples, or are freshly isolated T cells (such as in the form of freshly isolated MCs or PBMCs) which are used directly *ex vivo*, i.e. they are not cultured before being used in the method. Thus the T cells have not been restimulated in an antigen specific manner *in vitro*. However the T cells can be  
25 cultured before use, for example in the presence of one or more of the agents, and generally also exogenous growth promoting cytokines. During culturing the agent(s) are typically present on the surface of APCs, such as the APC used in the method. Pre-culturing of the T cells may lead to an increase in the sensitivity of the method. Thus the T cells can be converted into cell lines, such as short term  
30 cell lines (for example as described in Ota *et al* (1990) *Nature* **346**, p183-187).

The APC that is typically present in the method may be from the same individual as the T cell or from a different host. The APC may be a naturally occurring APC or an artificial APC. The APC is a cell that is capable of presenting the peptide to a T cell. It is typically a B cell, dendritic cell or  
35 macrophage. It is typically separated from the same sample as the T cell and is

5 typically co-purified with the T cell. Thus the APC may be present in MCs or PBMCs. The APC is typically a freshly isolated *ex vivo* cell or a cultured cell. It may be in the form of a cell line, such as a short term or immortalised cell line. The APC may express empty MHC class II molecules on its surface.

In the method one or more (different) agents may be used. Typically the T  
10 cells derived from the sample can be placed into an assay with all the agents which it is intended to test or the T cells can be divided and placed into separate assays each of which contain one or more of the agents.

The invention also provides the agents such as two or more of any of the agents mentioned herein (e.g. the combinations of agents which are present in the  
15 composition agent discussed above) for simultaneous separate or sequential use (eg. for *in vivo* use).

In one embodiment agent *per se* is added directly to an assay comprising T cells and APCs. As discussed above the T cells and APCs in such an assay could be in the form of MCs. When agents that can be recognised by the T cell  
20 without the need for presentation by APCs are used then APCs are not required. Analogues which mimic the original (i) or (ii) bound to a MHC molecule are an example of such an agent.

In one embodiment the agent is provided to the APC in the absence of the T cell. The APC is then provided to the T cell, typically after being allowed to  
25 present the agent on its surface. The peptide may have been taken up inside the APC and presented, or simply be taken up onto the surface without entering inside the APC.

The duration for which the agent is contacted with the T cells will vary depending on the method used for determining recognition of the peptide.  
30 Typically  $10^5$  to  $10^7$ , preferably  $5 \times 10^5$  to  $10^6$  PBMCs are added to each assay. In the case where agent is added directly to the assay its concentration is from  $10^{-1}$  to  $10^3$   $\mu\text{g/ml}$ , preferably 0.5 to 50  $\mu\text{g/ml}$  or 1 to 10  $\mu\text{g/ml}$ .

Typically the length of time for which the T cells are incubated with the agent is from 4 to 24 hours, preferably 6 to 16 hours. When using *ex vivo*

- 5 PBMCs it has been found that  $0.3 \times 10^6$  PBMCs can be incubated in  $10 \mu\text{g/ml}$  of peptide for 12 hours at  $37^\circ\text{C}$ .

The determination of the recognition of the agent by the T cells may be done by measuring the binding of the agent to the T cells (this can be carried out using any suitable binding assay format discussed herein). Typically T cells  
10 which bind the agent can be sorted based on this binding, for example using a FACS machine. The presence of T cells that recognise the agent will be deemed to occur if the frequency of cells sorted using the agent is above a "control" value. The frequency of antigen-experienced T cells is generally 1 in  $10^6$  to 1 in  $10^3$ , and therefore whether or not the sorted cells are antigen-experienced T cells can be  
15 determined.

The determination of the recognition of the agent by the T cells may be measured *in vivo*. Typically the agent is administered to the host and then a response which indicates recognition of the agent may be measured. The agent is typically administered intradermally or epidermally. The agent is typically  
20 administered by contacting with the outside of the skin, and may be retained at the site with the aid of a plaster or dressing. Alternatively the agent may be administered by needle, such as by injection, but can also be administered by other methods such as ballistics (e.g. the ballistics techniques which have been used to deliver nucleic acids). EP-A-0693119 describes techniques that can  
25 typically be used to administer the agent. Typically from 0.001 to 1000  $\mu\text{g}$ , for example from 0.01 to 100  $\mu\text{g}$  or 0.1 to 10  $\mu\text{g}$  of agent is administered.

In one embodiment a product can be administered which is capable of providing the agent *in vivo*. Thus a polynucleotide capable of expressing the agent can be administered, typically in any of the ways described above for the  
30 administration of the agent. The polynucleotide typically has any of the characteristics of the polynucleotide provided by the invention which is discussed below. The agent is expressed from the polynucleotide *in vivo*. Typically from 0.001 to 1000  $\mu\text{g}$ , for example from 0.01 to 100  $\mu\text{g}$  or 0.1 to 10  $\mu\text{g}$  of polynucleotide is administered.



5 Recognition of the agent administered to the skin is typically indicated by the occurrence of inflammation (e.g. induration, erythema or oedema) at the site of administration. This is generally measured by visual examination of the site.

The method of diagnosis based on the detection of an antibody that binds the agent is typically carried out by contacting a sample from the individual (such as any of the samples mentioned here, optionally processed in any manner mentioned herein) with the agent and determining whether an antibody in the sample binds the agent, such a binding indicating that the individual has, or is susceptible to coeliac disease. Any suitable format of binding assay may be used, such as any such format mentioned herein.

## 15 Therapy

The identification of the immunodominant epitope and other epitopes described herein allows therapeutic products to be made which target the T cells which recognise this epitope (such T cells being ones which participate in the immune response against gliadin). These findings also allow the prevention or treatment of coeliac disease by suppressing (by tolerisation) an antibody or T cell response to the epitope(s).

Certain agents of the invention bind the TCR that recognises the epitope of the invention (as measured using any of the binding assays discussed above) and cause tolerisation of the T cell that carries the TCR. Such agents, optionally in association with a carrier, can therefore be used to prevent or treat coeliac disease.

Generally tolerisation can be caused by the same peptides which can (after being recognised by the TCR) cause antigen specific functional activity of the T cell (such as any such activity mentioned herein, e.g. secretion of cytokines). Such agents cause tolerisation when they are presented to the immune system in a 'tolerising' context.

Tolerisation leads to a decrease in the recognition of a T cell or antibody epitope by the immune system. In the case of a T cell epitope this can be caused by the deletion or anergising of T cells that recognise the epitope. Thus T cell

activity (for example as measured in suitable assays mentioned herein) in response to the epitope is decreased. Tolerisation of an antibody response means that a decreased amount of specific antibody to the epitope is produced when the epitope is administered.

Methods of presenting antigens to the immune system in such a context are known and are described for example in Yoshida et al. Clin. Immunol. Immunopathol. 82, 207-215 (1997), Thureau et al. Clin. Exp. Immunol. 109, 370-6 (1997), and Weiner et al. Res. Immunol. 148, 528-33 (1997). In particular certain routes of administration can cause tolerisation, such as oral, nasal or intraperitoneal. Particular products which cause tolerisation may be administered (e.g. in a composition that also comprises the agent) to the individual. Such products include cytokines, such as cytokines that favour a Th2 response (e.g. IL-4, TGF- $\beta$  or IL-10). Products or agent may be administered at a dose that causes tolerisation.

The invention provides a protein that comprises a sequence able to act as an antagonist of the T cell (which T cell recognises the agent). Such proteins and such antagonists can also be used to prevent or treat coeliac disease. The antagonist will cause a decrease in the T cell response. In one embodiment, the antagonist binds the TCR of the T cell (generally in the form of a complex with HLA-DQ2 or -DQ8) but instead of causing normal functional activation causing an abnormal signal to be passed through the TCR intracellular signalling cascade, which causes the T cell to have decreased function activity (e.g. in response to recognition of an epitope, typically as measured by any suitable assay mentioned herein).

In one embodiment the antagonist competes with epitope to bind a component of MHC processing and presentation pathway, such as an MHC molecule (typically HLA-DQ2 or -DQ8). Thus the antagonist may bind HLA-DQ2 or -DQ8 (and thus be a peptide presented by this MHC molecule), such as peptide TP (Table 10) or a homologue thereof.

Methods of causing antagonism are known in the art. In one embodiment the antagonist is a homologue of the epitopes mentioned above and may have any

5 of the sequence, binding or other properties of the agent (particularly analogues). The antagonists typically differ from any of the above epitopes (which are capable of causing a normal antigen specific function in the T cell) by 1, 2, 3, 4 or more mutations (each of which may be a substitution, insertion or deletion). Such antagonists are termed “altered peptide ligands” or “APL” in the art. The  
10 mutations are typically at the amino acid positions that contact the TCR.

The antagonist may differ from the epitope by a substitution within the sequence that is equivalent to the sequence represented by amino acids 65 to 67 of A-gliadin (such antagonists are shown in Table 9). Thus preferably the antagonist has a substitution at the equivalent of position 64, 65 or 67. Preferably  
15 the substitution is 64W, 67W, 67M or 65T.

Since the T cell immune response to the epitope of the invention in an individual is polyclonal, more than one antagonist may need to be administered to cause antagonism of T cells of the response which have different TCRs. Therefore the antagonists may be administered in a composition which comprises  
20 at least 2, 4, 6 or more different antagonists, which each antagonise different T cells.

The invention also provides a method of identifying an antagonist of a T cell (which recognises the agent), comprising contacting a candidate substance with the T cell and detecting whether the substance causes a decrease in the  
25 ability of the T cell to undergo an antigen specific response (e.g. using any suitable assay mentioned herein), the detecting of any such decrease in said ability indicating that the substance is an antagonist.

In one embodiment, the antagonists (including combinations of antagonists to a particular epitope) or tolerising (T cell and antibody tolerising)  
30 agents are present in a composition comprising at least 2, 4, 6 or more antagonists or agents which antagonise or tolerate to different epitopes of the invention, for example to the combinations of epitopes discussed above in relation to the agents which are a product comprising more than one substance.

35 **Testing whether a composition is capable of causing coeliac disease**

5 As mentioned above the invention provides a method of determining whether a composition is capable of causing coeliac disease comprising detecting the presence of a protein sequence which is capable of being modified by a transglutaminase to as sequence comprising the agent or epitope of the invention (such transglutaminase activity may be a human intestinal transglutaminase activity). Typically this is performed by using a binding assay in which a moiety  
10 which binds to the sequence in a specific manner is contacted with the composition and the formation of sequence/moiety complex is detected and used to ascertain the presence of the agent. Such a moiety may be any suitable substance (or type of substance) mentioned herein, and is typically a specific  
15 antibody. Any suitable format of binding assay can be used (such as those mentioned herein).

In one embodiment, the composition is contacted with at least 2, 5, 10 or more antibodies which are specific for epitopes of the invention from different gliadins, for example a panel of antibodies capable of recognising the  
20 combinations of epitopes discussed above in relation to agents of the invention which are a product comprising more than one substance.

The composition typically comprises material from a plant that expresses a gliadin which is capable of causing coeliac disease (for example any of the gliadins or plants mentioned herein). Such material may be a plant part, such as a  
25 harvested product (e.g. seed). The material may be processed products of the plant material (e.g. any such product mentioned herein), such as a flour or food that comprises the gliadin. The processing of food material and testing in suitable binding assays is routine, for example as mentioned in Kricka LJ, J. Biolumin. Chemilumin. 13, 189-93 (1998).

30

### **Binding assays**

The determination of binding between any two substances mentioned herein may be done by measuring a characteristic of either or both substances that changes upon binding, such as a spectroscopic change.

5           The binding assay format may be a 'band shift' system. This involves determining whether the presence of one substance (such as a candidate substance) advances or retards the progress of the other substance during gel electrophoresis.

10           The format may be a competitive binding method which determines whether the one substance is able to inhibit the binding of the other substance to an agent which is known to bind the other substance, such as a specific antibody.

### **Mutant gliadin proteins**

15           The invention provides a gliadin protein in which an epitope sequence of the invention, or sequence which can be modified by a transglutaminase to provide such a sequence has been mutated so that it no longer causes, or is recognised by, a T cell response that recognises the epitope. In this context the term recognition refers to the TCR binding the epitope in such a way that normal (not antagonistic) antigen-specific functional activity of the T cell occurs.

20           Methods of identifying equivalent epitopes in other gliadins are discussed above. The wild type of the mutated gliadin is one which causes coeliac disease. Such a gliadin may have homology with SEQ ID NO:3, for example to the degree mentioned above (in relation to the analogue) across all of SEQ ID NO:3 or across 15, 30, 60, 100 or 200 contiguous amino acids of SEQ ID NO:3.

25           Likewise, for other non-A-gliadins, homology will be present between the mutant and the native form of that gliadin. The sequences of other natural gliadin proteins are known in the art.

          The mutated gliadin will not cause coeliac disease or will cause decreased symptoms of coeliac disease. Typically the mutation decreases the ability of the epitope to induce a T cell response. The mutated epitope may have a decreased binding to HLA-DQ2 or -DQ8, a decreased ability to be presented by an APC or a decreased ability to bind to or to be recognised (i.e. cause antigen-specific functional activity) by T cells that recognise the agent. The mutated gliadin or epitope will therefore show no or reduced recognition in any of the assays

35           mentioned herein in relation to the diagnostic aspects of the invention.

5           The mutation may be one or more deletions, additions or substitutions of length 1 to 3, 4 to 6, 6 to 10, 11 to 15 or more in the epitope, for example across sequence SEQ ID NO:2 or across any of SEQ ID NOS: 18-22, 31-36, 39-44, and 46; or across equivalents thereof. Preferably the mutant gliadin has at least one mutation in the sequence SEQ ID NO:1. A preferred mutation is at position 65 in  
10   A-gliadin (or in an equivalent position in other gliadins). Typically the naturally occurring glutamine at this position is substituted to any of the amino acids shown in Table 3, preferably to histidine, tyrosine, tryptophan, lysine, proline, or arginine.

          The invention thus also provides use of a mutation (such any of the  
15   mutations in any of the sequences discussed herein) in an epitope of a gliadin protein, which epitope is an epitope of the invention, to decrease the ability of the gliadin protein to cause coeliac disease.

          In one embodiment the mutated sequence is able to act as an antagonist. Thus the invention provides a protein that comprises a sequence which is able to  
20   bind to a T cell receptor, which T cell receptor recognises an agent of the invention, and which sequence is able to cause antagonism of a T cell that carries such a T cell receptor.

          The invention also provides proteins which are fragments of the above mutant gliadin proteins, which are at least 15 amino acids long (e.g. at least 30,  
25   60, 100, 150, 200, or 250 amino acids long) and which comprise the mutations discussed above which decrease the ability of the gliadin to be recognised. Any of the mutant proteins (including fragments) mentioned herein may also be present in the form of fusion proteins, for example with other gliadins or with non-gliadin proteins.

30           The equivalent wild type protein to the mutated gliadin protein is typically from a graminaceous monocotyledon, such as a plant of genus *Triticum*, e.g. wheat, rye, barley, oats or triticale. The protein is typically an  $\alpha$ ,  $\alpha\beta$ ,  $\beta$ ,  $\gamma$  or  $\omega$  gliadin. The gliadin may be an A-gliadin.

35   **Kits**

5           The invention also provides a kit for carrying out the method comprising one or more agents and optionally a means to detect the recognition of the agent by the T cell. Typically the different agents are provided for simultaneous, separate or sequential use. Typically the means to detect recognition allows or aids detection based on the techniques discussed above.

10           Thus the means may allow detection of a substance secreted by the T cells after recognition. The kit may thus additionally include a specific binding moiety for the substance, such as an antibody. The moiety is typically specific for IFN- $\gamma$ . The moiety is typically immobilised on a solid support. This means that after binding the moiety the substance will remain in the vicinity of the T cell which

15           secreted it. Thus “spots” of substance/moiety complex are formed on the support, each spot representing a T cell which is secreting the substance. Quantifying the spots, and typically comparing against a control, allows determination of recognition of the agent.

20           The kit may also comprise a means to detect the substance/moiety complex. A detectable change may occur in the moiety itself after binding the substance, such as a colour change. Alternatively a second moiety directly or indirectly labelled for detection may be allowed to bind the substance/moiety complex to allow the determination of the spots. As discussed above the second moiety may be specific for the substance, but binds a different site on the

25           substance than the first moiety.

          The immobilised support may be a plate with wells, such as a microtitre plate. Each assay can therefore be carried out in a separate well in the plate.

          The kit may additionally comprise medium for the T cells, detection moieties or washing buffers to be used in the detection steps. The kit may

30           additionally comprise reagents suitable for the separation from the sample, such as the separation of PBMCs or T cells from the sample. The kit may be designed to allow detection of the T cells directly in the sample without requiring any separation of the components of the sample.

          The kit may comprise an instrument which allows administration of the

35           agent, such as intradermal or epidermal administration. Typically such an

5 instrument comprises plaster, dressing or one or more needles. The instrument may allow ballistic delivery of the agent. The agent in the kit may be in the form of a pharmaceutical composition.

The kit may also comprise controls, such as positive or negative controls. The positive control may allow the detection system to be tested. Thus the  
10 positive control typically mimics recognition of the agent in any of the above methods. Typically in the kits designed to determine recognition *in vitro* the positive control is a cytokine. In the kit designed to detect *in vivo* recognition of the agent the positive control may be antigen to which most individuals should response.

15 The kit may also comprise a means to take a sample containing T cells from the host, such as a blood sample. The kit may comprise a means to separate mononuclear cells or T cells from a sample from the host.

#### **Polynucleotides, cells, transgenic mammals and antibodies**

20 The invention also provides a polynucleotide which is capable of expression to provide the agent or mutant gliadin proteins. Typically the polynucleotide is DNA or RNA, and is single or double stranded. The polynucleotide will preferably comprise at least 50 bases or base pairs, for example 50 to 100, 100 to 500, 500 to 1000 or 1000 to 2000 or more bases or  
25 base pairs. The polynucleotide therefore comprises a sequence which encodes the sequence of SEQ ID NO: 1 or 2 or any of the other agents mentioned herein. To the 5' and 3' of this coding sequence the polynucleotide of the invention has sequence or codons which are different from the sequence or codons 5' and 3' to these sequences in the corresponding gliadin gene.

30 5' and/or 3' to the sequence encoding the peptide the polynucleotide has coding or non-coding sequence. Sequence 5' and/or 3' to the coding sequence may comprise sequences which aid expression, such as transcription and/or translation, of the sequence encoding the agent. The polynucleotide may be capable of expressing the agent prokaryotic or eukaryotic cell. In one



5     embodiment the polynucleotide is capable of expressing the agent in a mammalian cell, such as a human, primate or rodent (e.g. mouse or rat) cell.

      A polynucleotide of the invention may hybridise selectively to a polynucleotide that encodes SEQ ID NO:3 at a level significantly above background. Selective hybridisation is typically achieved using conditions of  
10     medium to high stringency (for example 0.03M sodium chloride and 0.03M sodium citrate at from about 50°C to about 60°C). However, such hybridisation may be carried out under any suitable conditions known in the art (see Sambrook  
*et al* (1989), Molecular Cloning: A Laboratory Manual). For example, if high stringency is required, suitable conditions include 0.2 x SSC at 60°C. If lower  
15     stringency is required, suitable conditions include 2 x SSC at 60°C.

      Agents or proteins of the invention may be encoded by the polynucleotides described herein.

      The polynucleotide may form or be incorporated into a replicable vector. Such a vector is able to replicate in a suitable cell. The vector may be an  
20     expression vector. In such a vector the polynucleotide of the invention is operably linked to a control sequence which is capable of providing for the expression of the polynucleotide. The vector may contain a selectable marker, such as the ampicillin resistance gene.

      The polynucleotide or vector may be present in a cell. Such a cell may  
25     have been transformed by the polynucleotide or vector. The cell may express the agent. The cell will be chosen to be compatible with the said vector and may for example be a prokaryotic (bacterial), yeast, insect or mammalian cell. The polynucleotide or vector may be introduced into host cells using conventional techniques including calcium phosphate precipitation, DEAE-dextran  
30     transfection, or electroporation.

      The invention provides processes for the production of the proteins of the invention by recombinant means. This may comprise (a) cultivating a transformed cell as defined above under conditions that allow the expression of the protein; and preferably (b) recovering the expressed polypeptide. Optionally,  
35     the polypeptide may be isolated and/or purified, by techniques known in the art.

5           The invention also provides TCRs which recognise (or bind) the agent, or fragments thereof which are capable of such recognition (or binding). These can be present in the any form mentioned herein (e.g. purity) discussed herein in relation to the protein of the invention. The invention also provides T cells which express such TCRs which can be present in any form (e.g. purity) discussed  
10       herein for the cells of the invention.

          The invention also provides monoclonal or polyclonal antibodies which specifically recognise the agents (such as any of the epitopes of the invention) and which recognise the mutant gliadin proteins (and typically which do not recognise the equivalent wild-type gliadins) of the invention, and methods of  
15       making such antibodies. Antibodies of the invention bind specifically to these substances of the invention.

          For the purposes of this invention, the term "antibody" includes antibody fragments such as Fv, F(ab) and F(ab)<sub>2</sub> fragments, as well as single-chain antibodies.

20           A method for producing a polyclonal antibody comprises immunising a suitable host animal, for example an experimental animal, with the immunogen and isolating immunoglobulins from the serum. The animal may therefore be inoculated with the immunogen, blood subsequently removed from the animal and the IgG fraction purified. A method for producing a monoclonal antibody  
25       comprises immortalising cells which produce the desired antibody. Hybridoma cells may be produced by fusing spleen cells from an inoculated experimental animal with tumour cells (Kohler and Milstein (1975) *Nature* **256**, 495-497).

          An immortalized cell producing the desired antibody may be selected by a conventional procedure. The hybridomas may be grown in culture or injected  
30       intraperitoneally for formation of ascites fluid or into the blood stream of an allogenic host or immunocompromised host. Human antibody may be prepared by *in vitro* immunisation of human lymphocytes, followed by transformation of the lymphocytes with Epstein-Barr virus.

          For the production of both monoclonal and polyclonal antibodies, the  
35       experimental animal is suitably a goat, rabbit, rat or mouse. If desired, the

5 immunogen may be administered as a conjugate in which the immunogen is coupled, for example via a side chain of one of the amino acid residues, to a suitable carrier. The carrier molecule is typically a physiologically acceptable carrier. The antibody obtained may be isolated and, if desired, purified.

The polynucleotide, agent, protein or antibody of the invention, may carry  
10 a detectable label. Detectable labels which allow detection of the secreted substance by visual inspection, optionally with the aid of an optical magnifying means, are preferred. Such a system is typically based on an enzyme label which causes colour change in a substrate, for example alkaline phosphatase causing a colour change in a substrate. Such substrates are commercially available, e.g.  
15 from BioRad. Other suitable labels include other enzymes such as peroxidase, or protein labels, such as biotin; or radioisotopes, such as  $^{32}\text{P}$  or  $^{35}\text{S}$ . The above labels may be detected using known techniques.

Polynucleotides, agents, proteins, antibodies or cells of the invention may be in substantially purified form. They may be in substantially isolated form, in  
20 which case they will generally comprise at least 80% e.g. at least 90, 95, 97 or 99% of the polynucleotide, peptide, antibody, cells or dry mass in the preparation. The polynucleotide, agent, protein or antibody is typically substantially free of other cellular components. The polynucleotide, agent, protein or antibody may be used in such a substantially isolated, purified or free form in the method or be  
25 present in such forms in the kit.

The invention also provides a transgenic non-human mammal which expresses a TCR of the invention. This may be any of the mammals discussed herein (e.g. in relation to the production of the antibody). Preferably the mammal has, or is susceptible, to coeliac disease. The mammal may also express HLA-  
30 DQ2 or -DQ8 or HLA-DR3-DQ2 and/or may be given a diet comprising a gliadin which cause coeliac disease (e.g. any of the gliadin proteins mentioned herein). Thus the mammal may act as an animal model for coeliac disease.

The invention also provides a method of identifying a product which is therapeutic for coeliac disease comprising administering a candidate substance to  
35 a mammal of the invention which has, or which is susceptible to, coeliac disease

5 and determining whether substance prevents or treats coeliac disease in the mammal, the prevention or treatment of coeliac disease indicating that the substance is a therapeutic product. Such a product may be used to treat or prevent coeliac disease.

The invention provides therapeutic (including prophylactic) agents or  
10 diagnostic substances (the agents, proteins and polynucleotides of the invention). These substances are formulated for clinical administration by mixing them with a pharmaceutically acceptable carrier or diluent. For example they can be formulated for topical, parenteral, intravenous, intramuscular, subcutaneous, intraocular, intradermal, epidermal or transdermal administration. The substances  
15 may be mixed with any vehicle which is pharmaceutically acceptable and appropriate for the desired route of administration. The pharmaceutically carrier or diluent for injection may be, for example, a sterile or isotonic solution such as Water for Injection or physiological saline, or a carrier particle for ballistic delivery.

20 The dose of the substances may be adjusted according to various parameters, especially according to the agent used; the age, weight and condition of the patient to be treated; the mode of administration used; the severity of the condition to be treated; and the required clinical regimen. As a guide, the amount of substance administered by injection is suitably from 0.01 mg/kg to 30 mg/kg,  
25 preferably from 0.1 mg/kg to 10 mg/kg.

The routes of administration and dosages described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and dosage for any particular patient and condition.

The substances of the invention may thus be used in a method of  
30 treatment of the human or animal body, or in a diagnostic method practised on the human body. In particular they may be used in a method of treating or preventing coeliac disease. The invention also provide the agents for use in a method of manufacture of a medicament for treating or preventing coeliac disease. Thus the invention provides a method of preventing or treating coeliac disease comprising

- 5 administering to a human in need thereof a substance of the invention (typically a non-toxic effective amount thereof).

The agent of the invention can be made using standard synthetic chemistry techniques, such as by use of an automated synthesizer. The agent may be made from a longer polypeptide e.g. a fusion protein, which polypeptide  
10 typically comprises the sequence of the peptide. The peptide may be derived from the polypeptide by for example hydrolysing the polypeptide, such as using a protease; or by physically breaking the polypeptide. The polynucleotide of the invention can be made using standard techniques, such as by using a synthesiser.

15 **Plant cells and plants that express mutant gliadin proteins or express proteins comprising sequences which can act as antagonists**

The cell of the invention may be a plant cell, such as a cell of a graminaceous monocotyledonous species. The species may be one whose wild-type form expresses gliadins, such as any of the gliadin proteins mentioned herein  
20 (including gliadins with any degree of homology to SEQ ID NO:3 mentioned herein). Such a gliadin may cause coeliac disease in humans. The cell may be of wheat, maize, oats, rye, rice, barley, triticale, sorghum, or sugar cane. Typically the cell is of the Triticum genus, such as aestivum, spelta, polonicum or monococcum.

25 The plant cell of the invention is typically one which does not express a wild-type gliadin (such as any of the gliadins mentioned herein which may cause coeliac disease), or one which does not express a gliadin comprising a sequence that can be recognised by a T cell that recognises the agent. Thus if the wild-type plant cell did express such a gliadin then it may be engineered to prevent or  
30 reduce the expression of such a gliadin or to change the amino acid sequence of the gliadin so that it no longer causes coeliac disease (typically by no longer expressing the epitope of the invention).

This can be done for example by introducing mutations into 1, 2, 3 or more or all of such gliadin genes in the cell, for example into coding or non-  
35 coding (e.g. promoter regions). Such mutations can be any of the type or length

5 of mutations discussed herein (e.g., in relation to homologous proteins). The mutations can be introduced in a directed manner (e.g., using site directed mutagenesis or homologous recombination techniques) or in a random manner (e.g. using a mutagen, and then typically selecting for mutagenised cells which no longer express the gliadin (or a gliadin sequence which causes coeliac disease)).

10 In the case of plants or plant cells that express a protein that comprises a sequence able to act as an antagonist such a plant or plant cell may express a wild-type gliadin protein (e.g. one which causes coeliac disease). Preferably though the presence of the antagonist sequence will cause reduced coeliac disease symptoms (such as no symptoms) in an individual who ingests a food comprising  
15 protein from the plant or plant cell.

The polynucleotide which is present in (or which was transformed into) the plant cell will generally comprise promoter capable of expressing the mutant gliadin protein the plant cell. Depending on the pattern of expression desired, the promoter may be constitutive, tissue- or stage-specific; and/or inducible. For  
20 example, strong constitutive expression in plants can be obtained with the CAMV 35S, Rubisco ssu, or histone promoters. Also, tissue-specific or stage-specific promoters may be used to target expression of protein of the invention to particular tissues in a transgenic plant or to particular stages in its development. Thus, for example seed-specific, root-specific, leaf-specific, flower-specific etc  
25 promoters may be used. Seed-specific promoters include those described by Dalta *et al* (Biotechnology Ann. Rev. (1997), 3, pp.269-296). Particular examples of seed-specific promoters are napin promoters (EP-A-0 255, 378), phaseolin promoters, glutenine promoters, helianthene promoters (WO92/17580), albumin promoters (WO98/45460), oleosin promoters  
30 (WO98/45461) and ATS1 and ATS3 promoters (PCT/US98/06798).

The cell may be in any form. For example, it may be an isolated cell, e.g. a protoplast, or it may be part of a plant tissue, e.g. a callus, or a tissue excised from a plant, or it may be part of a whole plant. The cell may be of any type (e.g. of any type of plant part). For example, an undifferentiated cell, such as a callus  
35 cell; or a differentiated cell, such as a cell of a type found in embryos, pollen,

5 roots, shoots or leaves. Plant parts include roots; shoots; leaves; and parts involved in reproduction, such as pollen, ova, stamens, anthers, petals, sepals and other flower parts.

The invention provides a method of obtaining a transgenic plant cell comprising transforming a plant cell with a polynucleotide or vector of the invention to give a transgenic plant cell. Any suitable transformation method  
10 may be used (in the case of wheat the techniques disclosed in Vasil V et al, Biotechnology 10, 667-674 (1992) may be used). Preferred transformation techniques include electroporation of plant protoplasts and particle bombardment. Transformation may thus give rise to a chimeric tissue or plant in which some  
15 cells are transgenic and some are not.

The cell of the invention or thus obtained cell may be regenerated into a transgenic plant by techniques known in the art. These may involve the use of plant growth substances such as auxins, gibberellins and/or cytokinins to stimulate the growth and/or division of the transgenic cell. Similarly, techniques such as  
20 somatic embryogenesis and meristem culture may be used. Regeneration techniques are well known in the art and examples can be found in, e.g. US 4,459,355, US 4,536,475, US 5,464,763, US 5, 177,010, US 5, 187,073, EP 267,159, EP 604, 662, EP 672, 752, US 4,945,050, US 5,036,006, US 5,100,792, US 5,371,014, US 5,478,744, US 5,179,022, US 5,565,346, US 5,484,956, US  
25 5,508,468, US 5,538,877, US 5,554,798, US 5,489,520, US 5,510,318, US 5,204,253, US 5,405,765, EP 442,174, EP 486,233, EP 486,234, EP 539,563, EP 674,725, WO91/02071 and WO 95/06128.

In many such techniques, one step is the formation of a callus, i.e. a plant tissue comprising expanding and/or dividing cells. Such calli are a further aspect  
30 of the invention as are other types of plant cell cultures and plant parts. Thus, for example, the invention provides transgenic plant tissues and parts, including embryos, meristems, seeds, shoots, roots, stems, leaves and flower parts. These may be chimeric in the sense that some of their cells are cells of the invention and some are not. Transgenic plant parts and tissues, plants and seeds of the  
35 invention may be of any of the plant species mentioned herein.

5           Regeneration procedures will typically involve the selection of transformed cells by means of marker genes.

          The regeneration step gives rise to a first generation transgenic plant. The invention also provides methods of obtaining transgenic plants of further generations from this first generation plant. These are known as progeny  
10   transgenic plants. Progeny plants of second, third, fourth, fifth, sixth and further generations may be obtained from the first generation transgenic plant by any means known in the art.

          Thus, the invention provides a method of obtaining a transgenic progeny plant comprising obtaining a second-generation transgenic progeny plant from a  
15   first-generation transgenic plant of the invention, and optionally obtaining transgenic plants of one or more further generations from the second-generation progeny plant thus obtained.

          Progeny plants may be produced from their predecessors of earlier generations by any known technique. In particular, progeny plants may be  
20   produced by:

          obtaining a transgenic seed from a transgenic plant of the invention belonging to a previous generation, then obtaining a transgenic progeny plant of the invention belonging to a new generation by growing up the transgenic seed; and/or

25       propagating clonally a transgenic plant of the invention belonging to a previous generation to give a transgenic progeny plant of the invention belonging to a new generation; and/or

          crossing a first-generation transgenic plant of the invention belonging to a previous generation with another compatible plant to give a transgenic progeny  
30   plant of the invention belonging to a new generation; and optionally

          obtaining transgenic progeny plants of one or more further generations from the progeny plant thus obtained.

          These techniques may be used in any combination. For example, clonal propagation and sexual propagation may be used at different points in a process  
35   that gives rise to a transgenic plant suitable for cultivation. In particular,



5 repetitive back-crossing with a plant taxon with agronomically desirable characteristics may be undertaken. Further steps of removing cells from a plant and regenerating new plants therefrom may also be carried out.

Also, further desirable characteristics may be introduced by transforming the cells, plant tissues, plants or seeds, at any suitable stage in the above process,  
 10 to introduce desirable coding sequences other than the polynucleotides of the invention. This may be carried out by the techniques described herein for the introduction of polynucleotides of the invention.

For example, further transgenes may be selected from those coding for other herbicide resistance traits, e.g. tolerance to: Glyphosate (e.g. using an EPSP  
 15 synthase gene (e.g. EP-A-0 293,358) or a glyphosate oxidoreductase (WO 92/000377) gene); or tolerance to fosametin; a dihalobenzonitrile; glufosinate, e.g. using a phosphinothrycin acetyl transferase (PAT) or glutamine synthase gene (cf. EP-A-0 242,236); asulam, e.g. using a dihydropteroate synthase gene (EP-A-0 369,367); or a sulphonylurea, e.g. using an ALS gene); diphenyl ethers  
 20 such as acifluorfen or oxyfluorfen, e.g. using a protoporphyrinogen oxidase gene); an oxadiazole such as oxadiazon; a cyclic imide such as chlorophthalim; a phenyl pyrazole such as TNP, or a phenopylate or carbamate analogue thereof.

Similarly, genes for beneficial properties other than herbicide tolerance may be introduced. For example, genes for insect resistance may be introduced,  
 25 notably genes encoding *Bacillus thuringiensis* (*Bt*) toxins. Likewise, genes for disease resistance may be introduced, e.g. as in WO91/02701 or WO95/06128.

Typically, a protein of the invention is expressed in a plant of the invention. Depending on the promoter used, this expression may be constitutive or inducible. Similarly, it may be tissue- or stage-specific, i.e. directed towards a  
 30 particular plant tissue (such as any of the tissues mentioned herein) or stage in plant development.

The invention also provides methods of obtaining crop products by harvesting, and optionally processing further, transgenic plants of the invention. By crop product is meant any useful product obtainable from a crop plant.

5

**Products that contain mutant gliadin proteins or proteins that comprise sequence capable of acting as an antagonist**

The invention provides a product that comprises the mutant gliadin proteins or protein that comprises sequence capable of acting as an antagonist.

- 10 This is typically derived from or comprise plant parts from plants mentioned herein which express such proteins. Such a product may be obtainable directly by harvesting or indirectly, by harvesting and further processing the plant of the invention. Directly obtainable products include grains. Alternatively, such a product may be obtainable indirectly, by harvesting and further processing.
- 15 Examples of products obtainable by further processing are flour or distilled alcoholic beverages; food products made from directly obtained or further processed material, e.g. baked products (e.g. bread) made from flour. Typically such food products, which are ingestible and digestible (i.e. non-toxic and of nutrient value) by human individuals.

- 20 In the case of food products that comprise the protein which comprises an antagonist sequence the food product may also comprise wild-type gliadin, but preferably the antagonist is able to cause a reduction (e.g. completely) in the coeliac disease symptoms after such food is ingested.

The invention is illustrated by the following nonlimiting Examples:

25 **Example 1**

- We carried out epitope mapping in Coeliac disease by using a set of 51 synthetic 15-mer peptides that span the complete sequence of a fully characterized a-gliadin, "A-gliadin" (see Table 1). A-Gliadin peptides were also individually treated with tTG to generate products that might mimic those
- 30 produced in vivo<sup>3</sup>. We also sought to study Coeliac disease patients at the point of initiation of disease relapse to avoid the possibility that epitope "spreading" or "exhaustion" may have occurred, as described in experimental infectious and autoimmune diseases.

- 35 *Clinical and A-gliadin specific T cell responses with 3 and 10 day bread*

## 5 *challenge*

In a pilot study, two subjects with Coeliac disease in remission, defined by absence of serum anti-endomysial antibody (EMA), on a gluten free diet were fed four slices of standard gluten-containing white bread daily in addition to their usual gluten free diet. Subject 1 ceased bread because of abdominal pain, mouth  
 10 ulcers and mild diarrhoea after three days, but Subject 2 continued for 10 days with only mild nausea at one week. The EMA became positive in Subject 2 one week after the bread challenge, indicating the bread used had caused a relapse of Coeliac disease. But in Subject 1, EMA remained negative up to two months after bread challenge. In both subjects, symptoms that appeared with bread  
 15 challenge resolved within two days after returning to gluten free diet.

PBMC responses in IFN $\gamma$  ELISPOT assays to A-gliadin peptides were not found before or during bread challenge. But from the day after bread withdrawal (Day 4) in Subject 1 a single pool of 5 overlapping peptides spanning A-gliadin 51-85 (Pool 3) treated with tTG showed potent IFN $\gamma$  responses (see Figure 1a).  
 20 In Subject 1, the PBMC IFN $\gamma$  response to A-gliadin peptide remained targeted to Pool 3 alone and was maximal on Day 8. The dynamics and magnitude of the response to Pool 3 was similar to that elicited by  $\alpha$ -chymotrypsin digested gliadin. PBMC IFN $\gamma$  responses to tTG-treated Pool 3 were consistently 5 to 12-fold greater than Pool 3 not treated with tTG, and responses to  $\alpha$ -chymotrypsin  
 25 digested gliadin were 3 to 10-fold greater if treated with tTG. In Subject 2, Pool 3 treated with tTG was also the only immunogenic set of A-gliadin peptides on Day 8, but this response was weaker than Subject 1, was not seen on Day 4 and by Day 11 the response to Pool 3 had diminished and other tTG-treated pools of A-gliadin peptides elicited stronger IFN $\alpha$  responses (see Figure 1b).

30 The pilot study indicated that the initial T cell response in these Coeliac disease subjects was against a single tTG-treated A-gliadin pool of five peptides and was readily measured in peripheral blood. But if antigen exposure is continued for ten days instead of three, T cell responses to other A-gliadin peptides appear, consistent with epitope spreading.

### 5 *Coeliac disease-specific IFN- $\gamma$ induction by tTG-treated A-gliadin peptides*

In five out of six further Coeliac disease subjects on gluten free diet (see Table 1), bread challenge for three days identified tTG-treated peptides in Pool 3, and in particular, peptides corresponding to 56-70 (12) and 60-75 (13) as the sole A-gliadin components eliciting IFN $\gamma$  from PBMC (see Figure 2). IL-10  
 10 ELISPOT assays run in parallel to IFN $\gamma$  ELISPOT showed no IL-10 response to tTG-treated peptides 12 or 13. In one subject, there were no IFN $\gamma$  responses to any A-gliadin peptide or  $\alpha$ -chymotrypsin digested gliadin before, during or up to four days after bread challenge. In none of these Coeliac disease subjects did EMA status change from baseline when measured for up to two months after  
 15 bread challenge.

PBMC from four healthy, EMA-negative subjects with the HLA-DQ alleles  $\alpha 1^*0501$ ,  $\beta 1^*0201$  (ages 28-52, 2 females) who had been challenged for three days with bread after following a gluten free diet for one month, showed no IFN $\gamma$  responses above the negative control to any of the A-gliadin peptides with  
 20 or without tTG treatment. Thus, induction of IFN $\gamma$  in PBMC to tTG-treated Pool 3 and A-gliadin peptides 56-70 (12) and 60-75 (13) were Coeliac disease specific (7/8 vs. 0/4,  $p < 0.01$  by Chi-squared analysis).

### *Fine mapping of the minimal A-gliadin T cell epitope*

25 tTG-treated peptides representing truncations of A-gliadin 56-75 revealed that the same core peptide sequence QPQLP (SEQ ID NO:9) was essential for antigenicity in all of the five Coeliac disease subjects assessed (see Figure 3). PBMC IFN $\gamma$  responses to tTG-treated peptides spanning this core sequence beginning with the 7-mer PQPQLPY (SEQ ID NO:4) and increasing in length,  
 30 indicated that the tTG-treated 17-mer QLQPFQPQLPYQPQS (SEQ ID NO:10) (A-gliadin 57-73) possessed optimal activity in the IFN $\gamma$  ELISPOT (see Figure 4).

*Deamidation of Q65 by tTG generates the immunodominant T cell epitope in A-*  
 35 *gliadin*

5 HPLC analysis demonstrated that tTG treatment of A-gliadin 56-75 generated a single product that eluted marginally later than the parent peptide. Amino acid sequencing indicated that out of the six glutamine (Q) residues contained in A-gliadin 56-75, Q65 was preferentially deamidated by tTG (see Figure 5). Bioactivity of peptides corresponding to serial expansions from the core A-gliadin 62-68 sequence in which glutamate (E) replaced Q65, was  
 10 equivalent to the same peptides with Q65 after tTG-treatment (see Figure 4a). Replacement of Q57 and Q72 by E together or alone, with E65 did not enhance antigenicity of the 17-mer in the three Coeliac disease subjects studied (see Figure 6). Q57 and Q72 were investigated because glutamine residues followed  
 15 by proline in gliadin peptides are not deamidated by tTG in vitro (W. Vader et al, Proceedings 8th International Symposium Coeliac Disease). Therefore, the immunodominant T cell epitope was defined as QLQPFQPELPYPQPQS (SEQ ID NO:2).

20 *Immunodominant T cell epitope response is DQ2-restricted and CD4 dependent*

In two Coeliac disease subjects homozygous for HLA-DQ  $\alpha 1^*0501$ ,  $\beta 1^*0201$ , anti-DQ monoclonal antibody blocked the ELISPOT IFN $\gamma$  response to tTG-treated A-gliadin 56-75, but anti-DP and -DR antibody did not (see Figure 7). Anti-CD4 and anti-CD8 magnetic bead depletion of PBMC from two Coeliac  
 25 disease subjects indicated the IFN $\gamma$  response to tTG-treated A-gliadin 56-75 is CD4 T cell-mediated.

*Discussion*

In this study we describe a rather simple dietary antigen challenge using  
 30 standard white bread to elicit a transient population of CD4 T cells in peripheral blood of Coeliac disease subjects responsive to a tTG-treated A-gliadin 17-mer with the sequence: QLQPFQPELPYPQPQS (SEQ ID NO:2) (residues 57-73). The immune response to A-gliadin 56-75 (Q→E65) is restricted to the Coeliac disease-associated HLA allele, DQ  $\alpha 1^*0501$ ,  $\beta 1^*0201$ . Tissue transglutaminase  
 35 action in vitro selectively deamidates Q65. Elicited peripheral blood IFN $\gamma$

responses to synthetic A-gliadin peptides with the substitution Q→E65 is equivalent to tTG-treated Q65 A-gliadin peptides; both stimulate up to 10-fold more T cells in the IFN $\gamma$  ELISPOT than unmodified Q65 A-gliadin peptides.

We have deliberately defined this Coeliac disease-specific T cell epitope using in vivo antigen challenge and short-term ex vivo immune assays to avoid the possibility of methodological artifacts that may occur with the use of T cell clones in epitope mapping. Our findings indicate that peripheral blood T cell responses to ingestion of gluten are rapid but short-lived and can be utilized for epitope mapping. In vivo antigen challenge has also shown there is a temporal hierarchy of immune responses to A-gliadin peptides; A-gliadin 57-73 modified by tTG not only elicits the strongest IFN $\gamma$  response in PBMC but it is also the first IFN $\gamma$  response to appear.

Because we have assessed only peptides spanning A-gliadin, there may be other epitopes in other gliadins of equal or greater importance in the pathogenesis of Coeliac disease. Indeed, the peptide sequence at the core of the epitope in A-gliadin that we have identified PQQPLPY (SEQ ID NO:4) is shared by several other gliadins (SwissProt and TrEMBL accession numbers: P02863, Q41528, Q41531, Q41533, Q9ZP09, P04722, P04724, P18573). However, A-gliadin peptides that have previously been shown to possess bioactivity in biopsy challenge and in vivo studies (for example: 31-43, 44-55, and 206-217)<sup>4,5</sup> did not elicit IFN $\gamma$  responses in PBMC following three day bread challenge in Coeliac disease subjects. These peptides may be “secondary” T cell epitopes that arise with spreading of the immune response.

## Example 2

*The effect on T cell recognition of substitutions in the immunodominant epitope*

The effect of substituting the glutamate at position 65 in the 57-73 A-gliadin epitope was determined by measuring peripheral blood responses against the substituted epitopes in an IFN $\gamma$  ELISPOT assay using synthetic peptides (at 50  $\mu$ g/ml). The responses were measured in 3 Coeliac disease subjects 6 days after commencing gluten challenge (4 slices bread daily for 3 days). Results are

shown in table 3 and Figure 8. As can be seen substitution of the glutamate to histidine, tyrosine, tryptophan, lysine, proline or arginine stimulated a response whose magnitude was less than 10% of the magnitude of the response to the immunodominant epitope. Thus mutation of A-gliadin at this position could be used to produce a mutant gliadin with reduce or absent immunoreactivity.

### Example 3

*Testing the immunoreactivity of equivalent peptides from other naturally occurring gliadins*

The immunoreactivity of equivalent peptides from other naturally occurring wheat gliadins was assessed using synthetic peptides corresponding to the naturally occurring sequences which were then treated with transglutaminase. These peptides were tested in an ELISPOT in the same manner and with PBMCs from the same subjects as described in Example 2. At least five of the peptides show immunoreactivity comparable to the A-gliadin 57-73 E65 peptide (after transglutaminase treatment) indicating that other gliadin proteins in wheat are also likely to induce this Coeliac disease-specific immune response (Table 4 and Figure 9).

### Methods

**Subjects:** Patients used in the study attended a Coeliac Clinic in Oxford, United Kingdom. Coeliac disease was diagnosed on the basis of typical small intestinal histology, and normalization of symptoms and small intestinal histology with gluten free diet.

**Tissue typing:** Tissue typing was performed using DNA extracted from EDTA-anticoagulated peripheral blood. HLA-DQA and DQB genotyping was performed by PCR using sequence-specific primer mixes<sup>6-8</sup>.

**Anti-endomysial antibody assay:** EMA were detected by indirect

immunofluorescence using patient serum diluted 1:5 with monkey oesophagus,

5 followed by FITC-conjugated goat anti-human IgA. IgA was quantitated prior to EMA, none of the subjects were IgA deficient.

*Antigen Challenge:* Coeliac disease subjects following a gluten free diet, consumed 4 slices of gluten-containing bread (50g/slice, Sainsbury's "standard white sandwich bread") daily for 3 or 10 days. EMA was assessed the week before and up to two months after commencing the bread challenge. Healthy subjects who had followed a gluten free diet for four weeks, consumed their usual diet including four slices of gluten-containing bread for three days, then returned to gluten free diet for a further six days.

15 *IFN $\gamma$  and IL-10 ELISPOT:* PBMC were prepared from 50-100 ml of venous blood by Ficoll-Hypaque density centrifugation. After three washes, PBMC were resuspended in complete RPMI containing 10% heat inactivated human AB serum. ELISPOT assays for single cell secretion of IFN $\gamma$  and IL-10 were performed using commercial kits (Mabtech; Stockholm, Sweden) with 96-well plates (MAIP-S-45; Millipore, Bedford, MA) according to the manufacturers instructions (as described elsewhere<sup>9</sup>) with  $2.5 \times 10^5$  (IFN $\gamma$ ) or  $0.4 \times 10^5$  (IL-10) PBMC in each well. Peptides were assessed in duplicate wells, and Mycobacterium tuberculosis purified protein derivative (PPD RT49) (Serum Institute; Copenhagen, Denmark) (20  $\mu$ g/ml) was included as a positive control in all assays.

*Peptides:* Synthetic peptides were purchased from Research Genetics (Huntsville, Alabama) Mass-spectroscopy and HPLC verified peptides' authenticity and >70% purity. Digestion of gliadin (Sigma; G-3375) (100 mg/ml) with  $\alpha$ -chymotrypsin (Sigma; C-3142) 200:1 (w/w) was performed at room temperature in 0.1 M NH<sub>4</sub>HCO<sub>3</sub> with 2M urea and was halted after 24 h by heating to 98°C for 10 minutes. After centrifugation (13,000g, 10 minutes), the gliadin digest supernatant was filter-sterilized (0.2  $\mu$ m). Digestion of gliadin was verified by SDS-PAGE and protein concentration assessed.  $\alpha$ -Chymotrypsin-digested



- gliadin (640  $\mu\text{g/ml}$ ) and synthetic gliadin peptides (15-mers: 160  $\mu\text{g/ml}$ , other peptides: 0.1 mM) were individually treated with tTG (Sigma; T-5398) (50  $\mu\text{g/ml}$ ) in PBS +  $\text{CaCl}_2$  1 mM for 2 h at 37°C. Peptides and peptide pools were aliquotted into sterile 96-well plates and stored frozen at -20°C until use.
- Amino acid sequencing of peptides:* Reverse phase HPLC was used to purify the peptide resulting from tTG treatment of A-gliadin 56-75. A single product was identified and subjected to amino acid sequencing (automated sequencer Model 494A, Applied Biosystems, Foster City, California). The sequence of unmodified G56-75 was confirmed as: LQLQPFPPQLPYPQPQSFP (SEQ ID NO:5), and tTG treated G56-75 was identified as: LQLQPFPPQPELPYPQPQSFP (SEQ ID NO:11). Deamidation of glutamyl residues was defined as the amount (pmol) of glutamate recovered expressed as a percent of the combined amount of glutamine and glutamate recovered in cycles 2, 4, 8, 10, 15 and 17 of the amino acid sequencing. Deamidation attributable to tTG was defined as (% deamidation of glutamine in the tTG treated peptide - % deamidation in the untreated peptide) / (100 - % deamidation in the untreated peptide).
- CD4/CD8 and HLA Class II Restriction:* Anti-CD4 or anti-CD8 coated magnetic beads (Dynal, Oslo, Norway) were washed four times with RPMI then incubated with PBMC in complete RPMI containing 10% heat inactivated human AB serum (5x10<sup>6</sup> cells/ml) for 30 minutes on ice. Beads were removed using a magnet and cells remaining counted. In vivo HLA-class II restriction of the immune response to tTG-treated A-gliadin 56-75 was established by incubating PBMC (5x10<sup>6</sup> cells/ml) with anti-HLA-DR (L243), -DQ (L2), and -DP (B7.21) monoclonal antibodies (10  $\mu\text{g/ml}$ ) at room temperature for one hour prior to the addition of peptide.

#### Example 4

##### *Mucosal integrin expression by gliadin -specific peripheral blood lymphocytes*

Interaction between endothelial and lymphocyte adressins facilitates

homing

- 5 of organ-specific lymphocytes. Many adressesins are known. The heterodimer  $\alpha_4\beta_7$  is specific for lamina propria gut and other mucosal lymphocytes, and  $\alpha^E\beta_7$  is specific and intra-epithelial lymphocytes in the gut and skin. Approximately 30% of peripheral blood CD4 T cells express  $\alpha_4\beta_7$  and are presumed to be in transit to a mucosal site, while 5% of peripheral blood T cells express  $\alpha^E\beta_7$ .
- 10 Immunomagnetic beads coated with antibody specific for  $\alpha^E$  or  $\beta_7$  deplete PBMC of cells expressing  $\alpha^E\beta_7$  or  $\alpha^E\beta_7$  and  $\alpha_4\beta_7$ , respectively. In combination with ELISpot assay, immunomagnetic bead depletion allows determination of gliadin-specific T cell addressin expression that may identify these cells as homing to a mucosal surface. Interestingly, gluten challenge in vivo is associated with rapid
- 15 influx of CD4 T cells to the small intestinal lamina propria (not intra-epithelial sites), where over 90% lymphocytes express  $\alpha_4\beta_7$ .

Immunomagnetic beads were prepared and used to deplete PBMC from coeliac subjects on day 6 or 7 after commencing 3 day gluten challenge. FACS analysis demonstrated  $\alpha^E$  beads depleted approximately 50% of positive CD4 T

20 cells, while  $\beta_7$  beads depleted all  $\beta_7$  positive CD4 T cells. Depletion of PBMC using CD4- or  $\beta_7$ -beads, but not CD8- or  $\alpha^E$ -beads, abolished responses in the interferon gamma ELISpot. tTG gliadin and PPD responses were abolished by CD4 depletion, but consistently affected by integrin-specific bead depletion.

Thus A-gliadin 57-73 QE65-specific T cells induced after gluten

25 challenge in coeliac disease express the integrin,  $\alpha_4\beta_7$ , present on lamina propria CD4 T cells in the small intestine.

## Example 5

### *Optimal T cell Epitope Length*

- 30 Previous data testing peptides from 7 to 17 amino acids in length spanning the core of the dominant T cell epitope in A-gliadin indicated that the 17mer, A-gliadin 57-73 QE65 (SEQ ID NO:2) induced maximal responses in the interferon gamma Elispot using peripheral blood mononuclear cells (PBMC) from coeliac
- 35 volunteers 6 days after commencing a 3-day gluten challenge.

Peptides representing expansions form the core sequence of the dominant T cell epitope in A-gliadin were assessed in the IFN gamma ELISPOT using peripheral blood mononuclear cells (PBMC) from coeliac volunteers in 6 days after commencing a 3-day gluten challenge (n=4). Peptide 13: A-gliadin 59-71 QE65 (13mer), peptide 15: 58-72 QE65 (15mer), ..., peptide 27: 52-78 SE65 (27mer).

As shown in Figure 11 expansion of the A-gliadin 57-73 QE65 sequence does not substantially enhance response in the IFNgamma Elispot. Subsequent Examples characterise the agonist and antagonist activity of A-gliadin 57-73 QE65 using 17mer peptides.

### **Example 6**

*Comparison of A-gliadin 57-73 QE65 with other DQ2-restricted T cell epitopes in coeliac disease*

Dose response studies were performed using peptides corresponding to unmodified and transglutaminase-treated peptides corresponding to T cell epitopes of gluten-specific T cell clones and lines from intestinal biopsies of coeliac subjects. Responses to peptides were expressed as percent of response to A-gliadin 57-73 QE65. All subjects were HLA-DQ2+ (none were DQ8+).

The studies indicate that A-gliadin 57-73 QE65 is the most potent gliadin peptide for induction of interferon gamma in the ELISpot assay using coeliac PBMC after gluten challenge (see Figure 12a-h, and Tables 5 and 6). The second and third epitopes are suboptimal fragments of larger peptides i.e. A-gliadin 57-73 QE65 and GDA4\_WHEAT P04724-84-100 QE92. The epitope is only modestly bioactive (approximately 1/20<sup>th</sup> as active as A-gliadin 57-73 QE65 after blank is subtracted).

A-gliadin 57-73 QE65 is more potent than other known T cell epitopes in coeliac disease. There are 16 polymorphisms of A-gliadin 57-73 (including the sequence PQLPY (SEQ ID NO:12)) amongst sequenced gliadin genes, their bioactivity is assessed next.

## 5 **Example 7**

### *Comparison of gliadin- and A-gliadin 57-73 QE65-specific responses in peripheral blood*

The relative contribution of the dominant epitope, A-gliadin 57-73 QE65, to the total T cell response to gliadin in coeliac disease is a critical issue. Pepsin-  
 10 trypsin and chymotrypsin-digested gliadin have been traditionally used as antigen for development of T cell lines and clones in coeliac disease. However, it is possible that these proteases may cleave through certain peptide epitopes. Indeed, chymotrypsin digestion of recombinant  $\alpha$ 9-gliadin generates the peptide QLQPFQPELPY (SEQ ID NO:13), that is a truncation of the optimal epitope  
 15 sequence QLQPFQPELPYPQPQS (SEQ ID NO:2) (see above). Transglutaminase-treatment substantially increases the potency of chymotrypsin-digested gliadin in proliferation assays of gliadin-specific T cell clones and lines. Hence, transglutaminase-treated chymotrypsin-digested gliadin (tTG gliadin) may not be an ideal antigen, but responses against this mixture may approximate the  
 20 “total” number of peripheral blood lymphocyte specific for gliadin. Comparison of responses against A-gliadin 57-73 QE65 and tTG gliadin in the ELISpot assay gives an indication of the contribution of this dominant epitope to the overall immune response to gliadin in coeliac disease, and also be a measure of epitope spreading.

25 PBMC collected on day 6 or 7 after commencing gluten challenge in 4 coeliac subjects were assessed in dose response studies using chymotrypsin-digested gliadin +/- tTG treatment and compared with ELISpot responses to an optimal concentration of A-gliadin 57-73 QE65 (25mcg/ml). TTG treatment of gliadin enhanced PBMC responses in the ELISpot approximately 10-fold (tTG  
 30 was comparable to blank when assessed alone) (see Figure 13a-c). In the four coeliac subjects studied, A-gliadin 57-73 QE65 (25 mcg/ml) elicited responses between 14 and 115% those of tTG gliadin (500 mcg/ml), and the greater the response to A-gliadin 57-73 QE65 the greater proportion it represented of the tTG gliadin response.

5 Relatively limited data suggest that A-gliadin 57-73 QE65 responses are comparable to tTG gliadin in some subjects. Epitope spreading associated with more evolved anti-gliadin T cell responses may account for the smaller contribution of A-gliadin 57-73 QE65 to “total” gliadin responses in peripheral blood in some individuals. Epitope spreading may be maintained in individuals  
10 with less strictly gluten free diets.

### Example 8

*Definition of gliadin peptides bioactive in coeliac disease: polymorphisms of A-gliadin 57-73*

15 Overlapping 15mer peptides spanning the complete sequence of A-gliadin were assessed in order to identify the immunodominant sequence in coeliac disease. A-gliadin was the first fully sequenced alpha gliadin protein and gene, but is one of approximately 30-50 related alpha gliadin proteins in wheat. Twenty five distinct alpha-gliadin genes have been identified by searching protein  
20 data bases, Swiss-Prot and TREMBL describing a further 8 alpha-gliadins. Contained within these 25 alpha-gliadins, there are 16 distinct polymorphisms of the sequence corresponding to A-gliadin 57-73 (see Table 7).

Synthetic peptides corresponding to these 16 polymorphisms, in an unmodified form, after treatment with transglutaminase in vitro, as well as with  
25 glutamate substituted at position 10 (equivalent to QE65 in A-gliadin 57-73) were assessed using PBMC from coeliac subjects, normally following a gluten free diet, day 6 or 7 after gluten challenge in interferon gamma ELISpot assays. Glutamate-substituted peptides were compared at three concentrations (2.5, 25 and 250 mcg/ml), unmodified peptide and transglutaminase-treated peptides were  
30 assessed at 25 mcg/ml only. Bioactivity was expressed as % of response associated with A-gliadin 57-73 QE65 25 mcg/ml in individual subjects (n=4). (See Fig 14).

Bioactivity of “wild-type” peptides was substantially increased (>5-fold) by treatment with transglutaminase. Transglutaminase treatment of wild-type  
35 peptides resulted in bioactivity similar to that of the same peptides substituted

5 with glutamate at position 10. Bioactivities of five glutamate-substituted peptides (B, C, K, L, M), were >70% that of A-gliadin 57-73 QE65 (A), but none was significantly more bioactive than A-gliadin 57-73 QE65. PBMC responses to glutamate-substituted peptides at concentrations of 2.5 and 250 mcg/ml were comparable to those at 25 mcg/ml. Six glutamate-substituted gliadin peptides (H,  
10 I, J, N, O, P) were <15% as bioactive as A-gliadin 57-73 QE65. Other peptides were intermediate in bioactivity.

At least six gliadin-derived peptides are equivalent in potency to A-gliadin 57-73 QE65 after modification by transglutaminase. Relatively non-bioactive polymorphisms of A-gliadin 57-73 also exist. These data indicate that  
15 transglutaminase modification of peptides from several gliadins of *Triticum aestivum*, *T. uarzu* and *T. spelta* may be capable of generating the immunodominant T cell epitope in coeliac disease.

Genetic modification of wheat to generate non-coeliac-toxic wheat may likely require removal or modification of multiple gliadin genes. Generation of  
20 wheat containing gliadins or other proteins or peptides incorporating sequences defining altered peptide ligand antagonists of A-gliadin 57-73 is an alternative strategy to generate genetically modified wheat that is therapeutic rather than “non-toxic” in coeliac disease.

## 25 **Example 9**

### *Definition of Core Epitope Sequence:*

Comparison of peptides corresponding to truncations of A-gliadin 56-75 from the N- and C-terminal indicated that the core sequence of the T cell epitope is PELPY (A-gliadin 64-68). Attempts to define non-agonists and antagonists  
30 will focus on variants of A-gliadin that are substituted at residues that substantially contribute to its bioactivity.

Peptides corresponding to A-gliadin 57-73 QE65 with alanine (Figure 15) or lysine (Figure 16) substituted for residues 57 to 73 were compared in the IFN gamma ELISPOT using peripheral blood mononuclear cells (PBMC) from

5 coeliac volunteers 6 days after commencing a 3-day gluten challenge (n=8). (BL is blank, E is A-gliadin 57-73 QE65: QLQFPQPPELPYPQPQS (SEQ ID NO:2)).

It was found that residues corresponding to A-gliadin 60-70 QE65 (PFPQPPELPYPQ (SEQ ID NO:14)) contribute substantially to the bioactivity in A-gliadin 57-73 QE65. Variants of A-gliadin 57-73 QE65 substituted at positions 60-70 are assessed in a 2-step procedure. Initially, A-gliadin 57-73 QE65 substituted at positions 60-70 using 10 different amino acids with contrasting properties are assessed. A second group of A-gliadin 57-73 QE65 variants (substituted with all other naturally occurring amino acids except cysteine at positions that prove are sensitive to modification) are assessed in a second round.

### Example 10

#### *Agonist activity of substituted variants of A-gliadin 57-73 QE65*

A-gliadin 60-70 QE65 is the core sequence of the dominant T cell epitope in A-gliadin. Antagonist and non-agonist peptide variants of this epitope are most likely generated by modification of this core sequence. Initially, A-gliadin 57-73 QE65 substituted at positions 60-70 using 10 different amino acids with contrasting properties will be assessed in the IFN $\gamma$  ELISPOT using PBMC from coeliac subjects 6 days after starting 3 day gluten challenge. A second group of A-gliadin 57-73 QE65 variants (substituted with all other naturally occurring amino acids except cysteine) at positions 61-70 were also assessed. Both groups of peptides (all at 50 mcg/ml, in duplicate) were assessed using PBMC from 8 subjects and compared to the unmodified peptide (20 replicates per assay). Previous studies indicate that the optimal concentration for A-gliadin 57-73 QE65 in this assay is between 10 and 100 mcg/ml.

Results are expressed as mean response in spot forming cells (95% confidence interval) as % A-G 57-73 QE65 mean response in each individual. Unpaired t-tests will be used to compare ELISPOT responses of modified peptides with A-G 57-73 QE65. Super-agonists were defined as having a greater response than A-G 57-73 QE65 at a level of significance of  $p < 0.01$ ; partial

agonists as having a response less than A-G 57-73 QE65 at a level of significance of  $p < 0.01$ , and non-agonists as being not significantly different ( $p > 0.01$ ) from blank (buffer without peptide). Peptides with agonist activity 30% or less that of A-gliadin 57-73 QE65 were considered “suitable” partial or non-agonists to assess for antagonistic activity (see Table 8 and Figures 17-27).

The IFN $\gamma$  ELISPOT response of PBMC to A-gliadin 57-73 QE65 is highly specific at a molecular level. Proline at position 64 (P64), glutamate at 65 (E65) and leucine at position 66 (L66), and to a lesser extent Q63, P67, Y68 and P69 are particularly sensitive to modification. The substitutions Y61 and Y70 both generate super-agonists with 30% greater bioactivity than the parent peptide, probably by enhancing binding to HLA-DQ2 since the motif for this HLA molecule indicates a preference for bulky hydrophobic residues at positions 1 and 9. Eighteen non-agonist peptides were identified. Bioactivities of the variants (50 mcg/ml): P65, K64, K65 and Y65 (bioactivity 7-8%) were comparable to blank (7%). In total, 57 mutated variants of A-gliadin 57-73 QE65 were 30% or less bioactive than A-gliadin 57-73 QE65.

The molecular specificity of the peripheral blood lymphocyte (PBL) T cell response to the dominant epitope, A-gliadin 57-73 QE65, is consistently reproducible amongst HLA-DQ2+ coeliac subjects, and is highly specific to a restricted number of amino acids in the core 7 amino acids. Certain single-amino acid variants of A-gliadin 57-73 QE65 are consistently non-agonists in all HLA-DQ2+ coeliac subjects.

### Example 11

#### *Antagonist activity of substituted variants*

The homogeneity of the PBL T cell response to A-gliadin 57-73 QE65 in HLA-DQ2+ coeliac disease suggests that altered peptide ligands (APL) capable of antagonism in PBMC ex vivo may exist, even though the PBL T cell response is likely to be poly- or oligo-clonal. APL antagonists are generally weak agonists. Fifty-seven single amino acid-substituted variants of A-gliadin 57-73 QE65 with agonist activity 30% or less have been identified and are suitable



5 candidates as APL antagonists. In addition, certain weakly bioactive naturally occurring polymorphisms of A-gliadin 57-73 QE65 have also been identified (see below) and may be “naturally occurring” APL antagonists. It has also been suggested that competition for binding MHC may also antagonise antigen-specific T cell immune. Hence, non-gliadin peptides that do not induce  
 10 IFN $\gamma$  responses in coeliac PBMC after gluten challenge but are known to bind to HLA-DQ2 may be capable of reducing T cell responses elicited by A-gliadin 57-73 QE65. Two peptides that bind avidly to HLA-DQ2 are HLA class 1  $\alpha$  46-60 (HLA 1a) (PRAPWIEQEGPEYW (SEQ ID NO:15)) and thyroid peroxidase (tp) 632-645Y (IDVWLGGLLAENFLPY (SEQ ID NO:16)).

15 Simultaneous addition of peptide (50 $\mu$ g/ml) or buffer and A-gliadin 57-73 QE65 (10 $\mu$ g/ml) in IFN $\gamma$  ELISPOT using PBMC from coeliac volunteers 6 days after commencing 3 day gluten challenge (n=5). Results were expressed as response with peptide plus A-G 57-73 QE65 (mean of duplicates) as % response with buffer plus A-G 57-73 QE65 (mean of 20 replicates). (See Table 9).

20 Four single amino acid-substituted variants of A-gliadin 57-73 QE65 reduce the interferon gamma PBMC ELISPOT response to A-gliadin 57-73 QE65 ( $p < 0.01$ ) by between 25% and 28%, 13 other peptide variants reduce the ELISPOT response by between 18% and 24% ( $p < 0.06$ ). The HLA-DQ2 binder, thyroid peroxidase (tp) 632-645Y reduces PBMC interferon gamma responses to  
 25 A-gliadin 57-73 QE65 by 31% ( $p < 0.0001$ ) but the other HLA-DQ2 binder, HLA class 1  $\alpha$  46-60, does not alter responses (see Tables 9 and 10). The peptide corresponding to a transglutaminase-modified polymorphism of A-gliadin 57-73, SwissProt accession no.: P04725 82-98 QE90 (PQPQFPPELPYPQPQS (SEQ ID NO:17)) reduces responses to A-gliadin 57-73 QE65 by 19% ( $p < 0.009$ ) (see  
 30 Table 11).

Interferon gamma responses of PBMC to A-gliadin 57-73 QE65 in ELISPOT assays are reduced by co-administration of certain single-amino acid A-gliadin 57-73 QE65 variants, a polymorphism of A-gliadin 57-73 QE65, and an unrelated peptide known to bind HLA-DQ2 in five-fold excess. These finding  
 35 suggest that altered peptide ligand antagonists of A-gliadin 57-73 QE65 exist.

- 5 Not only putative APL antagonists but also certain peptides that bind HLA-DQ2 effectively reduce PBL T cell responses to A-gliadin 57-73 QE65.

These findings support two strategies to interrupt the T cell response to the dominant A-gliadin epitope in HLA-DQ2+ coeliac disease.

1. Optimisation of APL antagonists by substituting amino acids at more than  
10 one position (64-67) for use as “traditional” peptide pharmaceuticals or for specific genetic modification of gliadin genes in wheat.
2. Use of high affinity HLA-DQ2 binding peptides to competitively inhibit presentation of A-gliadin 57-73 QE65 in association with HLA-DQ2.

- 15 These two approaches may be mutually compatible. Super-agonists were generated by replacing F61 and Q70 with tyrosine residues. It is likely these super-agonists resulted from improved binding to HLA-DQ2 rather than enhanced contact with the T cell receptor. By combining these modifications with other substitutions that generate modestly effective APL antagonists might  
20 substantially enhance the inhibitory effect of substituted A-gliadin 57-73 QE65 variants.

## Example 12

- Development of interferon gamma ELISpot using PBMC and A-gliadin 57-73  
25 QE65 and P04724 84-100 QE92 as a diagnostic for coeliac disease: Definition of immune-responsiveness in newly diagnosed coeliac disease*

- Induction of responsiveness to the dominant A-gliadin T cell epitope in PBMC measured in the interferon gamma ELISpot follows gluten challenge in almost all DQ2+ coeliac subjects following a long term strict gluten free diet  
30 (GFD) but not in healthy DQ2+ subjects after 4 weeks following a strict GFD. A-gliadin 57-73 QE65 responses are not measurable in PBMC of coeliac subjects before gluten challenge and pilot data have suggested these responses could not be measured in PBMC of untreated coeliacs. These data suggest that in coeliac disease immune-responsiveness to A-gliadin 57-73 QE65 is restored following  
35 antigen exclusion (GFD). If a diagnostic test is to be developed using the

ELISpot assay and PBMC, it is desirable to define the duration of GFD required before gluten challenge is capable of inducing responses to A-gliadin 57-73 QE65 and other immunoreactive gliadin peptides in blood.

Newly diagnosed DQ2+ coeliac subjects were recruited from the gastroenterology outpatient service. PBMC were prepared and tested in interferon gamma ELISpot assays before subjects commenced GFD, and at one or two weeks after commencing GFD. In addition, gluten challenge (3 days consuming 4 slices standard white bread, 200g/day) was performed at one or two weeks after starting GFD. PBMC were prepared and assayed on day six after commencing gluten challenge. A-gliadin 57-73 QE65 (A), P04724 84-100 QE92 (B) (alone and combined) and A-gliadin 57-73 QP65 (P65) (non-bioactive variant, see above) (all 25 mcg/ml) were assessed.

All but one newly diagnosed coeliac patient was DQ2+ (one was DQ8+) (n=11). PBMC from newly diagnosed coeliacs that were untreated, or after 1 or 2 weeks following GFD did not show responses to A-gliadin 57-73 QE65 and P04724 84-100 QE92 (alone or combined) that were not significantly different from blank or A-gliadin 57-73 QP65 (n=9) (see Figure 28). Gluten challenge in coeliacs who had followed GFD for only one week did not substantially enhance responses to A-gliadin 57-73 QE65 or P04724 84-100 QE92 (alone or combined). But gluten challenge 2 weeks after commencing GFD did induce responses to A-gliadin 57-73 QE65 and P04724 84-100 QE92 (alone or combined) that were significantly greater than the non-bioactive variant A-gliadin 57-73 QP65 and blank. Although these responses after gluten challenge at 2 weeks were substantial they appear to be less than in subjects >2 months after commencing GFD. Responses to A-gliadin 57-73 QE65 alone were equivalent or greater than responses to P04724 84-100 QE92 alone or when mixed with A-gliadin 57-73 QE65. None of the subjects experienced troubling symptoms with gluten challenge.

Immune responsiveness (as measured in PBMC after gluten challenge) to A-gliadin is partially restored 2 weeks after commencing GFD, implying that "immune unresponsiveness" to this dominant T cell epitope prevails in untreated

coeliac disease and for at least one week after starting GFD. The optimal timing of a diagnostic test for coeliac disease using gluten challenge and measurement of responses to A-gliadin 57-73 QE65 in the ELISpot assay is at least 2 weeks after commencing a GFD.

Interferon gamma-secreting T cells specific to A-gliadin 57-73 QE65 cannot be measured in the peripheral blood in untreated coeliacs, and can only be induced by gluten challenge after at least 2 weeks GFD (antigen exclusion). Therefore, timing of a diagnostic test using this methodology is crucial and further studies are needed for its optimization. These finding are consistent with functional anergy of T cells specific for the dominant epitope, A-gliadin 57-73 QE65, reversed by antigen exclusion (GFD). This phenomenon has not been previously demonstrated in a human disease, and supports the possibility that T cell anergy may be inducible with peptide therapy in coeliac disease.

### Example 13

#### *Comprehensive Mapping of Wheat Gliadin T Cell Epitopes*

Antigen challenge induces antigen-specific T cells in peripheral blood. In coeliac disease, gluten is the antigen that maintains this immune-mediated disease. Gluten challenge in coeliac disease being treated with a gluten free diet leads to the appearance of gluten-specific T cells in peripheral blood, so enabling determination of the molecular specificity of gluten T cell epitopes. As described above, we have identified a single dominant T cell epitope in a model gluten protein, A-gliadin (57-73 deamidated at Q65). In this Example, gluten challenge in coeliac patients was used to test all potential 12 amino acid sequences in every known wheat gliadin protein derived from 111 entries in Genbank. In total, 652 20mer peptides were tested in HLA-DQ2 and HLA-DQ8 associated coeliac disease. Seven of the 9 coeliac subjects with the classical HLA-DQ2 complex (HLA-DQA1\*05, HLA-DQB1\*02) present in over 90% of coeliacs had an inducible A-gliadin 57-73 QE65- and gliadin-specific T cell response in peripheral blood. A-gliadin 57-73 was the only significant  $\alpha$ -gliadin T cell epitope, as well as the most potent gliadin T cell epitope, in HLA-DQ2-associated

5 coeliac disease. In addition, there were as many as 5 families of structurally related peptides that were between 10 and 70% as potent as A-gliadin 57-73 in the interferon- $\gamma$  ELISpot assay. These new T cell epitopes were derived from  $\gamma$ - and  $\omega$ -gliadins and included common sequences that were structurally very similar, but not identical to the core sequence of A-gliadin 57-73 (core sequence: 10 FPQPQLPYP (SEQ ID NO:18)), for example: FPQPQQPFP (SEQ ID NO:19) and PQQPQQPFP (SEQ ID NO:20). Although no homologues of A-gliadin 57-73 have been found in rye or barley, the other two cereals toxic in coeliac disease, the newly defined T cell epitopes in  $\gamma$ - and  $\omega$ -gliadins have exact matches in rye and barley storage proteins (secalins and hordeins, respectively).

15 Coeliac disease not associated with HLA-DQ2 is almost always associated with HLA-DQ8. None of the seven HLA-DQ8+ coeliac subjects had inducible A-gliadin 57-73-specific T cell responses following gluten challenge, unless they also possessed the complete HLA-DQ2 complex. Two of 4 HLA-DQ8+ coeliac subjects who did not possess the complete HLA-DQ2 complex, 20 had inducible gliadin peptide-specific T cell responses following gluten challenge. In one HLA-DQ8 subject, a novel dominant T cell epitope was identified with the core sequence LQPQNPSQQPQ (SEQ ID NO:21). The transglutaminase-deamidated version of this peptide was more potent than the non-deamidated peptide. Previous studies suggest that the transglutaminase-deamidated peptide would have the sequence LQPENPSQEQPE (SEQ ID 25 NO:22); but further studies are required to confirm this sequence. Amongst the healthy HLA-DQ2 (10) and HLA-DQ8 (1) subjects who followed a gluten free diet for a month, gliadin peptide-specific T cell responses were uncommon, seldom changed with gluten challenge, and were never potent T cell epitopes revealed with gluten challenge in coeliac subjects. In conclusion, there are 30 unlikely to be more than six important T cell epitopes in HLA-DQ2-associated coeliac disease, of which A-gliadin 57-73 is the most potent. HLA-DQ2- and HLA-DQ8-associated coeliac disease do not share the same T cell specificity.

We have shown that short-term gluten challenge of individuals with 35 coeliac disease following a gluten free diet induces gliadin-specific T cells in

5 peripheral blood. The frequency of these T cells is maximal in peripheral blood on day 6 and then rapidly wanes over the following week. Peripheral blood gliadin-specific T cells express the integrin  $\alpha 4\beta 7$  that is associated with homing to the gut lamina propria. We exploited this human antigen-challenge design to map T cell epitopes relevant to coeliac disease in the archetypal gluten  $\alpha$ -gliadin protein, A-gliadin. Using 15mer peptides overlapping by 10 amino acids with  
 10 and without deamidation by transglutaminase (tTG), we demonstrated that T cells induced in peripheral blood initially target only one A-gliadin peptide, residues 57-73 in which glutamine at position 65 is deamidated. The epitope is HLA-DQ2-restricted, consistent with the intimate association of coeliac disease with  
 15 HLA-DQ2.

Coeliac disease is reactivated by wheat, rye and barley exposure. The  $\alpha/\beta$ -gliadin fraction of wheat gluten is consistently toxic in coeliac disease, and most studies have focused on these proteins. The gene cluster coding for  $\alpha/\beta$ -gliadins is located on wheat chromosome 6C. There are no homologues of  $\alpha/\beta$ -gliadins in rye or barley. However, all three of the wheat gliadin subtypes ( $\alpha/\beta$ ,  $\gamma$ ,  
 20 and  $\omega$ ) are toxic in coeliac disease. The  $\gamma$ - and  $\omega$ -gliadin genes are located on chromosome 1A in wheat, and are homologous to the secalins and hordeins in rye and barley.

There are now genes identified for 61  $\alpha$ -gliadins in wheat (*Triticum aestivum*). The  $\alpha$ -gliadin sequences are closely homologous, but the dominant epitope in A-gliadin derives from the most polymorphic region in the  $\alpha$ -gliadin sequence. Anderson et al (1997) have estimated that there are a total of about 150 distinct  $\alpha$ -gliadin genes in *T. aestivum*, but many are pseudogenes. Hence, it is unlikely that T-cell epitopes relevant to coeliac disease are not included within  
 30 known  $\alpha$ -gliadin sequences.

Our work has identified a group of deamidated  $\alpha$ -gliadin peptides almost identical to A-gliadin 57-73 as potent T cell epitopes specific to coeliac disease. Over 90% of coeliac patients are HLA-DQ2+, and so far, we have only assessed HLA-DQ2+ coeliac subjects after gluten challenge. However, coeliac patients  
 35 who do not express HLA-DQ2 nearly all carry HLA-DQ8. Hence, it is critical to

know whether A-gliadin 57-73 and its homologues in other wheat, rye and barley gluten proteins are the only T-cell epitopes recognized by T cells induced by gluten challenge in both HLA-DQ2+ and HLA-DQ8+ coeliac disease. If this were the case, design of peptide therapeutics for coeliac disease might only require one peptide.

#### *Homologues of A-gliadin 57-73 as T-cell epitopes*

Initial searches of SwissProt and Trembl gene databases for cereal genes coding for the core sequence of A-gliadin 57-73 (PQLPY <SEQ ID NO:12>) only revealed  $\alpha/\beta$ -gliadins. However, our fine-mapping studies of the A-gliadin 57-73 QE65 epitope revealed a limited number of permissive point substitutions in the core region (PQLP) (note Q65 is actually deamidated in the epitope). Hence, we extended our search to genes in SwissProt or Trembl databases encoding for peptides with the sequence XXXXXXXPQ[ILMP][PST]XXXXXX (SEQ ID NO:23). Homologues were identified amongst  $\gamma$ -gliadins, glutenins, hordeins and secalins (see Table 12). A further homologue was identified in  $\omega$ -gliadin by visual search of the three  $\omega$ -gliadin entries in Genbank.

These homologues of A-gliadin 57-73 were assessed after deamidation by tTG (or synthesis of the glutamate(QE)-substituted variant in four close homologues) using the IFN $\gamma$  ELISpot assay with peripheral blood mononuclear cells after gluten challenge in coeliac subjects. The  $\omega$ -gliadin sequence (AAG17702 141-157) was the only bioactive peptide, approximately half as potent as A-gliadin 57-73 (see Table 12, and Figure 29). Hence, searches for homologues of the dominant A-gliadin epitope failed to account for the toxicity of  $\gamma$ -gliadin, secalins, and hordeins.

#### *Methods*

##### *Design of a set of peptides spanning all possible wheat gliadin T-cell epitopes*

In order to identify all possible T cell epitopes coded by the known wheat (*Triticum aestivum*) gliadin genes or gene fragments (61  $\alpha/\beta$ -, 47  $\gamma$ -, and 3  $\omega$ -gliadin entries in Genbank), gene-derived protein sequences were aligned using the ClustalW software (MegAlign) and arranged into phylogenetic groupings (see

5 Table 22). Many entries represented truncations of longer sequences, and many gene segments were identical except for the length of polyglutamine repeats or rare substitutions. Hence, it was possible to rationalize all potential unique 12 amino acid sequences encoded by known wheat genes to be included in a set of 652 20mer peptides. (Signal peptide sequences were not included). Peptide  
10 sequences are listed in Table 23.

*Comprehensive epitope mapping*

Healthy controls (HLA-DQ2+ n=10, and HLA-DQ8+ n=1) who had followed a gluten free diet for 4 weeks, and coeliac subjects (six HLA-DQ2, four complex heterozygotes HLA-DQ2/8, and three HLA-DQ8/X) (see Table 13)  
15 following long-term gluten free diet were studied before and on day 6 and 7 after 3-day gluten challenge (four 50g slices of standard white bread – Sainsbury's sandwich bread, each day). Peripheral blood (a total of 300ml over seven days) was collected and peripheral blood mononuclear cells (PBMC) were separated by Lymphoprep density gradient. PBMC were incubated with pools of 6 or 8 20mer  
20 peptides, or single peptides with or without deamidation by tTG in overnight interferon gamma (IFN $\gamma$ ) ELISpot assays.

Peptides were synthesized in batches of 96 as Pepsets (Mimotopes Inc., Melbourne Australia). Approximately 0.6 micromole of each of 652 20mers was provided. Two marker 20mer peptides were included in each set of 96  
25 (VLQQHNIAHGSSQVLQESTY – peptide 161 (SEQ ID NO:24), and IKDFHVYFRESRDALWKGPG (SEQ ID NO:25)) and were characterized by reverse phase-HPLC and amino acid sequence analysis. Average purities of these marker peptides were 50% and 19%, respectively. Peptides were initially dissolved in acetonitrile (10%) and Hepes 100mM to 10mg/ml.

30 The final concentration of individual peptides in pools (or alone) incubated with PBMC for the IFN $\gamma$  ELISpot assays was 20  $\mu$ g/ml. Five-times concentrated solutions of peptides and pools in PBS with calcium chloride 1mM were aliquotted and stored in 96-well plates according to the template later used in ELISpot assays. Deamidated peptides and pools of peptides were prepared by  
35 incubation with guinea pig tissue tTG (Sigma T5398) in the ratio 100:32  $\mu$ g/ml



5 for two hours at 37°C. Peptides solutions were stored at -20°C and freshly thawed prior to use.

Gliadin (Sigma G3375) (100 mg/ml) in endotoxin-free water and 2M urea was boiled for 10 minutes, cooled to room temperature and incubated with filter (0.2 µm)-sterilised pepsin (Sigma P6887) (2 mg/ml) in HCl 0.02M or  
 10 chymotrypsin (C3142) (4mg/ml) in ammonium bicarbonate (0.2M). After incubation for 4 hours, pepsin-digested gliadin was neutralized with sodium hydroxide, and then both pepsin- and chymotrypsin-digested gliadin were boiled for 15 minutes. Identical incubations with protease in which gliadin was omitted were also performed. Samples were centrifuged at 15 000g, then protein  
 15 concentrations were estimated in supernatants by the BCA method (Pierce, USA). Before final use in IFNγ ELISpot assays, aliquots of gliadin-protease were incubated with tTG in the ratio 2500:64 µg/ml.

IFNγ ELISpot assays (Mabtech, Sweden) were performed in 96-well plates (MAIP S-45, Millipore) in which each well contained 25µl of peptide  
 20 solution and 100µl of PBMC ( $2 \times 10^5$ /well) in RPMI containing 10% heat inactivated human AB serum. Deamidated peptide pools were assessed in one 96-well ELISpot plate, and peptides pools without deamidation in a second plate (with an identical layout) on both day 0 and day 6. All wells in the plate containing deamidated peptides included tTG (64 µg/ml). In each ELISpot plate  
 25 there were 83 wells with peptide pools (one unique pool in each well), and a series of wells for “control” peptides (peptides all >90% purity, characterized by MS and HPLC, Research Genetics): P04722 77-93 (QLQPFQPQLPYQPQP (SEQ ID NO:26)), P04722 77-93 QE85 (in duplicate) (QLQPFQPQLPYQPQP (SEQ ID NO:27)), P02863 77-93 (QLQPFQPQLPYSQPQP (SEQ ID NO:28)),  
 30 P02863 77-93 QE85 (QLQPFQPQLPYSQPQP (SEQ ID NO:29)), and chymotrypsin-digested gliadin (500 µg/ml), pepsin-digested gliadin (500 µg/ml), chymotrypsin (20 µg/ml) alone, pepsin (10 µg/ml) alone, and blank (PBS+/-tTG) (in triplicate).

After development and drying, IFNγ ELISpot plates were assessed using  
 35 the MAIP automated ELISpot plate counter. In HLA-DQ2 healthy and coeliac

5 subjects, induction of spot forming cells (sfc) by peptide pools in the IFN $\gamma$  ELISpot assay was tested using a one-tailed Wilcoxon Matched-Pairs Signed-Ranks test (using SPSS software) applied to spot forming cells (sfc) per million PBMC minus blank on day 6 versus day 0 (“net response”). Significant induction of an IFN $\gamma$  response to peptide pools in PBMC by *in vivo* gluten challenge was  
 10 defined as a median “net response” of at least 10 sfc/million PBMC and  $p < 0.05$  level of significance. Significant response to a particular pool of peptides on day 6 was followed by assessment of individual peptides within each pool using PBMC drawn the same day or on day 7.

For IFN $\gamma$  ELISpot assays of individual peptides, bioactivity was  
 15 expressed as a percent of response to P04722 77-93 QE85 assessed in the same ELISpot plate. Median response to blank (PBS alone) was 0.2 (range 0-5) sfc per well, and the positive control (P04722 77-93 QE85) 76.5 (range: 25-282) sfc per well using a median of 0.36 million (range: 0.3-0.72) PBMC. Hence, median response to blank expressed as a percentage of P04722 77-93 QE65 was 0.2%  
 20 (range: 0-6.7). Individual peptides with mean bioactivity greater than 10% that of P04722 QE85 were analyzed for common structural motifs.

## Results

### *Healthy HLA-DQ2 subjects*

None of the healthy HLA-DQ2+ subjects following a gluten free diet for a  
 25 month had IFN $\gamma$  ELISpot responses to homologues of A-gliadin 57-73 before or after gluten challenge. However, in 9/10 healthy subjects, gluten challenge was associated with a significant increase in IFN $\gamma$  responses to both peptic- and chymotryptic-digests of gliadin, from a median of 0-4 sfc/million on day 0 to a median of 16-29 sfc/million (see Table 14). Gliadin responses in healthy subjects  
 30 were unaffected by deamidation (see Table 15). Amongst healthy subjects, there was no consistent induction of IFN $\gamma$  responses to specific gliadin peptide pools with gluten challenge (see Figure 30, and Table 16). IFN $\gamma$  ELISpot responses were occasionally found, but these were weak, and not altered by deamidation. Many of the strongest responses to pools were also present on day 0 (see Table  
 35 17, subjects H2, H8 and H9). Four healthy subjects did show definite responses

to pool 50, and the two with strongest responses on day 6 also had responses on day 0. In both subjects, the post-challenge responses to pool 50 responses were due to peptide 390 (QQTYPQRPQQPFQTQQPQQ (SEQ ID NO:30)).

#### *HLA-DQ2 coeliac subjects*

Following gluten challenge in HLA-DQ2+ coeliac subjects, median IFN $\gamma$  ELISpot responses to P04722 77-93 E85 rose from a median of 0 to 133 sfc/million (see Table 4). One of the six coeliac subjects (C06) did not respond to P04722 77-93 QE85 (2 sfc/million) and had only weak responses to gliadin peptide pools (maximum: Pool 50+tTG 27 sfc/million). Consistent with earlier work, bioactivity of wild-type P04722 increased 6.5 times with deamidation by tTG (see Table 15). Interferon-gamma responses to gliadin-digests were present at baseline, but were substantially increased by gluten challenge from a median of 20 up to 92 sfc/million for chymotryptic-gliadin, and from 44 up to 176 sfc/million for peptide-gliadin. Deamidation of gliadin increased bioactivity by a median of 3.2 times for chymotryptic-gliadin and 1.9 times for peptic-gliadin (see Table 15). (Note that the acidity required for digestion by pepsin is likely to result in partial deamidation of gliadin.)

In contrast to healthy subjects, gluten challenge induced IFN $\gamma$  ELISpot responses to 22 of the 83 tTG-treated pools including peptides from  $\alpha$ -,  $\gamma$ - and  $\omega$ -gliadins (see Figure 31, and Table 17). Bioactivity of pools was highly consistent between subjects (see Table 18). IFN $\gamma$  ELISpot responses elicited by peptide pools were almost always increased by deamidation (see Table 17). But enhancement of bioactivity of pools by deamidation was not as marked as for P04722 77-73 Q85, even for pools including homologues of A-gliadin 57-73. This suggests that Pepset peptides were partially deamidated during synthesis or in preparation, for example the Pepset peptides are delivered as salts of trifluoroacetic acid (TFA) after lyophilisation from a TFA solution.

One hundred and seventy individual tTG-deamidated peptides from 21 of the most bioactive pools were separately assessed. Seventy-two deamidated peptides were greater than 10% as bioactive as P04722 77-93 QE85 at an equivalent concentration (20  $\mu$ g/ml) (see Table 19). The five most potent

5 peptides (85-94% bioactivity of P04722 QE85) were previously identified  $\alpha$ -gliadin homologues A-gliadin 57-73. Fifty of the bioactive peptides were not homologues of A-gliadin 57-73, but could be divided into six families of structurally related sequences (see Table 20). The most bioactive sequence of each of the peptide families were: PQQPQQPQQPFPQPPQFPW (SEQ ID NO:31) (peptide 626, median 72% bioactivity of P04722 QE85),  
 10 QQPQQPFPQQPQQQLPFPQQ (SEQ ID NO:32) (343, 34%),  
QAFPPQQQTFFPHQPPQQQFPQ (SEQ ID NO:33) (355, 27%),  
TQQPQQPFPQQPQQPFPQTQ (SEQ ID NO:34) (396, 23%),  
PIQPPQQPFPQQPQQPQQPFP (SEQ ID NO:35) (625, 22%),  
 15 PQQSFSYQQQFPQQPYPQQ (SEQ ID NO:36) (618, 18%) (core sequences are underlined). All of these sequences include glutamine residues predicted to be susceptible to deamidation by transglutaminase (e.g. QXP, QXPF (SEQ ID NO:37), QXX[FY] (SEQ ID NO:38)) (see Vader et al 2002). Some bioactive peptides contain two core sequences from different families.

20 Consistent with the possibility that different T-cell populations respond to peptides with distinct core sequences, bioactivity of peptides from different families appear to be additive. For example, median bioactivity of tTG-treated Pool 81 was 141% of P04722 QE85, while bioactivity of individual peptides was in rank order: Peptide 631 (homologue of A-gliadin 57-73) 61%, 636 (homologue  
 25 of 626) 51%, and 635 19%, 629 16%, and 634 13% (all homologues of 396).

Although likely to be an oversimplification, the contribution of each “peptide family” to the summed IFN $\gamma$  ELISpot response to gliadin peptides was compared in the HLA-DQ2+ coeliac subjects (see Figure 32). Accordingly, the contribution of P04722 77-73 E85 to the summed response to gliadin peptides is  
 30 between 1/5 and 2/3.

Using the peptide homology search programme, WWW PepPepSearch, which can be accessed through the world wide web of the internet at, for example, “cbrg.inf.ethz.ch/subsection3\_1\_5.html.”, and by direct comparison with Genbank sequences for rye secalins, exact matches were found for the core  
 35 sequences QQPFPQPQQPFP (SEQ ID NO:39) in barley hordeins (HOR8) and

rye secalins (A23277, CAA26449, AAG35598), QQPFPQQPQQPFP (SEQ ID NO:40) in barley hordeins (HOG1 and HOR8), and for PIQPQQPFPQQP (SEQ ID NO:41) also in barley hordeins (HOR8).

#### *HLA-DQ8-associated coeliac disease*

Seven HLA-DQ8+ coeliac subjects were studied before and after gluten challenge. Five of these HLA-DQ8+ (HLA-DQA0\*0301-3, HLA-DQB0\*0302) subjects also carried one or both of the coeliac disease-associated HLA-DQ2 complex (DQA0\*05, DQB0\*02). Two of the three subjects with both coeliac-associated HLA-DQ complexes had potent responses to gliadin peptide pools (and individual peptides including P04722 77-93 E85) that were qualitatively and quantitatively identical to HLA-DQ2 coeliac subjects (see Figures 33 and 34, and Table 18). Deamidated peptide pool 74 was bioactive in both HLA-DQ2/8 subjects, but only in one of the 6 HLA-DQ2/X subjects. Pretreatment of pool 74 with tTG enhances bioactivity between 3.8 and 22-times, and bioactivity of tTG-treated pool 74 in the three responders is equivalent to between 78% and 350% the bioactivity of P04722 77-93 E85. Currently, it is not known which peptides are bioactive in Pool 74 in subject C02, C07, and C08.

Two of the four HLA-DQ8 coeliac subjects that lacked both or one of the HLA-DQ2 alleles associated with coeliac disease showed very weak IFN $\gamma$  ELISpot responses to gliadin peptide pools, but the other two did respond to both protease-digested gliadin and specific peptide pools. Subject C12 (HLA-DQ7/8) responded vigorously to deamidated Pools 1-3 (see Figure 35). Assessment of individual peptides in these pools identified a series of closely related bioactive peptides including the core sequence LQPQNPSQQQPQ (SEQ ID NO:42) (see Table 20). Previous work (by us) has demonstrated that three glutamine residues in this sequence are susceptible to tTG-mediated deamidation (underlined). Homology searches using WWW PepPepSearch have identified close matches to LQPQNPSQQQPQ (SEQ ID NO:43) only in wheat  $\alpha$ -gliadins.

The fourth HLA-DQ8 subject (C11) had inducible IFN $\gamma$  ELISpot responses to tTG-treated Pool 33 (see Figure 36). Pools 32 and 33 include

5 polymorphisms of a previously defined HLA-DQ8 restricted gliadin epitope (QQYPSGQGSFQPSQQNPQ (SEQ ID NO:44)) active after deamidation by tTG (underlined Gln are deamidated and convey bioactivity) (van der Wal et al 1998). Currently, it is not known which peptides are bioactive in Pool 33 in subject C11.

Comprehensive T cell epitope mapping in HLA-DQ2-associated coeliac  
 10 disease using in vivo gluten challenge and a set of 652 peptides spanning all known 12 amino acid sequences in wheat gliadin has thus identified at least 72 peptides at 10% as bioactive as the known  $\alpha$ -gliadin epitope, A-gliadin 57-73 E65. However, these bioactive peptides can be reduced to a set of perhaps as few as 5 distinct but closely related families of peptides. Almost all these peptides are  
 15 rich in proline, glutamine, phenylalanine, and/or tyrosine and include the sequence PQ(QL)P(FY)P (SEQ ID NO:45). This sequence facilitates deamidation of Q in position 2 by tTG. By analogy with deamidation of A-gliadin 57-68 (Arentz-Hansen 2000), the enhanced bioactivity of these peptides generally found with deamidation by tTG may be due to increased affinity of  
 20 binding for HLA-DQ2.

Cross-reactivity amongst T cells in vivo recognizing more than one of these bioactive gliadin peptides is possible. However, if each set of related peptides does activate a distinct T cell population in vivo, the epitope corresponding to A-gliadin 57-73 E65 is the most potent and is generally  
 25 recognized by at least 40% of the peripheral blood T cells that secrete IFN $\gamma$  in response to gliadin after gluten challenge.

No gliadin-peptide specific responses were found in HLA-DQ2/8 coeliac disease that differed qualitatively from those in HLA-DQ2/X-associated coeliac disease. However, peripheral blood T cells in HLA-DQ8+ coeliac subjects  
 30 without both HLA-DQ2 alleles did not recognize A-gliadin 57-73 E65 homologues. Two different epitopes were dominant in two HLA-DQ8+ coeliacs. The dominant epitope in one of these HLA-DQ8+ individuals has not been identified previously (LQPQNPSQQQPQ (SEQ ID NO:46)).

Given the teaching herein, design of an immunotherapy for coeliac  
 35 disease utilizing all the commonly recognised T cell epitopes is practical and may

- 5 include fewer than six distinct peptides. Epitopes in wheat  $\gamma$ - and  $\omega$ -gliadins are also present in barley hordeins and rye secalins.

#### Example 14

- Several ELISpot assays were performed as previously described and  
10 yielded the following results and/or conclusions:

##### *Examination of multiple $\alpha$ -gliadin polymorphisms with PQLPY*

Potent agonists of A-gliadin 57-73QE (G01) include

- QLQPFPPQELPYPQPQS (G01) (SEQ ID NO:2),  
PQL-Y-----P (G10), and  
15 PQPQPFL----- (G12).

Less potent include

- L-----P (G04),  
-----R-----P (G05), and  
-----S-----P (G06).

- 20 Less potent yet include

-----L-----S-----P (G07),  
-----S-----S-----P (G08),  
-----S--S-----P (G09), and  
PQPQPFP----- (G13).

- 25 Dashes indicate identity with the G01 sequence in the particular position.

##### *Gluten challenge induces A-gliadin 57-73 QE65 T cells only after two weeks of gluten-free diet in newly diagnosed coeliac disease*

- Additional analyses indicated that tTG-deamidated gliadin responses  
30 change after two weeks of gluten-free diet in newly diagnosed coeliac disease.  
Other analyses indicated that deamidated gliadin-specific T cells are CD4<sup>+</sup> $\alpha_4\beta_7$ <sup>+</sup>  
HLA-DQ2 restricted.

##### *Optimal epitope (clones versus gluten challenge)*

5 A “dominant” epitope is defined by  $\gamma$ IFN ELISpot after gluten challenge.  
 QLQPFQPPELPYPQPQS (SEQ ID NO:2) (100% ELISpot response). Epitopes  
 defined by intestinal T cell clones: QLQPFQPPELPY (SEQ ID NO:13) (27%),  
 PQPELPYPQPPELPY (SEQ ID NO:47) (52%), and  
 QQLPQPEQPQSFPEQERPF (SEQ ID NO:48)(9%).

10

### *Dominance among individual peptide responses*

Dominance depends on wheat or rye. For wheat, dominant peptides  
 include peptide numbers 89, 90 and 91 (referring to sequence numbers in Table  
 15 23). For rye, dominant peptides include peptide numbers 368, 369, 370, 371, and  
 372 (referring to sequence numbers in Table 23). Some peptides, including 635  
 and 636 (referring to sequence numbers in Table 23) showed activity in both rye  
 and wheat.

### *In vivo gluten challenge allows T cell epitope hierarchy to be defined for coeliac disease*

The epitope hierarchy is consistent among HLA-DQ2<sup>+</sup> coeliacs but  
 different for HLA-DQ8<sup>+</sup> coeliacs. The hierarchy depends on what cereal is  
 consumed. Deamidation generates almost all gliadin epitopes. HLA-DQ2, DQ8,  
 25 and DR4 present deamidated peptides. HLA-DQ2/8-associated coeliac disease  
 preferentially present DQ2-associated gliadin epitopes. Gliadin epitopes are  
 sufficiently restricted to justify development of epitope-based therapeutics.

Other analyses indicated the following: HLA-DR3-DQ2 (85-95%) and  
 HLA-DR4-DQ8 (5-15%).

30 Other analyses indicated the following:

| HLA-DQ | HLA-DQA1 | HLA-DQB1 | Duodenal  | Gluten | EMA    |
|--------|----------|----------|-----------|--------|--------|
| on     |          |          |           |        |        |
|        | allele   | allele   | histology | free   | gluten |
|        |          |          |           |        | (on    |

35 GFD)



|    |     |       |              |               |     |       |        |
|----|-----|-------|--------------|---------------|-----|-------|--------|
| 5  | C01 | 2,6   | 102/6, 501   | 201, 602      | SVA | 1 yr  | +(-)   |
|    | C02 | 2,2   | 501          | 201           | SVA | 1 yr  | +(-)   |
|    | C03 | 2,5   | 101/4/5, 501 | 201, 501      | PVA | 1 yr  | +(-)   |
|    | C04 | 2,5   | 101/4/5, 501 | 201, 501      | SVA | 7 yr  | +(-)   |
|    | C05 | 2,2   | 201, 501     | 201, 202      | SVA | 4 mo  |        |
| 10 |     | +(ND) |              |               |     |       |        |
|    | C06 | 2,2   | 201, 501     | 201, 202      | SVA | 2 yr  | +(-)   |
|    | C07 | 2,8   | 301-3, 501   | 201, 302      | SVA | 1 yr  | +(-)   |
|    | C08 | 2,8   | 301-3, 501   | 201,302/8     | SVA | 11 yr | ND (-) |
| 15 | C09 | 2,8   | 301-3, 501   | 201, 302      | SVA | 29 yr | +(-)   |
|    | C10 | 2,8   | 201, 301-3   | 202, 302      | IEL | 1 yr  | +(-)   |
|    | C11 | 6,8   | 102/6, 301-3 | 602/15, 302/8 | IEL | 9 mo  | -      |
|    |     | (ND)  |              |               |     |       |        |
|    | C12 | 8,7   | 301-3, 505   | 302, 301/9-10 | SVA | 2 yr  | - (-)  |
| 20 | C13 | 8,8   | 301          | 302           | SVA | 1 yr  | + (+)  |

Another analysis was carried out to determine the bioactivity of individual tTG-deamidated peptides in pools 1-3 in subject C12. The results are as follows (sequence numbers refer to the peptides listed in Table 23): Sequence 8 (100%),

25 Sequence 5 (85%), Sequence 6 (82%), Sequence 3 (77%), Sequence 1 (67%), Sequence 2 (59%), Sequence 9 (49%), Sequence 7 (49%), Sequence 10 (33%), Sequence 4 (15%), Sequence 12 (8%), Sequence 11 (0%), Sequence 23 (26%), Sequence 14 (18%), Sequence 15 (18%), Sequence 17 (18%), Sequence 16 (13%), Sequence 14 (8%), Sequence 22 (5%), Sequence 18 (3%), Sequence 19

30 (3%), Sequence 20 (0%), Sequence 21 (0%). The predicted deamidated sequence is LQPENPSQEQPE(SEQ ID NO:22).

*Individual ELISpot responses by PBMC (Spot forming cells determined by ELISpot Reader)*

35 Peptide (see Table 23) C01 C02 C03 C04 C05

|    |    |     |    |    |    |    |
|----|----|-----|----|----|----|----|
| 5  | 65 | 16  | 2  | 1  | 2  | 3  |
|    | 66 | 32  | 6  | 13 | 0  | 6  |
|    | 67 | 16  | 3  | 4  | 0  | 4  |
|    | 68 | 25  | 8  | 4  | 2  | 2  |
|    | 69 | 4   | 0  | 0  | 0  | 0  |
| 10 | 70 | 2   | 1  | 0  | 0  | 0  |
|    | 71 | 1   | 1  | 0  | 0  | 1  |
|    | 72 | 0   | 0  | 0  | 0  | 0  |
|    | 73 | 95  | 21 | 42 | 31 | 31 |
|    | 74 | 122 | 15 | 29 | 21 | 28 |
| 15 | 75 | 5   | 1  | 2  | 2  | 5  |
|    | 76 | 108 | 13 | 28 | 16 | 22 |
|    | 77 | 3   | 0  | 1  | 0  | 1  |
|    | 78 | 21  | 2  | 3  | 5  | 3  |
|    | 79 | 20  | 0  | 2  | 0  | 2  |
| 20 | 80 | 5   | 2  | 0  | 0  | 3  |
|    | 81 | 4   | 1  | 2  | 3  | 1  |
|    | 82 | 3   | 3  | 5  | 2  | 2  |
|    | 83 | 14  | 2  | 0  | 0  | 1  |
|    | 84 | 3   | 0  | 0  | 0  | 0  |
| 25 | 85 | 14  | 1  | 2  | 1  | 2  |
|    | 86 | 11  | 0  | 2  | 0  | 2  |

### *Cross-reactivity*

To deal with data from 652 peptides in 29 subjects, or to determine when a particular response is a true positive peptide-specific T-cell response, or to determine when a response to a peptide is due to cross-reactivity with another structurally related peptide, expression of a particular peptide response can be as a percentage of a “dominant” peptide response. Alternately, the expression can be a “relatedness” as correlation coefficients between peptide responses, or via bioinformatics.

5

*Additional epitopes*

A representative result is as follows:

Combination of peptides with P04722E (all 20mcg/ml) (n=4)

|            | Alone | P04722E+ |
|------------|-------|----------|
| 10 Pep 626 | 60    | 135      |
| P04722E    | 100   | 110      |
| HLAa       | 0     | 85       |

(expressed as percent P04722E)

626+tT: PQQPQQPQQPFPQPQQPFPW (SEQ ID NO:31)

15 P04724E: QLQPFQPELPYPQPQL (SEQ ID NO:49)

TTG-deamidation of peptide 626 (n=12)

No tTG = 100%

20 TTG = 170%

*Substitution at particular positions*

Substitution of Peptide 626 PQQP[Q1]QP[Q2]QPFQ[PQ3]QPFPV (n=12)

|       | Glu | Arg |
|-------|-----|-----|
| 25 Q1 | 95  | 90  |
| Q2    | 145 | 80  |
| Q3    | 155 | 10  |

(expressed as percent wild-type peptide)

30 Bioactivity of tTG-treated 15mers spanning Peptide 626/627

(PQQPQQPQQPFPQPQQPFPWQP(SEQ ID NO:31)) (n=8)

|          |    |
|----------|----|
| P1-15    | 5  |
| P2-16    | 4  |
| P3-17    | 3  |
| 35 P4-18 | 38 |

|   |       |    |
|---|-------|----|
| 5 | P5-19 | 65 |
|   | P6-20 | 95 |
|   | P7-21 | 65 |
|   | P8-22 | 90 |

(expressed as percent of maximal 15mer response)

10

*Multiple epitopes:*

P04724E: QLQPFPPQLPYPQPQL (SEQ ID NO:51)

626+TGTG: PQQPQQPQQPFPQQPFPW (SEQ ID NO:31)

Minimal epitope: QPQQPFPQQPFPW (SEQ ID NO:52)

15 Immunomagnetic depletion of PBMC by beads coated with anti-CD4 and by anti-integrin  $\beta_7$  depleted IFN $\gamma$  ELISpot responses, while immunomagnetic depletion of PBMC by beads coated with anti-CD8 or anti- $\alpha$ E integrin. Thus, the PBMC secreting IFN $\gamma$  are CD4+ and  $\alpha_4\beta_7$ +, associated with homing to the lamina propria in the gut.

20 Blocked by anti-DQ antibody but not by anti-DR antibody in heterozygotes and homozygotes for HLA-DQ2. This may imply multiple epitopes within one sequence.

*T cell epitopes in coeliac disease*

25 Other investigators have characterized certain intestinal T cell clone epitopes. See, e.g., Vader et al., Gastroenterology 2002, 122:1729-37; Arentz-Hansen et al., Gastroenterology 2002, 123:803-809. These are examples of epitopes whose relevance is at best unclear because of the in vitro techniques used to clone T cells.

30

*Intestinal versus peripheral blood clones*

Intestinal: 1) intestinal biopsies, 2) T cell clones raised against peptic-tryptic digest of gluten, 3) all HLA-DQ2 restricted, 4) clones respond to gliadin deamidated by transglutaminase.

- 5 Peripheral blood: 1) T cell clones raised against gluten are HLA-DR, DQ and DP restricted. Result: Intestinal T cell clones can be exclusively used to map coeliac disease associated epitopes

GDA\_9Wheat 307 aa Definition Alpha/Beta-Gliadin MM1 Precursor (Prolamin)

- 10 Accession P18573 -- Genbank (which is incorporated herein by reference in its entirety)

#### *Intestinal T cell clone epitopes*

- A definition of intestinal T cell clone epitopes can be found in, for  
 15 example, Arentz-Hansen et al., J Exp Med. 2000, 191:603-12. Also disclosed therein are gliadin epitopes for intestinal T cell clones. Deamidated QLQPFPPQQLPY (SEQ ID NO:53) is an epitope, with a deamidated sequence of QLQPFPPQPELPY (SEQ ID NO:13). There is an HLA-DQ2 restriction. A homology search shows other bioactive rAlpha-gliadins include PQQQLPY (SEQ  
 20 ID NO:4) singly or duplicated. A majority of T cell clones respond to either/or DQ2- $\alpha$ I: QLQPFPPQPELPY (SEQ ID NO:13) DQ2- $\alpha$ II: PQPELPYPQPELPY (SEQ ID NO:47)

#### *Dominant gliadin T cell epitopes*

- 25 All deamidated by transglutaminase.  
 Peripheral blood day 6 after gluten challenge: A-gliadin 57-73:  
 QLQPFPPQPELPYPQPQS (SEQ ID NO:2)  
 Intestinal T cell clones: DQ2- $\alpha$ I: QLQPFPPQPELPY (SEQ ID NO:13) DQ2- $\alpha$ II:  
 PQPELPYPQPELPY (SEQ ID NO:47)

30

#### *Intestinal T-cell Clone Epitope Mapping*

- |                    |         |                            |
|--------------------|---------|----------------------------|
| $\alpha$ -Gliadins | A1      | PFPQPQLPY (SEQ ID NO:54)   |
|                    | A2      | PQPQLPYPQ (SEQ ID NO:55)   |
|                    | A3      | PYPQPQLPY (SEQ ID NO:56)   |
| 35                 | Glia-20 | PQQPYPQPQPQ (SEQ ID NO:57) |

|    |            |         |                                  |
|----|------------|---------|----------------------------------|
| 5  | Γ-Gliadins | G1      | PQQSFPQQQ(SEQ ID NO:58)          |
|    |            | G2      | IIPQQPAQ(SEQ ID NO:59)           |
|    |            | G3      | FPQQPQQPYQPQ(SEQ ID NO:60)       |
|    |            | G4      | FSQPQQQFPQPQ(SEQ ID NO:61)       |
|    |            | G5      | LQPQQPFPPQQQPYQPQQ(SEQ ID NO:62) |
| 10 | Glu-21     |         | QSEQSQQPFPPQQF(SEQ ID NO:63)     |
|    |            | Glu-5   | Q(IL)PQQPQQF(SEQ ID NO:64)       |
|    | Glutenin   | Glt-156 | PFSQQQQSPF(SEQ ID NO:65)         |
|    |            | Glt-17  | PFSQQQQQ(SEQ ID NO:66)           |

15

*Gluten exposure and induction of IFN $\gamma$ -secreting A-Gliadin 57-73QE65-specific T cells in peripheral blood*

20        Untreated coeliac disease, followed by gluten free diet for 1, 2, or 8 weeks, followed by gluten exposure (3 days bread 200g/day), followed by gluten free diet

      Result 1: Duration of gluten free diet and IFN $\gamma$  ELISpot responses on day 0 and day 6 of gluten challenge: A-gliadin 57-73 QE65 (results expressed as IFN $\gamma$  specific spots/million PPBMC)

25        Day 0: none (5), 1 week (1), 2 weeks (2), 8 weeks (1)

      Day 6: none (0), 1 week (4), 2 weeks (28), 8 weeks (48)

      Result 2: Duration of gluten free diet and IFN $\gamma$  ELISpot responses on day 0 and day 6 of gluten challenge: tTG-gliadin (results expressed as IFN $\gamma$  specific spots/million PPBMC)

30        Day 0: none (45), 1 week (62), 2 weeks (5), 8 weeks (5)

      Day 6: none (0), 1 week (67), 2 weeks (40), 8 weeks (60)

      Result 3: Duration of gluten free diet and IFN $\gamma$  ELISpot responses on day 0 and day 6 of gluten challenge: A-gliadin 57-73 P65 (results expressed as IFN $\gamma$  specific spots/million PPBMC)

35        Day 0: none (1), 1 week (2), 2 weeks (1), 8 weeks (1)

- 5 Day 6: none (0), 1 week (0), 2 weeks (0), 8 weeks (0)  
 Result 4: Duration of gluten free diet and IFN $\gamma$  ELISpot responses on day  
 0 and day 6 of gluten challenge: PPD (results expressed as IFN $\gamma$  specific  
 spots/million PPBMC)  
 Day 0: none (90), 1 week (88), 2 weeks (210), 8 weeks (150)  
 10 Day 6: none (0), 1 week (100), 2 weeks (210), 8 weeks (100)  
 Result 5: Duration of gluten free diet and IFN $\gamma$  ELISpot responses on day  
 0 and day 6 of gluten challenge: tTG (results expressed as IFN $\gamma$  specific  
 spots/million PPBMC)  
 Day 0: none (5), 1 week (4), 2 weeks (3), 8 weeks (2)  
 15 Day 6: none (0), 1 week (4), 2 weeks (1), 8 weeks (2)

*Gluten challenge in HLA-DQ2 coeliac disease on long term gluten*

- Characterization of anti-gliadin T cell response was carried out in  
 peripheral blood on day 6-8 after 3-day gluten challenge.  
 20 Result 1: PBMC Day 6 Long-term gluten free diet (preincubation with anti-HLA-  
 DR and -DQ antibody) (expressed as % inhibition)  
 DR-: tTG-gliadin 100 mcg/ml (105), A-gliadin 57-73 QE65 50 mcg/ml  
 (90), PPD 5 mcg/ml (30)  
 DQ-: tTG-gliadin 100 mcg/ml (5), A-gliadin 57-73 QE65 50 mcg/ml  
 25 (22), PPD 5 mcg/ml (78).  
 Result 2: PBMC Day 6 Long-term gluten free diet (expressed as % CD8-  
 depleted PBMC response)  
 B7 depletion: tTG-gliadin n=6 (7), A-gliadin 57-73 n=9 (6), PPD n=8  
 (62)  
 30 AE depletion: tTG-gliadin n=6 (120), A-gliadin 57-73 n=9 (80), PPD n=8  
 (110).  
 CD4 depletion: tTG-gliadin n=6 (10), A-gliadin 57-73 n=9 (9), PPD n=8  
 (10).

- 35 *Therapeutic peptides include, but are not limited to*

- 5 QLQPFPPQLPYQPQS(SEQ ID NO:10) (AG01)  
 QLQPFPPQLPYQPQP(SEQ ID NO:26) (AG02)  
 QLQPFPPQLPYQPQL(SEQ ID NO:51) (AG03)  
 QLQPFPPQLPYLQPQP (SEQ ID NO:67) (AG04)  
 QLQPFPPQLPYQPQP (SEQ ID NO:68) (AG05)  
 10 QLQPFPPQLPYSQPQP(SEQ ID NO:28) (AG06)  
 QLQPFPPQLPYSQPQP(SEQ ID NO:69) (AG07)  
 QLQPFPPQLPYSQPQP(SEQ ID NO:70) (AG08)  
 QLQPFPPQLSYSQPQP(SEQ ID NO:71) (AG09)  
 PQLPYPPQLPYQPQP(SEQ ID NO:72) (AG10)  
 15 PQLPYPPQLPYQPQL(SEQ ID NO:73) (AG11)  
 PQPQFPPQLPYQPQS(SEQ ID NO:74) (AG12)  
 PQPQFPPQLPYQPQS(SEQ ID NO:75) (AG13)  
 PQPQFPPQLPYQYQP(SEQ ID NO:76) (AG14)  
 PQPQFPPQLPYQPPP(SEQ ID NO:77) (AG015)

20

Briefly after oral antigen challenge, specificities of peripheral blood T cells reflect those of intestinal T cell clones. In peripheral blood, epitopes of intestinal T cell clones are sub-optimal compared to A-gliadin 57-73 QE65, which is an optimal  $\alpha$ -gliadin epitope.

25

### Example 15

ELISpot assays were also carried out for mapping purposes as follows.

*Fine-mapping the dominant DQ-8 associated epitope*

- |    | Sequence / sfc             | tTG-treated sequence / sfc |
|----|----------------------------|----------------------------|
| 30 | VPQLQPQNPSQQQPQEQV / 76(A) | RWPVPQLQPQNPSQQ / 60(L)    |
|    |                            | WPVPQLQPQNPSQQQ / 90(M)    |
|    | VPQLQPENPSQQQPQEQV / 3(B)  | PVPQLQPQNPSQQQP / 130(N)   |
|    | VPQLQPRNPSQQQPQEQV / 76(C) | VPQLQPQNPSQQQPQ / 140(O)   |
| 35 |                            | PQLQPQNPSQQQPQE / 59(P)    |



|    |                             |                         |
|----|-----------------------------|-------------------------|
| 5  | VPQLQPQNPSQEQPQEQV / 100(D) | QLQPQNPSQQQPQEQ / 95(Q) |
|    | VPQLQPQNPSQRQPQEQV / 1(E)   | LQPQNPSQQQPQEQV / 30(R) |
|    |                             | QPQNPSQQQPQEQVP / 4(S)  |
|    | VPQLQPQNPSQQQPQEQV / 71(F)  |                         |
| 10 | VPQLQPQNPSQQQPQEQV / 27(G)  | DQ8 Gliadin Epitope     |
|    |                             | GDA09 202Q / 6          |
|    | VPQLQPQNPSQEQPQEQV / 81(H)  | GDA09 202E / 83         |
|    | VPQLQPENPSQQQPQEQV / 2(I)   | GDA09 202Q+tTG / 17     |
|    | VPQLQPENPSQEQPQEQV / 6(J)   | BI + tTG / 0            |
| 15 | VPQLQPENPSQEQPQEQV / 5(K)   | BI / 0                  |
|    | A=SEQ ID NO:78              |                         |
|    | B=SEQ ID NO:79              |                         |
|    | C=SEQ ID NO:80              |                         |
| 20 | D=SEQ ID NO:81              |                         |
|    | E=SEQ ID NO:82              |                         |
|    | F=SEQ ID NO:83              |                         |
|    | G=SEQ ID NO:84              |                         |
|    | H=SEQ ID NO:85              |                         |
| 25 | I=SEQ ID NO:86              |                         |
|    | J=SEQ ID NO:87              |                         |
|    | K=SEQ ID NO:88              |                         |
|    | L=SEQ ID NO:89              |                         |
|    | M=SEQ ID NO:90              |                         |
| 30 | N=SEQ ID NO:91              |                         |
|    | O=SEQ ID NO:92              |                         |
|    | P=SEQ ID NO:93              |                         |
|    | Q=SEQ ID NO:94              |                         |
|    | R=SEQ ID NO:95              |                         |
| 35 | S=SEQ ID NO:96              |                         |

5

*Fine-mapping dominant epitope (2)*

Pool 33 (deamidated) / sfc

A2b3 301 qqyp sgqg ffqp sqqn pqaq (SEQ ID NO:359)/ 2

A2b5 301 qqyp sgqg ffqp fqqn pqaq (SEQ ID NO:360)/ 1

10 A3a1 301 qqyp sgqg ffqp sqqn pqaq (SEQ ID NO:361)/ 0

A3b1 301 qqyp ssqv sfqp sqln pqaq (SEQ ID NO:362)/ 0

A3b2 301 qqyp ssqg sfqp sqqn pqaq (SEQ ID NO:363)/ 2

A4a 301 eqyp sgqv sfqs sqqn pqaq (SEQ ID NO:364)/ 28

A1b1 309 sfrp sqqn plaq gsvq pqqI (SEQ ID NO:365)/ 2

15 A1a1 309 sfrp sqqn pqaq gsvq pqqI (SEQ ID NO:366)/ 2

**Example 16***Bioactivity of gliadin epitopes in IFN $\gamma$ -ELISpot (25 mcg/ml, n=6) (expressed as % A-gliadin 57-73 QE65 response)*

20 DQ2-AII: wild type (WT) (4), WT + tTG (52), Glu-substituted (52)

DQ2-AI: wild type (WT) (2), WT + tTG (22), Glu-substituted (28)

GDA09: wild type (WT) (1), WT + tTG (7), Glu-substituted (8)

A-G31-49: wild type (WT) (2), WT + tTG (3), Glu-substituted (0)

25 *Dose response of A-Gliadin 57-73 QE65 (G01E) (n=8) (expressed as %G01E maximal response)*

0.025 mcg/ml (1), 0.05 mcg/ml (8), 0.1 mcg/ml (10), 0.25 mcg/ml (22),

0.5 mcg/ml (38), 1 mcg/ml (43), 2.5 mcg/ml (52), 5 mcg/ml (70), 10 mcg/ml (81),

25 mcg/ml (95), 50 mcg/ml (90), 100 mcg/ml (85).

30 IFN $\gamma$  ELISpot response to gliadin epitopes alone or mixed with A-gliadin 57-75 (G01E) (all 50 mcg/ml, tTG-gliadin 100 mcg/ml, PPD 5 mcg/ml; n=9) (expressed as % G01E response)

Alone: DQ2-A1 (20), DQ2-A2 (55), Omega G1 (50), tTG Gliadin (80),

PPD (220), DQ2 binder (0)

5 G01E+: DQ2-A1 (90), DQ2-A2 (95), Omega G1 (100), tTG Gliadin  
(120), PPD (280), DQ2 binder (80)

*Effect of alanine and lysine substitution of A-gliadin 57-73 QE65 on IFN $\gamma$   
ELISpot responses in individual coeliac subjects (n=8)*

10 Epitope sequence: QLQFPQPPELPYPQPQS(SEQ ID NO:2)

Alanine substitution at positions 57-59 and 72-73 showed little to no  
decrease in % A-gliadin 57-73 QE65 response. Alanine substitution at positions  
60-62 and 68-71 showed moderate decrease in % A-gliadin 57-73 QE65  
response. Alanine substitution at positions 63-67 showed most decrease in % A-  
15 gliadin 57-73 QE65 response.

Effect of lysine substitution of A-gliadin 57-73 QE65 on IFN $\gamma$  ELISpot  
responses in individual coeliac subjects (n=8);

Epitope sequence: QLQFPQPPELPYPQPQS(SEQ ID NO:2)

Lysine substitution at positions 57-59 and 71-73 showed little to no  
20 decrease in % A-gliadin 57-73 QE65 response. Lysine substitution at positions  
60-61 and 69-70 showed moderate decrease in % A-gliadin 57-73 QE65  
response. Lysine substitution at positions 62-68 showed most decrease in % A-  
gliadin 57-73 QE65 response.

## 25 **Example 17**

Table 24 shows the results of analyses examining the 652 peptides with  
several patients challenged with wheat or rye.

## 30 References

1. Molberg O, et al. Nature Med. 4, 713-717 (1998).
2. Quarsten H, et al. Eur. J. Immunol. 29, 2506-2514 (1999).
3. Greenberg CS et al. FASEB 5, 3071-3077 (1991).
4. Mantzaris G, Jewell D. Scand. J. Gastroenterol. 26, 392-398 (1991).
- 35 5. Mauri L, et al. Scand. J. Gastroenterol. 31, 247-253 (1996).

- 5 6. Bunce M, et al. *Tissue Antigens* 46, 355-367 (1995).
7. Olerup O, et al. *Tissue antigens* 41, 119-134 (1993).
8. Mullighan CG, et al. *Tissue-Antigens*. 50, 688-92 (1997).
9. Plebanski M et al. *Eur. J. Immunol.* 28, 4345-4355 (1998).
- 10 10. Anderson DO, Greene FC. The alpha-gliadin gene family. II. DNA and protein sequence variation, subfamily structure, and origins of pseudogenes. *Theor Appl Genet* (1997) 95:59-65.
11. Arentz-Hansen H, Korner R, Molberg O, Quarsten H, Van der Wal Y, Kooy YMC, Lundin KEA, Koning F, Roepstorff P, Sollid LM, McAdam SN. The intestinal T cell response to alpha-gliadin in adult celiac disease is focused on a single deamidated glutamine targeted by tissue transglutaminase. *J Exp Med*. 2000; 191:603-12.
- 15 12. Vader LW, de Ru A, van der Wal, Kooy YMC, Benckhuijsen W, Mearin ML, Drijfhout JW, van Veelen P, Koning F. Specificity of tissue transglutaminase explains cereal toxicity in celiac disease. *J Exp Med* 2002; 195:643-649.
- 20 13. van der Wal Y, Kooy Y, van Veelen P, Pena S, Mearin L, Papadopoulos G, Koning F. Selective deamidation by tissue transglutaminase strongly enhances gliadin-specific T cell reactivity. *J Immunol*. 1998; 161:1585-8.
- 25 14. van der Wal Y, Kooy Y, van Veelen P, Pena S, Mearin L, Molberg O, Lundin KEA, Sollid L, Mutis T, Benckhuijsen WE, Drijfhout JW, Koning F. *Proc Natl Acad Sci USA* 1998; 95:10050-10054.
- 30 15. Vader W, Kooy Y, Van Veelen P et al. The gluten response in children with celiac disease is directed toward multiple gliadin and glutenin peptides. *Gastroenterology* 2002, 122:1729-37
- 35 16. Arentz-Hansen H, McAdam SN, Molberg O, et al. Celiac lesion T cells recognize epitopes that cluster in regions of gliadin rich in proline residues. *Gastroenterology* 2002, 123:803-809.

40

Each of the PCT publications, U.S. patents, other patents, journal references, and any other publications cited or referred to herein is incorporated herein by reference in their entirety.

5

**Table 1. A-Gliadin protein sequence (based on amino acid sequencing)  
(SEQ ID NO:3)**

10 VRVPVQLQP QNPSQQQPE QVPLVQQQF PGQQQQFPQP QPYPQFPFP SQQPYLQLQP FPQPQLPYQP  
1 11 21 31 41 51 61  
PQSFPQPQPY PQPQPQYSQP QQPISQQQAQ QQQQQQQQQ QQQILQQILQ QQLIPCMDVV LQQHINAHAR  
71 81 91 101 111 121 131  
SQVLQGSTYQ LLQELCCQHL WQIPEQSQCQ AIHNVVHAII LHQQQKQQQQ PSSQVSFQQP LQQYP LGQGS  
141 151 161 171 181 191 201  
15 FRPSQQNPQA QGSVPQQQLP QFEEIRNLAL QILPAMCNVY IAPYCTIAPF GIFGTN  
211 221 231 241 251 261

**Table 2. Coeliac disease subjects studied**

|   | Age<br>Sex | Gluten<br>free diet | HLA-DQ2      | Bread<br>challenge | Symptoms<br>with bread                                     |
|---|------------|---------------------|--------------|--------------------|--|
| 1 | 64 f       | 14 yr               | Homozygote   | 3 days             | Abdominal<br>pain, lethargy,<br>mouth ulcers,<br>diarrhoea |
| 2 | 57 m       | 1 yr                | Heterozygote | 10 days            | Lethargy,<br>nausea  |
| 3 | 35 f       | 7 yr                | Heterozygote | 3 days             | Nausea   |
| 4 | 36 m       | 6 wk                | Homozygote   | 3 days             | Abdominal<br>pain, mouth<br>ulcers,<br>diarrhoea           |
| 5 | 26 m       | 19 yr               | Heterozygote | 3 days             | None   |
| 6 | 58 m       | 35 yr               | Heterozygote | 3 days             | None   |
| 7 | 55 m       | 1 yr                | Heterozygote | 3 days             | Diarrhoea  |
| 8 | 48 f       | 15 yr               | Homozygote   | 3 days             | Abdominal<br>pain, diarrhoea                               |

20

5

| Aminoacid at position | Range   | Mean |
|-----------------------|---------|------|
| Glutamate             | (100)   | 100% |
| Asparagine            | (50-84) | 70%  |
| Aspartate             | (50-94) | 65%  |
| Alanine               | (44-76) | 64%  |
| Cysteine              | (45-83) | 62%  |
| Serine                | (45-75) | 62%  |
| Valine                | (24-79) | 56%  |
| Threonine             | (46-66) | 55%  |
| Glycine               | (34-47) | 40%  |
| Leucine               | (8-46)  | 33%  |
| Glutamine             | (16-21) | 19%  |
| Isoleucine            | (3-25)  | 14%  |
| Methionine            | (3-32)  | 14%  |
| Phenylalacine         | (0-33)  | 12%  |
| Histidine             | (0-13)  | 8%   |
| Tyrosine              | (0-17)  | 8%   |
| Tryptophan            | (0-17)  | 8%   |
| Lysine                | (0-11)  | 4%   |
| Proline               | (0-4)   | 2%   |
| Arginine              | (0-2)   | 1%   |

Table 3

| pt response | Peptides sequence                                | Corresponding residues in gliadin protein sequences<br>(Accession no)        |
|-------------|--|--|
| 13)         | QLQPSFPQLPYQPQ<br>(SEQ ID NO:10)                 | 57-73 $\alpha$ -Gliadin (T.aestivum) Q41545                                  |
| 100(100)    | QLQFFFPQLPVPQPS<br>(SEQ ID NO:2)                 | 57-73 $\alpha$ -Gliadin (T.aestivum) Q41545                                  |
| 7)          | QLQFFFPQLPYSQPQP<br>(SEQ ID NO:28)               | 77-93 $\alpha$ / $\beta$ -Gliadin precursor (Triticum.aestivum) P02863       |
|             |  | 76-92 $\alpha$ -Gliadin (T.aestivum) Q41528                                  |
|             |  | 77-93 $\alpha$ -Gliadin storage protein (T.aestivum) Q41531                  |
|             |  | 57-73 $\alpha$ -Gliadin mature peptide (T.aestivum) Q41533                   |
|             |  | 77-93 $\alpha$ -Gliadin precursor (T. spelta) Q92P09                         |
| -20)        | 83 (61-113) QLQFFFPQLPYQPQP'<br>(SEQ ID NO:26)   | 77-93 $\alpha$ / $\beta$ -Gliadin A-II precursor (T.aestivum) P0472          |
| -337)       | 83(74-97) QLQFFFPQLFVFPQL<br>(SEQ ID NO:51)      | 77-93 $\alpha$ / $\beta$ -Gliadin A-IV precursor (T.aestivum) P04724         |
|             |  | 77-93 $\alpha$ / $\beta$ -Gliadin MM1 precursor (T.aestivum)P18573           |
|             | 109(41-152)) PQLFYYPQLFYYPQPQP<br>(SEQ ID NO:72) | 84-100 $\alpha$ / $\beta$ -Gliadin A-IV precursor (T.aestivum) P04724        |
|             | PQLFYYPQLFYYPQPQL<br>(SEQ ID NO:73)              | 84-100 $\alpha$ / $\beta$ -Gliadin MM1 precursor (T.aestivum)P18573          |
| 3(0-7)      | QLQPFLQQLPYQPQP<br>(SEQ ID NO:69)                | 77-93 $\alpha$ / $\beta$ -Gliadin A-I precursor (T.aestivum) P04721          |
|             |  | 77-93 $\alpha$ -Gliadin (T.aestivum) Q41509                                  |
| 2(0-7)      | QLQFFSQQLPYSPQFP<br>(SEQ ID NO:70)               | 77-93 $\alpha$ -Gliadin storage protein (T.aestivum) Q41530                  |
|             | PQPQFFFPQLPYQTQP<br>(SEQ ID NO:97)               | 77-93 $\alpha$ / $\beta$ -Gliadin A-III precursor (T.aestivum) P04723        |
| -40)        | 24(11-43) PQPQFFFPQLPYQPQS<br>(SEQ ID NO:75)     | 82-98 $\alpha$ / $\beta$ -Gliadin A-V precursor (T.aestivum) P04725          |
| -30)        | 19(11-33) PQPQFFFPQLPYQPQP<br>(SEQ ID NO:77)     | 82-98 $\alpha$ / $\beta$ -Gliadin clone FW1215 precursor (T.aestivum) P04726 |
|             |  | 82-98 $\alpha$ / $\beta$ -Gliadin (T. urartu) Q41632                         |
| -30)        | 21(11-33) PQPQFLPQLPYQPQPQS<br>(SEQ ID NO:74)    | 79-95 $\alpha$ / $\beta$ -Gliadin clone FW8142 precursor (T.aestivum) P04726 |
|             |  | 79-95 $\alpha$ -Gliadin (T.aestivum) Q41529                                  |
|             |  | 79-95 $\alpha$ / $\beta$ -Gliadin precursor (T.aestivum) Q41546              |

## Table 4

5

Table 5. Table 5 T cell epitopes described in coeliac disease

| Source        | Restriction | Frequency     | Sequence*                   |
|---------------|-------------|---------------|-----------------------------|
| Gamma-gliadin | DQ2         | 3/NS (iTCC)   | QQLPQPFQPPQSSPPEQBRRPF(SEQ  |
| Alpha-gliadin | DQ2         | 12/17 (iTCL)  | QLQPELPQPELPY(SEQ ID NO:13) |
| Alpha-gliadin | DQ2         | 11/17 (iTCL)  | POPELPYPQPELPY(SEQ ID       |
| Alpha-gliadin | DQ2         | 1/23 (bTCC)   | LGQQQPFPPQPPYPQPPF(SEQ      |
| Alpha-gliadin | DQ8         | 3/NS (iTCC)   | QQYPSGEGSFPQSPENPQ(SEQ ID   |
| Glutenin      | DQ8         | 1/1 (iTCC)    | GQQGYYPISPPQSSGQ(SEQ ID     |
| Alpha-gliadin | DQ2         | 11/12 in vivo | QLQPFQPELPYPQPPQ(SAQ ID     |

NS not stated in original publication, iTCC intestinal T cell clone, iTCL intestinal polyclonal T cell line, bTCC peripheral blood T cell clone. All peptides are the products of transglutaminase modifying wild type gluten peptides except the fourth and sixth peptides

10 Table 6. Relative bioactivity of gliadin T cell epitopes in coeliac PBMC after gluten challenge  
ELISpot response as % A-gliadin 57-73 QE65 (all 2Smcg/ml)

| Sequence                             | Wild type | Wildtype+ITG | E-substituted |
|--------------------------------------|-----------|--------------|---------------|
| QQLPQPFQPPQSSPPEQBRRPF(SEQ ID NO:48) | 9(3)      | 18(7)        | 10(5)         |
| QLQPFQPELPY(SEQ ID NO:13)            | 6(2)      | 19(1)        | 8(3)          |
| PQPELPYPQPELPY(SEQ ID NO:47)         | 13(6)     | 53(8)        | 48(9)         |
| QQYPSGEGSFPQSPENPQ(SEQ ID NO:14)     | 10(3)     | 9(3)         | 14(8)         |
| QLQPFQPELPYPQPPQ(SAQ ID NO:2)        | 18(7)     | 87(7)        | 100           |
| PQLPYPQPELPYPQPPQ(SEQ ID NO:101)     | 14(4)     | 80 (17)      | 69(20)        |

Sequence refers that of transglutaminase (ITG) modified peptide and the T cell epitope. Wild type is the unmodified gliadin peptide. Data from 4 subjects. Blank was 5 (1) %.

15

**Table 7. Polymorphisms of A-gliadin 57-73****A. Sequences derived from Nordic autumn wheat strain Mjoelner**

| Alpha-gliadin protein (single letter code refers to Fig. 14 peptides) | Polymorphism                       |
|---|------------------------------------|
| Q41545 A-gliadin (from sequenced protein) 57-73 (A)                   | QLQPFPQQLPYYPQPS (SEQ ID NO:10)    |
| Gli alpha 1,6: (EMBL: AJ133605 & AJ133602 58-74) (J)                  | QBPQFPFPQLPYYPQTQP (SEQ ID NO:105) |
| Gli alpha 3,4,5: (EMBL: AJ133606, AJ133607, AJ133608 57-73) (I)       | QLQPFPQQLPYYPQPS (SEQ ID NO:28)    |
| Gli alpha 7: (EMBL: AJ133604 57-73) (E)                               | QLQPFPQQLPYYPQPS (SEQ ID NO:71)    |
| Gli alpha 8, 9, 11: (EMBL: ) (F)                                      | QLQPFPQQLPYYPQPS (SEQ ID NO:68)    |
| Gli alpha 10: (EMBL: AJ133610 57-73) (D)                              | QLQPFPQQLPYYPQPS (SEQ ID NO:28)    |
|   | QLQPFPQQLPYYPQPS (SEQ ID NO:104)   |

5 **B. SWISSPROT and TREMBL scan (10.12.99) for gliadins containing the sequence: XXXXXXXXPQLPYXXXXX**

| Wheat (Triticum aestivum unless stated) gliadin accession number | Polymorphism                      |
|--|-----------------------------------|
| Q41545 A-gliadin (from sequenced protein) 57-73 (A)              | QLQPFPQQLPYYPQPS (SEQ ID NO:10)   |
| SWISSPROT:   |                                   |
| GDA0_WHEAT P02863 77-93 (F)                                      | QLQPFPQQLPYYPQPS (SEQ ID NO:28)   |
| GDA1_WHEAT P04721 77-93 (G)                                      | QLQPFPQQLPYYPQPS (SEQ ID NO:69)   |
| GDA2_WHEAT P04722 77-93 (B)                                      | QLQPFPQQLPYYPQPS (SEQ ID NO:26)   |
| GDA3_WHEAT P04723 77-93 (O)                                      | PQBPFPFPQLPYYPQTQP (SEQ ID NO:97) |
| GDA4_WHEAT P04724 77-93 (C)                                      | QLQPFPQQLPYYPQPS (SEQ ID NO:51)   |
| GDA4_WHEAT P04724 84-100 (K)                                     | PQLPYYPQQLPYYPQPS (SEQ ID NO:72)  |
| GDA5_WHEAT P04725 82-98 (N)                                      | PQBPFPFPQLPYYPQPS (SEQ ID NO:75)  |
| GDA6_WHEAT P04726 82-98 (P)                                      | PQBPFPFPQLPYYPQPS (SEQ ID NO:77)  |
| GDA7_WHEAT P04727 79-95 (M)                                      | PQBPFPFPQLPYYPQPS (SEQ ID NO:74)  |
| GDA9_WHEAT P18573 77-93 (C)                                      | QLQPFPQQLPYYPQPS (SEQ ID NO:51)   |
| GDA9_WHEAT P18573 84-100 (L)                                     | PQLPYYPQQLPYYPQPS (SEQ ID NO:73)  |
| GDA9_WHEAT P18573 91-107 (K)                                     | PQLPYYPQQLPYYPQPS (SEQ ID NO:72)  |
| TREMBL   |                                   |
| Q41509 ALPHA-GLIADIN 77-93 (G)                                   | QLQPFPQQLPYYPQPS (SEQ ID NO:69)   |
| Q41528 ALPHA-GLIADIN 76-92 (F)                                   | QLQPFPQQLPYYPQPS (SEQ ID NO:28)   |
| Q41529 ALPHA-GLIADIN 79-95 (M)                                   | PQBPFPFPQLPYYPQPS (SEQ ID NO:74)  |
| Q41530 ALPHA-GLIADIN 77-93 (H)                                   | QLQPFPQQLPYYPQPS (SEQ ID NO:70)   |
| Q41531 ALPHA-GLIADIN 77-93 (F)                                   | QLQPFPQQLPYYPQPS (SEQ ID NO:28)   |
| Q41533 ALPHA-GLIADIN 57-73 (F)                                   | QLQPFPQQLPYYPQPS (SEQ ID NO:28)   |
| Q41546 ALPHA/BETA-GLIADIN 79-95 (M)                              | PQBPFPFPQLPYYPQPS (SEQ ID NO:74)  |
| Q41632 ALPHA/BETA-TYPE GLIADIN. Triticum uratur 82-98 (P)        | PQBPFPFPQLPYYPQPS (SEQ ID NO:77)  |
| Q9ZP09 ALPHA-GLIADIN Triticum spelta 77-93 (F)                   | QLQPFPQQLPYYPQPS (SEQ ID NO:28)   |



Table 8. Bioactivity of substituted variants of A-gliadin 57-73 QE65 (Subst) compared to unmodified A-gliadin 57-73 QE65 (G) (mean 100%, 95% CI 97-104) and blank (no peptide, bl) (mean 7.1%, 95% CI: 5.7-8.5)

| Subst            | %   | P vs G  | Subst | %  | P vs G  | Subst | %  | P vs G  | Subst        | %  | P vs G  | P vs bl |
|------------------|-----|---------|-------|----|---------|-------|----|---------|--------------|----|---------|---------|
| Super-agonists   |     |         |       |    |         |       |    |         |              |    |         |         |
| Y61              | 129 | <0.0001 | F62   | 71 | 0.001   | H62   | 47 | <0.0001 | N66          | 24 | <0.0001 |         |
|                  |     |         | V63   | 70 | <0.0001 | G69   | 47 | <0.0001 | R64          | 24 | <0.0001 |         |
| I                |     |         |       |    |         |       |    |         |              |    |         |         |
| Y70              | 129 | 0.0006  | S69   | 70 | <0.0001 | N63   | 47 | <0.0001 | K63          | 23 | <0.0001 |         |
|                  |     |         | H63   | 70 | <0.0001 | H68   | 47 | <0.0001 | V65          | 23 | <0.0001 |         |
| W70              | 119 | 0.017   | F63   | 70 | 0.008   | M68   | 46 | <0.0001 | H66          | 23 | <0.0001 |         |
| K57              | 118 | 0.02    | P70   | 69 | <0.0001 | D68   | 46 | <0.0001 | H67          | 22 | <0.0001 |         |
| Y59              | 117 | 0.04    | T62   | 69 | <0.0001 | V69   | 46 | <0.0001 | L64          | 22 | <0.0001 |         |
| A57              | 116 | 0.046   | L61   | 69 | <0.0001 | G63   | 45 | <0.0001 | S66          | 22 | <0.0001 |         |
| S70              | 116 | 0.045   | S61   | 69 | <0.0001 | V64   | 45 | <0.0001 | F67          | 21 | <0.0001 |         |
| K58              | 114 | 0.08    | T61   | 69 | <0.0001 | E61   | 45 | <0.0001 | W66          | 21 | <0.0001 |         |
| W59              | 110 | 0.21    | T63   | 69 | <0.0001 | A69   | 43 | <0.0001 | G64          | 21 | <0.0001 |         |
| A73              | 109 | 0.24    | M66   | 68 | <0.0001 | R62   | 42 | <0.0001 | G65          | 21 | <0.0001 |         |
| I59              | 108 | 0.37    | T69   | 67 | <0.0001 | G68   | 42 | <0.0001 | D64          | 21 | <0.0001 |         |
| G59              | 108 | 0.34    | K60   | 66 | <0.0001 | A64   | 42 | <0.0001 | I65          | 21 | <0.0001 |         |
| A58              | 108 | 0.35    | S62   | 66 | <0.0001 | C65   | 42 | <0.0001 | M64          | 20 | <0.0001 | <0.0001 |
| W60              | 105 | 0.62    | M61   | 66 | <0.0001 | N67   | 41 | <0.0001 | G67          | 19 | <0.0001 | <0.0001 |
| A59              | 104 | 0.61    | P61   | 65 | <0.0001 | W63   | 41 | <0.0001 | T65          | 19 | <0.0001 | 0.003   |
| K72              | 104 | 0.65    | M62   | 64 | <0.0001 | F69   | 41 | <0.0001 | A66          | 19 | <0.0001 | <0.0001 |
| S59              | 103 | 0.76    | Q61   | 64 | <0.0001 | N68   | 40 | <0.0001 | I64          | 19 | <0.0001 | 0.0003  |
| K73              | 102 | 0.8     | G61   | 64 | <0.0001 | V66   | 40 | <0.0001 | R63          | 19 | <0.0001 | <0.0001 |
| A70              | 102 | 0.81    | A63   | 64 | <0.0001 | H69   | 40 | <0.0001 | W67          | 19 | <0.0001 | <0.0001 |
| Y60              | 101 | 0.96    | L62   | 60 | <0.0001 | M69   | 40 | <0.0001 | K68          | 18 | <0.0001 | <0.0001 |
| A72              | 100 | 0.94    | I68   | 60 | <0.0001 | R69   | 40 | <0.0001 | H64          | 18 | <0.0001 | <0.0001 |
| S63              | 98  | 0.67    | S67   | 59 | <0.0001 | W69   | 40 | <0.0001 | W64          | 18 | <0.0001 | 0.0001  |
| K59              | 96  | 0.46    | N61   | 59 | <0.0001 | Q69   | 39 | <0.0001 | Q65          | 18 | <0.0001 | 0.0002  |
| I60              | 96  | 0.5     | I69   | 59 | <0.0001 | L67   | 38 | <0.0001 | F64          | 16 | <0.0001 | 0.0008  |
| G70              | 95  | 0.41    | V61   | 58 | <0.0001 | K69   | 38 | <0.0001 | L65          | 16 | <0.0001 | 0.0022  |
| D65              | 95  | 0.44    | D61   | 58 | <0.0001 | K62   | 38 | <0.0001 | N64          | 16 | <0.0001 | <0.0001 |
| E70              | 93  | 0.27    | E60   | 57 | <0.0001 | E67   | 37 | <0.0001 | F65          | 16 | <0.0001 | 0.12    |
| I63              | 92  | 0.19    | A61   | 57 | <0.0001 | L69   | 37 | <0.0001 | Q67          | 15 | <0.0001 | 0.0012  |
| S60              | 92  | 0.23    | Q62   | 56 | <0.0001 | S64   | 36 | <0.0001 | M65          | 14 | <0.0001 | 0.015   |
| P59              | 88  | 0.08    | F68   | 56 | <0.0001 | G62   | 36 | <0.0001 | D66          | 14 | <0.0001 | 0.013   |
| M63              | 87  | 0.03    | N65   | 56 | <0.0001 | E69   | 36 | <0.0001 | R67          | 14 | <0.0001 | 0.002   |
| K71              | 85  | 0.047   | A62   | 56 | <0.0001 | E68   | 36 | <0.0001 | Non-agonists |    |         |         |
| 10               |     |         |       |    |         |       |    |         |              |    |         |         |
| V62              | 84  | 0.04    | A68   | 53 | <0.0001 | V67   | 35 | <0.0001 | P63          | 13 | <0.0001 | 0.012   |
| I70              | 84  | 0.04    | P66   | 53 | <0.0001 | D62   | 35 | <0.0001 | E64          | 12 | <0.0001 | 0.053   |
| I61              | 83  | 0.01    | R61   | 53 | <0.0001 | R68   | 34 | <0.0001 | W65          | 11 | <0.0001 | 0.24    |
| V68              | 82  | 0.0045  | S68   | 53 | <0.0001 | Q66   | 34 | <0.0001 | Q64          | 11 | <0.0001 | 0.15    |
| E59              | 81  | 0.01    | V63   | 52 | <0.0001 | A67   | 33 | <0.0001 | G66          | 11 | <0.0001 | 0.07    |
| Partial agonists |     |         |       |    |         |       |    |         |              |    |         |         |
|                  |     |         | N69   | 51 | <0.0001 | N62   | 32 | <0.0001 | R65          | 11 | <0.0001 | 0.26    |
| W61              | 79  | 0.002   | E63   | 51 | <0.0001 | F66   | 31 | <0.0001 | Y67          | 10 | <0.0001 | 0.13    |
| A60              | 78  | 0.002   | T64   | 51 | <0.0001 | E62   | 31 | <0.0001 | E66          | 10 | <0.0001 | 0.17    |
| Y62              | 78  | 0.006   | T67   | 51 | <0.0001 | D69   | 31 | <0.0001 | K66          | 10 | <0.0001 | 0.21    |
| G60              | 77  | 0.003   | Y69   | 50 | <0.0001 | D67   | 30 | <0.0001 | R66          | 10 | <0.0001 | 0.23    |
| A71              | 77  | 0.003   | D63   | 50 | <0.0001 | M67   | 29 | <0.0001 | K67          | 10 | <0.0001 | 0.11    |
| W62              | 76  | 0.0009  | A65   | 49 | <0.0001 | Y66   | 28 | <0.0001 | P65          | 8  | <0.0001 | 0.57    |
| Q60              | 76  | 0.001   | K61   | 49 | <0.0001 | I67   | 28 | <0.0001 | K64          | 8  | <0.0001 | 0.82    |
| L63              | 74  | 0.0002  | I66   | 49 | <0.0001 | H65   | 26 | <0.0001 | K65          | 8  | <0.0001 | 0.63    |
| I62              | 74  | 0.0005  | T68   | 48 | <0.0001 | P68   | 26 | <0.0001 | Y65          | 7  | <0.0001 | 0.9     |
| K70              | 74  | 0.001   | S65   | 48 | <0.0001 | Y64   | 25 | <0.0001 |              |    |         |         |
| H61              | 72  | <0.0001 | L68   | 48 | <0.0001 | EK65  | 25 | <0.0001 |              |    |         |         |
| W68              | 72  | <0.0001 | Q68   | 48 | <0.0001 | T66   | 25 | <0.0001 |              |    |         |         |



5 **Table 9. Antagonism of A-gliadin 57-73 QE65 interferon gamma ELISPOT response by substituted variants of A-gliadin 57-73 QE65 (Subst) (P is significance level in unpaired t-test). Agonist activity (% agonist) of peptides compared to A-gliadin 57-73 QE65 is also shown.**

| Subst                 | % Inhibit. | P      | % agonist. | Subst | % Inhibit. | P    | % agonist. |
|-----------------------|------------|--------|------------|-------|------------|------|------------|
| Antagonists           |            |        |            | 13    | 0.18       | 11   |            |
| 65T                   | 28         | 0.004  | 19         | 65M   | 13         | 0.16 | 14         |
| 67M                   | 27         | 0.0052 | 29         | 68P   | 13         | 0.16 | 26         |
| 64W                   | 26         | 0.007  | 18         | 63R   | 13         | 0.19 | 19         |
| 67W                   | 25         | 0.0088 | 19         | 66G   | 12         | 0.19 | 11         |
| Potential antagonists |            |        |            | 65Q   | 12         | 0.2  | 18         |
| 67I                   | 24         | 0.013  | 10         | 65Y   | 12         | 0.22 | 7          |
| 67Y                   | 24         | 0.013  | 21         | 66S   | 12         | 0.22 | 22         |
| 64G                   | 21         | 0.03   | 21         | 67F   | 11         | 0.25 | 21         |
| 64D                   | 21         | 0.029  | 16         | 66R   | 10         | 0.29 | 10         |
| 65L                   | 20         | 0.046  | 26         | 67K   | 10         | 0.29 | 10         |
| 66N                   | 20         | 0.037  | 24         | 64F   | 10         | 0.29 | 16         |
| 65H                   | 20         | 0.038  | 16         | 65F   | 9          | 0.41 | 16         |
| 64N                   | 19         | 0.05   | 16         | 63P   | 8          | 0.42 | 13         |
| 64Y                   | 19         | 0.06   | 25         | 65EK  | 8          | 0.39 | 25         |
| 66Y                   | 19         | 0.048  | 28         | 64Q   | 7          | 0.49 | 11         |
| 64E                   | 19         | 0.049  | 12         | 64I   | 5          | 0.6  | 21         |
| 67A                   | 18         | 0.058  | 30         | 68K   | 5          | 0.56 | 19         |
| 67H                   | 18         | 0.052  | 22         | 67Q   | 5          | 0.61 | 18         |
| Non-antagonists       |            |        |            | 65G   | 5          | 0.62 | 15         |
| 65V                   | 17         | 0.07   | 23         | 64M   | 4          | 0.7  | 20         |
| 65I                   | 17         | 0.086  | 21         | 66H   | 4          | 0.66 | 23         |
| 66T                   | 17         | 0.069  | 25         | 66 E  | 3          | 0.76 | 10         |
| 65W                   | 15         | 0.11   | 11         | 66D   | 1          | 0.9  | 14         |
| 67R                   | 15         | 0.13   | 14         | 63K   | 1          | 0.88 | 23         |
| 65P                   | 15         | 0.13   | 8          | 64H   | 1          | 0.93 | 18         |
| 65K                   | 15         | 0.11   | 8          | 66K   | 0          | 0.98 | 10         |
| 66W                   | 15         | 0.12   | 21         | 64K   | -2         | 0.88 | 8          |
| 67G                   | 14         | 0.14   | 19         | 64L   | -11        | 0.26 | 22         |
| 66A                   | 14         | 0.14   | 19         |       |            |      |            |

**Table 10. Inhibition of A-gliadin 57-73 QE65 interferon gamma ELISPOT response by peptides known to bind HLA-DQ2 (P is significance level in unpaired t-test).**

20

| Peptide | % Inhibit. | P       |
|---------|------------|---------|
| TP      | 31         | <0.0001 |
| HLA1a   | 0          | 0.95    |

Table 11. Antagonism of A-gliadin 57-73 QE65 interferon gamma ELISpot response by naturally occurring polymorphisms of A-gliadin 57-73 QE65 (P is significance level in unpaired t-test).

| A-gliadin 57-73 QE65 polymorphism                           | % Inhibit. | P     |
|---|------------|-------|
| P04725 82-98 QE90 <u>PQPOPFPELPYPQPQS</u><br>(SEQ ID NO:17) | 19         | 0.009 |
| Q41509 77-93 QE85 QLQPF <u>L</u> QPELPYSQPQP                | 11         | 0.15  |
| Gli $\forall$ 1,6 58-74 QE66 QPQPFP <u>P</u> PELPYPQTQP     | 11         | 0.11  |
| P04723 77-93 QE85 <u>PQPQFPPELPYPQTQP</u>                   | 10         | 0.14  |
| Gli $\forall$ 3-5 57-73 QE65 QLQPFQPEL <u>S</u> VSQPQP      | 7          | 0.34  |
| P02863 77-93 QE85 QLQPFQPELPYSQP <u>Q</u> P                 | 6          | 0.35  |
| Q41509 77-93 QE85 QLQPF <u>L</u> QPELPYSQPQP                | 6          | 0.41  |
| (SEQ ID NO:29)  |            |       |
| P04727 79-95 QE65 <u>PQPOPEL</u> PELPYPQPQS                 | 6          | 0.39  |
| P04726 82-98 QE90 <u>PQPOPFPELPYPQPPP</u>                   | 5          | 0.43  |

5

10

**Table 12. Prolamin homologues of A-gliadin 57-73 (excluding alpha/beta-gliadins)**

| <b>Prolamin</b>           | <b>Accession number</b> | <b>Sequence</b>                         | <b>% Bioactivity*</b> |
|---------------------------|-------------------------|---|-----------------------|
| Wheat: $\alpha$ -gliadin  | A-gliadin (57-73)       | <b>QLQPFPPQQLPYPPQSQ</b> (SEQ ID NO:10) | 100 (0)               |
| Wheat: $\omega$ -gliadin  | AAG17702 (141-157)      | PQ.....F.....QSE                        | 32 (6.4)              |
| Barley: C-hordein         | Q40055 (166-182)        | ...QPPL.....F.....Q                     | 2.3 (2.0)             |
| Wheat $\gamma$ -gliadin   | P21292 (96-112)         | ...QTTPQ.....F.....QPQ                  | 2.1 (4.2)             |
| Rye: secalin              | Q43639 (335-351)        | ...QSPQ.....F.....Q                     | 1.6 (1.4)             |
| Barley: $\gamma$ -hordein | P80198 (52-68)          | ...QPFPQ.....HQQHFP                     | -1.0 (1.8)            |
| Wheat: LMW glutenin       | P16315 (67-83)          | LQ...QPIL.....FS...Q...Q                | -0.9 (1.0)            |
| Wheat: HMW glutenin       | P08489 (718-734)        | HGYYPIS.....SGQGQRP                     | 6.4 (4.0)             |
| Wheat $\gamma$ -gliadin   | P04730 (120-136)        | ...QCCQQL.....L...QSSRYQ                | 0.7 (0.9)             |
| Wheat: LMW glutenin       | P10386 (183-199)        | ...QCCQQL.....L...QSSRYE                | -0.7 (0.5)            |
| Wheat: LMW glutenin       | O49958 (214-230)        | ...QCCQQL.....L...EQSRYD                | -1.1 (0.3)            |
| Barley: B1-hordein        | P06470 (176-192)        | ...QCCQQL.....L...EQFRHE                | 1.8 (1.4)             |
| Barley: B-hordein         | Q40026 (176-192)        | ...QCCQQL.....L...EQFRHE                | 0.5 (0.9)             |

\*Bioactivity is expressed as 100x(spot forming cells with peptide 25mcg/ml plus tTG 8mcg/ml minus blank)/(spot forming cells with A-gliadin 57-73 25mcg/ml plus tTG 8mcg/ml minus blank) (mean (SEM), n=5). Peptides were preincubated with tTG for 2h 37°C. Note, Q is deamidated in A-gliadin 57-73 by tTG.

**Table 13. Clinical details of coeliac subjects.**

|     | HLA-DQ | HLA-DQA1<br>alleles | HLA-DQB1<br>alleles | Duodenal<br>histology | Gluten free | EMA on gluten<br>(on GFD) |
|-----|--------|---------------------|---------------------|-----------------------|-------------|---------------------------|
| C01 | 2, 6   | 102/6, 501          | 201, 602            | SVA                   | 1 yr        | + (-)                     |
| C02 | 2, 2   | 501                 | 201                 | SVA                   | 1 yr        | + (-)                     |
| C03 | 2, 5   | 101/4/5, 501        | 201, 501            | PVA                   | 1 yr        | + (-)                     |
| C04 | 2,5    | 101/4-5, 501        | 201, 501            | SVA                   | 7 yr        | + (-)                     |
| C05 | 2, 2   | 201, 501            | 201, 202            | SVA                   | 4 mo        | + (ND)                    |
| C06 | 2, 2   | 201, 501            | 201, 202            | SVA                   | 2 yr        | + (-)                     |
| C07 | 2, 8   | 301-3, 501          | 201, 302            | SVA                   | 1 yr        | + (-)                     |
| C08 | 2, 8   | 301-3, 501          | 201, 302/8          | SVA                   | 11 yr       | ND (-)                    |
| C09 | 2, 8   | 301-3, 501          | 201, 302            | SVA                   | 29 yr       | + (-)                     |
| C10 | 2, 8   | 201, 301-3          | 202, 302            | IEL                   | 1 yr        | + (-)                     |
| C11 | 6,8    | 102/6, 301-3        | 602/15, 302/8       | IEL                   | 9 mo        | - (ND)                    |
| C12 | 8,7    | 301-3, 505          | 302, 301/9-10       | SVA                   | 2 yr        | - (-)                     |
| C13 | 8, 8   | 301                 | 302                 | SVA                   | 1 yr        | + (+)                     |

SVA subtotal villous atrophy, PVA partial villous atrophy, IEL increased intra-epithelial atrophy, GFD gluten-free diet, ND not done.

**Table 14. HLA-DQ2+ Coeliac (C01-6) and healthy control (H01-10) IFN $\gamma$  ELISpot responses to control peptides (20  $\mu$ g/ml) and gliadin (500  $\mu$ g/ml) before and after gluten challenge (sfc/million PBMC minus response to PBS alone)**

| Peptide                    | Healthy Day 0 | Healthy Day 6     | Coeliac Day 0  | Coeliac Day 6      |
|----------------------------|---------------|-------------------|----------------|--------------------|
| P04722 77-93               | 0 (-4 to 17)  | 0 (-5 to 9)       | -2 (-3 to 0)   | 27 (0-100)*        |
| P04722 77-93 + tTG         | 0 (-5 to 4)   | 0 (-9 to 3)       | 0 (-4 to 11)   | 141 (8 to 290)**   |
| P04722 77-93 QE85          | 0 (-5 to 5)   | 0 (-3 to 4)       | 0 (-6 to 14)   | 133 (10 to 297)*   |
| P02863 77-93               | 0 (-4 to 13)  | 2 (-3 to 5)       | -2 (-3 to 2)   | 8 (-2 to 42)**     |
| P02863 77-93 + tTG         | -1 (-5 to 4)  | -1 (-4 to 11)     | 1 (-4 to 6)    | 65 (8-164)**       |
| P02863 77-93 QE85          | 0 (-4 to 13)  | 0 (-4 to 14)      | -1 (-4 to 6)   | 42 (-2 to 176)*    |
| Gliadin chymotrypsin       | 2 (-5 to 20)  | 18 (0 to 185)*    | 20 (11 to 145) | 92 (50 to 154)     |
| Gliadin chymotrypsin + tTG | 0 (-1 to 28)  | 16 (-9 to 171)*   | 55 (29 to 248) | 269 (206 to 384)** |
| Chymotrypsin               | 0 (-4 to 5)   | 1 (-4 to 11)      | -2 (-5 to 5)   | 1 (-4 to 8)        |
| Chymotrypsin + tTG         | 0 (-5 to 8)   | 6 (0 to 29)       | -2 (-3 to 11)  | 2 (-3 to 18)*      |
| Gliadin pepsin             | 4 (-4 to 28)  | 29 (0 to 189)***  | 44 (10 to 221) | 176 (54 to 265)**  |
| Gliadin pepsin + tTG       | 2 (-3 to 80)  | 27 (-4 to 241)*** | 61 (8 to 172)  | 280 (207 to 406)** |
| Pepsin                     | 0 (-4 to 10)  | 0 (-3 to 12)      | 0 (-2 to 3)    | 2 (-2 to 8)        |
| Pepsin + tTG               | 0 (-3 to 8)   | 0 (-5 to 9)       | 1 (-6 to 3)    | 0 (-3 to 14)       |
| PBS alone                  | 4 (0 to 6)    | 2 (0 to 6)        | 4 (1 to 12)    | 4 (0 to 4)         |
| PBS + tTG                  | 3 (0 to 8)    | 3 (0 to 11)       | 4 (2 to 10)    | 4 (2 to 11)        |

Day 6 vs. Day 0: \*P<0.05 \*\*P<0.02, \*\*\*P<0.01 by one-tailed Wilcoxon Matched-Pairs Signed-Ranks test

**Table 15. Effect of deamidation by tTG to gliadin (0.5 mg/ml) and A-gliadin 57-73 homologues on IFN $\gamma$  ELISpot responses in HLA-DQ2+ coeliac (C01-6) and healthy control subjects (H01-10) (median ratio tTG:no tTG pretreatment, range)**

| Peptide              | Healthy Day 6  | Coeliac Day 0  | Coeliac Day 6    |
|----------------------|----------------|----------------|------------------|
| Gliadin chymotrypsin | 0.94 (0.4-9.0) | 2.1 (0.8-6.8)* | 3.2 (1.8 -4.2)** |
| Gliadin pepsin       | 1.4 (0.5-1.4)  | 1.4 (0.8-4.0)* | 1.9 (1.1-4.4)**  |
| P04722 77-93 Q85     |                |                | 6.5 (2.3-12)**   |
| P04722 77-93 E85     |                |                | 0.7 (0.6-1.1)    |
| P02863 77-93 Q85     |                |                | 7.5 (3.9-19.9)** |
| P02863 77-93 E85     |                |                | 1.0 (0.8-1.2)    |

5      TTG>no tTG: \*P<0.05 \*\*P,0.02, \*\*\*P<0.01 by one-tailed Wilcoxon Matched-Pairs Signed-Ranks test



**Table 16. Healthy subjects: IFN $\gamma$  ELISpot Responses (>10 sfc/million PBMC and >4 x buffer only) to tTG-treated gliadin peptide Pools on Day 6 of gluten challenge (sfc/million PBMC) (*italic*: response also present on Day 0):**

**Group 1 – HLA-DQ2 (DQA1\*0501-5, DQB1\*0201)**

- 5 **Group 2 – HLA-DQ8 (DQA1\*0301, DQB1\*0302) and absent or “incomplete” DQ2 (only DQA1\*0501-5 or DQB1\*0201)**

| Subject | Group 1 |      |      |      |      |      |      |      |      |      | Group 2 |   |
|---------|---------|------|------|------|------|------|------|------|------|------|---------|---|
|         | H01     | H02  | H03  | H04  | H05  | H06  | H07  | H08  | H09  | H10  | H11     |   |
| HLA-DQ  | 2, 6    | 2, 7 | 2, 8 | 2, 5 | 2, 6 | 2, 6 | 2, 6 | 2, 7 | 2, 5 | 2, 5 | 8, 8    |   |
| Pool 1  | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 2       | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 3       | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 4       | .       | .    | .    | .    | .    | .    | .    | .    | 13   | .    | .       | . |
| 5       | .       | 17   | .    | .    | .    | .    | .    | .    | 24   | .    | .       | . |
| 6       | .       | .    | .    | .    | .    | .    | .    | .    | 31   | .    | .       | . |
| 7       | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 8       | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 9       | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 10      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 11      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 12      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 13      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 14      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 15      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 16      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 17      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 18      | .       | .    | .    | .    | .    | .    | 20   | .    | .    | .    | .       | . |
| 19      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 20      | .       | 11   | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 21      | .       | 11   | .    | .    | .    | .    | .    | .    | 27   | .    | .       | . |
| 22      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 23      | .       | 43   | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 24      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 25      | .       | 11   | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 26      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 27      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 28      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 29      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 30      | .       | .    | .    | .    | .    | .    | .    | 23   | .    | .    | .       | . |
| 31      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 32      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 33      | .       | 20   | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 34      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 35      | .       | 11   | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 36      | .       | .    | .    | .    | .    | .    | .    | .    | .    | .    | .       | . |
| 37      | .       | .    | .    | .    | .    | .    | .    | 18   | .    | .    | .       | . |
| 38      | 14      | .    | .    | .    | .    | .    | .    | 12   | .    | .    | .       | . |
| 39      | .       | .    | .    | .    | .    | .    | .    | 11   | .    | .    | .       | . |
| 40      | .       | 14   | .    | .    | .    | .    | .    | 17   | .    | .    | .       | . |

|                |     |     |    |    |    |   |    |    |    |    |
|----------------|-----|-----|----|----|----|---|----|----|----|----|
| 41             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 42             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 43             | .   | .   | .  | .  | .  | . | .  | .  | 11 | .  |
| 44             | .   | 14  | .  | .  | .  | . | .  | .  | .  | .  |
| 45             | .   | 11  | .  | .  | .  | . | .  | .  | .  | .  |
| 46             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 47             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 48             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 49             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 50             | .   | 14  | .  | .  | 12 | . | .  | 22 | .  | 14 |
| 51             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 52             | .   | 14  | .  | .  | .  | . | .  | .  | .  | .  |
| 53             | .   | 26  | .  | .  | .  | . | .  | .  | .  | .  |
| 54             | .   | .   | .  | .  | .  | . | .  | 12 | .  | .  |
| 55             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 56             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 57             | .   | 23  | .  | .  | .  | . | 12 | .  | .  | .  |
| 58             | .   | 14  | .  | .  | .  | . | .  | .  | .  | .  |
| 59             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 60             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 61             | .   | 23  | .  | .  | .  | . | .  | 11 | 11 | .  |
| 62             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 63             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 64             | .   | 20  | .  | .  | .  | . | .  | .  | .  | .  |
| 65             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 66             | .   | 14  | .  | .  | .  | . | .  | .  | .  | .  |
| 67             | .   | 11  | .  | .  | .  | . | .  | .  | .  | .  |
| 68             | .   | 20  | .  | .  | .  | . | .  | 20 | .  | .  |
| 69             | .   | 20  | .  | .  | .  | . | .  | .  | .  | .  |
| 70             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 71             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 72             | .   | 11  | .  | .  | .  | . | .  | .  | .  | .  |
| 73             | .   | 14  | .  | .  | .  | . | .  | .  | .  | .  |
| 74             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 75             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 76             | .   | 14  | .  | .  | .  | . | .  | .  | .  | .  |
| 77             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 78             | .   | 11  | .  | .  | .  | . | .  | .  | .  | .  |
| 79             | .   | 11  | .  | .  | 19 | . | .  | .  | .  | .  |
| 80             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 81             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 82             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| 83             | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| P04722 77-93   | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| P04722 77-93 E | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| P04722 77-93 E | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| P02863 77-93   | .   | .   | .  | .  | .  | . | .  | 11 | .  | .  |
| P02863 77-93 E | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |
| Gliadin+C      | 171 | 40  | 25 | 16 | 10 | . | 18 | 14 | .  | 17 |
| Chymotrypsin   | 29  | 26  | 18 | .  | .  | . | .  | 22 | .  | .  |
| Gliadin+Pepsin | 241 | 151 | 29 | 24 | 48 | . | 16 | 45 | .  | 19 |
| Pepsin         | .   | .   | .  | .  | .  | . | .  | .  | .  | .  |

**Table 17: tTG-deamidated gliadin peptide pools showing significant increase in IFN gamma responses between Day 0 and Day 6 of gluten challenge in HLA-DQ2 coeliac subjects C01-6 (Day 6 –Day 0 response, and ratio of responses to tTG-deamidated pool and same pool without tTG treatment)**

| IFN $\gamma$ ELISpot |                      |                         | IFN $\gamma$ ELISpot |                      |                         |
|----------------------|----------------------|-------------------------|----------------------|----------------------|-------------------------|
| Pool                 | (Median sfc/million) | tTG: no tTG<br>(Median) | Pool                 | (Median sfc/million) | tTG: no tTG<br>(Median) |
| 9                    | 59***                | 1.0                     | 49                   | 46***                | 1.4                     |
| 10                   | 116**                | 1.7                     | 50                   | 50***                | 4.6                     |
| 11                   | 24***                | 2.5                     | 51                   | 40***                | 1.7                     |
| 12                   | 133***               | 1.1                     | 52                   | 30***                | 3.1                     |
| 13                   | 26**                 | 2.1                     | 53                   | 27**                 | 1.4                     |
| 42                   | 30**                 | 1.2                     | 76                   | 17***                | 1.1                     |
| 43                   | 32***                | 1.3                     | 79                   | 20***                | 0.9                     |
| 44                   | 24***                | 1.5                     | 80                   | 83***                | 1                       |
| 45                   | 10***                | 1.1                     | 81                   | 141***               | 1.1                     |
| 46                   | 12***                | 2.1                     | 82                   | 22***                | 1.5                     |
| 48                   | 17***                | 1.4                     | 83                   | 16**                 | 1.8                     |

Day 6 vs. Day 0 \*\*P<0.02, \*\*\*P<0.01 by one-tailed Wilcoxon Matched-Pairs Signed-Ranks test

**Table 18. Coeliac subjects: IFN $\gamma$  ELISpot Responses >10 sfc/million PBMC and >4 x buffer only to tTG-treated Pepset Pools on Day 6 of gluten challenge (sfc/million**

**PBMC) (*italic*: response also present on Day 0):**

**Group 1 – HLA-DQ2 (DQA1\*0501-5, DQB1\*0201/2),**

**5 Group 2 – HLA-DQ2/8 (DQA1\*0501-5, \*0301, and DQB1\*0201/2, \*0302), and**

**Group 3 – HLA-DQ8 (DQA1\*0301, DQB1\*0302) and absent or “incomplete” DQ2 (only DQA1\*0501-5 or DQB1\*0201/2)**

| Group 1: |      |      |      |      |      |      | Group 2: |      |      | Group 3 |      |      |      |
|----------|------|------|------|------|------|------|----------|------|------|---------|------|------|------|
| Subject  | C01  | C02  | C03  | C04  | C05  | C06  | C07      | C08  | C09  | C10     | C11  | C12  | C13  |
| HLA-DQ   | 2, 6 | 2, 2 | 2, 5 | 2, 5 | 2, 2 | 2, 2 | 2, 8     | 2, 8 | 2, 8 | 2, 8    | 6, 8 | 7, 8 | 8, 8 |
| Pool 1   | .    | .    | .    | .    | .    | .    | 23       | .    | .    | .       | .    | 223  | .    |
| 2        | .    | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | 155  | .    |
| 3        | .    | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | 41   | .    |
| 4        | 11   | .    | .    | .    | .    | .    | .        | .    | .    | 22      | .    | .    | .    |
| 5        | .    | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 6        | 18   | .    | .    | 21   | .    | .    | 20       | 17   | .    | .       | .    | .    | .    |
| 7        | .    | .    | .    | .    | .    | .    | 353      | .    | .    | .       | .    | .    | .    |
| 8        | 11   | 64   | .    | .    | .    | 14   | 20       | 480  | .    | .       | .    | 13   | .    |
| 9        | 93   | 127  | .    | 92   | 25   | .    | 32       | 460  | .    | .       | .    | 18   | .    |
| 10       | 175  | 491  | 58   | 200  | 48   | .    | 84       | 787  | .    | .       | .    | .    | .    |
| 11       | 32   | 118  | .    | 33   | 14   | .    | 26       | 27   | .    | 12      | .    | .    | .    |
| 12       | 204  | 379  | 54   | 225  | 61   | .    | 129      | 587  | .    | 12      | .    | .    | .    |
| 13       | 93   | 142  | .    | 29   | 18   | .    | 60       | .    | .    | .       | .    | 11   | .    |
| 14       | .    | 45   | .    | 21   | .    | .    | 17       | .    | .    | .       | .    | .    | .    |
| 15       | 18   | 30   | .    | .    | .    | .    | 38       | 43   | .    | .       | .    | .    | .    |
| 16       | .    | .    | .    | .    | .    | .    | 37       | .    | .    | .       | .    | .    | .    |
| 17       | .    | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 18       | .    | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 19       | 11   | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 20       | 11   | 215  | .    | .    | .    | .    | 51       | 167  | .    | .       | .    | .    | .    |
| 21       | .    | .    | .    | .    | .    | .    | .        | .    | 11   | .       | .    | .    | .    |
| 22       | .    | 21   | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 23       | .    | 18   | .    | 21   | .    | .    | .        | .    | .    | 12      | .    | .    | .    |
| 24       | .    | 15   | .    | .    | .    | .    | 10       | .    | .    | .       | .    | .    | .    |
| 25       | .    | 15   | .    | .    | .    | .    | .        | .    | .    | 12      | .    | .    | .    |
| 26       | .    | 18   | .    | .    | .    | .    | 13       | .    | .    | 12      | .    | .    | .    |
| 27       | .    | 15   | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 28       | .    | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | 11   | .    |
| 29       | .    | .    | .    | .    | .    | .    | 11       | .    | .    | .       | .    | .    | .    |
| 30       | 11   | .    | .    | .    | .    | .    | 11       | .    | .    | .       | .    | .    | .    |
| 31       | .    | 70   | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 32       | .    | 18   | .    | .    | .    | .    | 20       | .    | .    | .       | .    | .    | .    |
| 33       | 11   | .    | .    | 10   | .    | .    | 14       | .    | 11   | .       | 40   | .    | 11   |
| 34       | .    | .    | .    | .    | .    | .    | .        | 11   | .    | .       | .    | .    | .    |
| 35       | .    | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 36       | .    | .    | .    | .    | .    | .    | .        | .    | .    | .       | .    | .    | .    |
| 37       | .    | .    | .    | 23   | .    | 14   | .        | .    | .    | .       | .    | .    | .    |



**Table 19. Deamidated peptides with mean bioactivity > 10% of P04722 E85 (20 µg/ml) in HLA-DQ2 coeliac subjects C01-5**

(Peptide Numbers in Table 19 correspond to Peptide Numbers and accompanying SEQ ID NOs. identified in Table 24)

| Rank | No. | Sequence             | Mean<br>(SEM) | Rank | No. | Sequence             | Mean<br>(SEM) |
|------|-----|----------------------|---------------|------|-----|----------------------|---------------|
|      | 89  | PQLPYTQFQLPYTQFQLPYT | 94 (18)       | 37   | 413 | SEQQQQFFPQQQQQSTQ    | 18 (4)        |
| *2   | 91  | PQFFPQLPYTQFQLPYTQF  | 89 (12)       | 38   | 380 | QFQQPQQFFPQQPQLPFP   | 18 (6)        |
| *3   | 74  | MQLEFFPQFQLPYTQFQLPY | 88 (14)       | 39   | 618 | PQQSTSYQQQFFPQQPYTQQ | 18 (7)        |
| *4   | 90  | PQLPYTQFQLPYTQFQFTR  | 87 (16)       | *40  | 78  | LQLQFFPQQLPYTQFQFTR  | 17 (8)        |
| *5   | 76  | LQLQFFPQQLPYTQFQFTR  | 85 (15)       | 41   | 390 | QQTYTQRQQFFPQTQQPQR  | 17 (9)        |
| 6    | 626 | PQQPQQPQQFFPQQPQFFW  | 72 (23)       | 42   | 348 | QQTYTQRQQTFTHHQQQQF  | 16 (10)       |
| 7    | 627 | QFFPQQPQQFQWQFQFFQ   | 66 (30)       | 43   | 409 | QQPQQFQLQQFQQLPQQP   | 16 (2)        |
| *8   | 631 | FPQQPQQFQQLPYTQFQ    | 61 (12)       | 44   | 382 | QQFFPQQPQQFFPQTQQPQQ | 16 (6)        |
| 9    | 636 | PQQPQQFQFQFQFQFQFQ   | 51 (10)       | 45   | 629 | FFPQTQQSFLQQFQFFQ    | 16 (5)        |
| *10  | 73  | LQLQFFPQQLPYTQFQLPY  | 49 (11)       | 46   | 643 | FLQFQQFFPQQPQQFFPQQP | 16 (6)        |
| 11   | 412 | SQQPQQFQFQFQFQFQFQ   | 34 (19)       | 47   | 389 | QQFFPQTQQFQFQFQFQFQ  | 16 (6)        |
| 12   | 343 | QQPQQFQFQFQFQFQFQ    | 34 (11)       | 48   | 350 | QQFFPQFQQTFFHQFQQAFF | 15 (8)        |
| *13  | 68  | LQLQFFPQQLPYTQFQFTR  | 33 (10)       | 49   | 65  | FFFSQQPQTQQFQFQFQF   | 15 (5)        |
| *14  | 66  | LQLQFFPQQLPYTSQFQFTR | 32 (7)        | 50   | 349 | QQFFPQFQQTFFHQFQQQF  | 15 (9)        |
| *15  | 96  | PQFFPQLPYTQFQFQFQ    | 28 (6)        | 51   | 610 | PWWQQQLPQQQSTQSQFQF  | 15 (11)       |
| 16   | 393 | QLFFPQQPQQFQFQFQFQ   | 27 (8)        | *52  | 81  | PQPQFFPQLPYTQFQFTR   | 15 (5)        |
| 17   | 355 | QAFFPQQTFFHQFQFQFQ   | 27 (15)       | *53  | 75  | MQQLPFFPQFQFQFQLPYTQ | 14 (5)        |
| *18  | 67  | LQLQFFPQQLPYTSQFQFTR | 26 (6)        | 54   | 368 | QQFFPQQPQQFQFQFQFQ   | 14 (7)        |
| 19   | 335 | QQQFFPQFQFQFQFQFQ    | 25 (11)       | *55  | 82  | PQPQFFPQFQFQFQLPYTQ  | 14 (3)        |
| *20  | 95  | PQFFLQLPYTQFQFQFQ    | 24 (6)        | *56  | 80  | LQLQFFPQFQFQFQLPYTQ  | 14 (4)        |
| 21   | 396 | TQQPQQFQFQFQFQFQ     | 23 (9)        | 57   | 624 | FTQQPQTFFHQFQFQFQ    | 14 (6)        |
| 22   | 609 | SCISGLERPWQQQLPQQ    | 23 (18)       | 58   | 407 | QFQQPQQSQFQFQFQFQ    | 14 (5)        |
| 23   | 385 | QQFFPQFQFQFQFQFQ     | 23 (7)        | 59   | 337 | QQFFPQFQFQFQFQFQ     | 13 (4)        |
| 24   | 375 | PQQFQFQFQFQFQFQFQ    | 23 (10)       | 60   | 634 | PQQQLQFQFQLPQFQFQFQ  | 13 (3)        |
| 25   | 406 | QFQFQFQFQFQFQFQFQ    | 22 (8)        | 61   | 388 | QQPYTQFQFQFQFQFQFQ   | 13 (3)        |
| 26   | 625 | FPQFQFQFQFQFQFQFQ    | 22 (9)        | 62   | 641 | FFELQFQFQFQFQFQFQ    | 13 (7)        |
| 27   | 378 | QQQFQFQFQFQFQFQFQ    | 22 (10)       | 63   | 399 | QQFQFQFQFQFQFQFQ     | 13 (5)        |
| 28   | 371 | PQQFQFQFQFQFQFQFQ    | 22 (10)       | 64   | 387 | QQTYTQRQQFQFQFQFQ    | 13 (4)        |
| 29   | 642 | PQQPQQFQFQFQFQFQFQ   | 20 (8)        | 65   | 628 | FFPWWQFQFQFQFQFQFQ   | 12 (4)        |
| 30   | 635 | FLQFQFQFQFQFQFQFQ    | 19 (5)        | *66  | 88  | PQFFPQLPYTSQFQFQFQ   | 12 (3)        |
| *31  | 93  | PQFFPQLPYTQFQFQFQ    | 19 (5)        | 67   | 408 | QFQFQFQFQFQFQFQFQ    | 12 (5)        |
| 32   | 377 | PQQFQFQFQFQFQFQFQ    | 19 (9)        | *68  | 77  | LQLQFQFQFQFQFQFQFQ   | 11 (4)        |
| 33   | 411 | LQQPQQFQFQFQFQFQFQ   | 19 (4)        | 69   | 370 | PQQQLPQQFQFQFQFQFQ   | 11 (5)        |
| 34   | 415 | SQQPQQFQFQFQFQFQFQ   | 18 (5)        | *70  | 79  | LQLQFQFQFQFQFQFQFQ   | 11 (5)        |
| *35  | 94  | PQFFPQLPYTQFQFQFQ    | 18 (3)        | 71   | 379 | QQPQQFQFQFQFQFQFQ    | 11 (5)        |
| 36   | 329 | PSQQVQWQQFQFQFQFQ    | 18 (4)        | 72   | 397 | PQQPQQFQFQFQFQFQFQ   | 11 (3)        |

**Table 20. Peptides >10% as bioactive as P04722 QE65 grouped by structure. (Peptide Numbers in Table 20 correspond to Peptide Numbers and their accompanying SEQ ID NOs. identified in Table 24)**

| Rank  | Peptide no.<br>(Pool)<br>Gliadin-subtype | Sequence                 | IFN $\gamma$ ELISpot<br>response<br>compared to<br>P04722 77-93<br>QE85: mean<br>(SEM) |
|---|--|--------------------------|--|
| <b>Group 1: Homologues of A-gliadin 57-73</b> |  |                          |  |
|   | P04722 77-93                             | <b>QLQFFPQQLPYQPQP</b>   |  |
| 1   | 89 (12) $\alpha$                         | PQL...Y.....LPYP         | 94 (18)  |
| 2   | 91 (12) $\alpha$                         | PQPFPQL...Y.....         | 89 (12)  |
| 3   | 74 (10) $\alpha$                         | M.....LPY                | 88 (14)  |
| 4   | 90 (12) $\alpha$                         | PQL...Y.....PFRP         | 87 (16)  |
| 5   | 76 (10) $\alpha$                         | L.....PFR                | 85 (15)  |
| 8   | 631 (81) $\omega$                        | FPQQPQ.....F....QS       | 61 (12)  |
| 10  | 73 (10) $\alpha$                         | L.....LPY                | 49 (11)  |
| 13  | 68 (9) $\alpha$                          | L.....PFR                | 33 (10)  |
| 14  | 66 (9) $\alpha$                          | L.....S.....PFR          | 32 (7)   |
| 18  | 67 (9) $\alpha$                          | L.....S.....QFR          | 26 (6)   |
| 20  | 95 (13) $\alpha$                         | PQPFL.....FPQQ           | 24 (6)   |
| 31  | 93 (12) $\alpha$                         | PQPFP.....PFRPQQ         | 19 (5)   |
| 35  | 94 (12) $\alpha$                         | PQFPF.....PPFSQQ         | 18 (3)   |
| 40  | 78 (10) $\alpha$                         | L.....R.....PFR          | 17 (8)   |
| 52  | 81 (11) $\alpha$                         | PQPQFP.....T...PFPF      | 15 (5)   |
| 53  | 75 (10) $\alpha$                         | MLQFPFPQFPF.....         | 14 (5)   |
| 55  | 82 (11) $\alpha$                         | PQPFPFPQFPF.....         | 14 (3)   |
| 56  | 80 (10) $\alpha$                         | LQLQFPFPQFPF.....        | 14 (4)   |
| 66  | 88 (11) $\alpha$                         | PQFPF.....S.....PFRPQQ   | 12 (3)   |
| 68  | 77 (10) $\alpha$                         | LQLQFPFPQFPF.....        | 11 (4)   |
| 70  | 79 (10) $\alpha$                         | LQLQFPFPQFPFL.....       | 11 (5)   |
| <b>Group 2: Homologues of peptide 626</b>     |  |                          |  |
|   |  | <b>QQFFPQFPQFPF</b>      |  |
| 6   | 626(80) $\omega$                         | PQQPQFP.....W            | 72 (23)  |
| 7   | 627(80) $\omega$                         | .....WQPQQFPFQ           | 66 (30)  |
| 9   | 636(81) $\omega$                         | PQQP.....L...VQFP        | 51 (10)  |
| 11  | 412(53) $\gamma$                         | SQQP.....Q.....PQQ       | 34 (19)  |
| 33  | 411(53) $\gamma$                         | LQQP.....Q.....PQQ       | 19 (4)   |
| 36  | 329(42) $\gamma$                         | PSGQVQWPO.....           | 18 (4)   |
| 41  | 390(50) $\gamma$                         | QQTYPQP.....T.....QQ     | 17 (9)   |
| 59  | 337(43) $\gamma$                         | Q.....CQQPQRTI           | 13 (4)   |
| 61  | 388(50) $\gamma$                         | QQPYPQQF.....T.....QQ    | 13 (3)   |
| <b>Group 3: Homologues of peptide 355</b>     |  |                          |  |
|   |  | <b>FPQPQQTFPHQPQQFPF</b> |  |
| 17  | 355(46) $\gamma$                         | QA.....Q                 | 27 (15)  |
| 42  | 348(45) $\gamma$                         | QQT.....                 | 16 (10)  |
| 48  | 350(45) $\gamma$                         | QQL.....A.....           | 15 (8)   |
| 50  | 349(45) $\gamma$                         | QQL.....                 | 15 (9)   |
| <b>Group 4: Homologues of Peptide 396</b>     |  |                          |  |
|   |  | <b>QQFFPQFPQFPF</b>      |  |
| 21  | 396(51) $\gamma$                         | TQQP.....QTQ             | 23 (9)   |
| 27  | 378(49) $\gamma$                         | QQP.....PQQQ             | 22 (10)  |

|  |           |                             |         |
|--|-----------|-----------------------------|---------|
| 28   | 371(48) γ | PQQQFIQP.....TY             | 22 (10) |
| 29   | 642(82) ω | PQQP.....L.....QQP          | 20 (8)  |
| 30   | 635(81) ω | PLQP.....QPQ                | 19 (5)  |
| 44   | 382(49) γ | .....QTQQPQQ                | 16 (6)  |
| 45   | 629(81) ω | PFPT.....S.....L.....QQ     | 16 (5)  |
| 46   | 643(82) ω | PLQP.....QQP                | 16 (6)  |
| 60   | 634(81) ω | PQQL.....L.....QQP          | 13 (3)  |
| 64   | 387(50) γ | .....T.....L.....QQPQQPF    | 13 (4)  |
| 62   | 641(82) ω | FPEI.....L.....LQP          | 13 (7)  |
| <b>Group 5: Homologues of Peptide 343 (overlap Groups 2 and 4)</b> |           |                             |         |
|  |           | <b>QQPFPPQQPQLPFPQ</b>      |         |
| 12   | 343(44) γ | QQP.....Q                   | 34 (11) |
| 16   | 393(51) γ | QLPFPQQP.....               | 27 (8)  |
| 19   | 335(43) γ | QQ.....Q.....PQ             | 25 (11) |
| 23   | 385(50) γ | .....QPPQ                   | 23 (7)  |
| 24   | 375(48) γ | P.....Q.....PQQ             | 23 (10) |
| 25   | 406(52) γ | QP.....L.....PQ             | 22 (8)  |
| 32   | 377(49) γ | P.....Q.....QPQ             | 19 (9)  |
| 34   | 415(53) γ | SQQP.....QS.....            | 18 (5)  |
| 37   | 413(53) γ | SKQP.....QS.....            | 18 (4)  |
| 38   | 380(49) γ | QPQQP.....                  | 18 (6)  |
| 43   | 409(53) γ | QP.....L.....Q.....L.....PQ | 16 (2)  |
| 47   | 389(50) γ | .....T.....Q.....QPPQ       | 16 (6)  |
| 58   | 407(52) γ | QP.....S.....Q.....PQ       | 14 (5)  |
| 63   | 399(51) γ | .....T.....Q.....LQQP       | 13 (5)  |
| 67   | 408(52) γ | QP.....SK.....Q.....PQ      | 12 (5)  |
| 71   | 379(49) γ | QQP.....Q.....P             | 11 (5)  |
| 72   | 397(51) γ | PQQP.....T.....Q.....       | 11 (3)  |
| <b>Group 6: Peptide 625</b>  |           |                             |         |
|  |           | <b>PIQPQQPFPQP</b>          |         |
| 26   | 625(80) ω | .....QQPQQPFP               | 22 (9)  |
| 57   | 624(80) ω | FTQPQQPT.....               | 14 (6)  |
| 65   | 628(80) ω | PF...W.....TQQSFPLQ         | 12 (4)  |
| <b>Group 7: Peptide 618</b>  |           |                             |         |
| 39   | 618(79) ω | PQQSFYQQQFPFPQYPQQ          | 18 (7)  |



**Table 21. Bioactivity of individual tTG-deamidated Pools 1-3 peptides in Subject C12:**  
(Peptide Numbers in Table 21 correspond to Peptide Numbers and their accompanying SEQ ID NOs, identified in Table 24)

| No.   | Sequence                      | %   | No. | Sequence                     | %  |
|---|-------------------------------|-----|-----|------------------------------|----|
| 8   | AVRWVPVQ <u>LQPQNPSQQQ</u> PQ | 100 | 23  | <u>LQPQNPSQQQ</u> PQEQVPLMQQ | 26 |
|   |                               | 85  |     |                              | 18 |
| 5   | MVRVTVPQ.....                 |     | 14  | .....EQVPLVQQ                |    |
|   |                               | 82  |     |                              | 18 |
| 6   | AVRVSVPQ.....                 |     | 15  | .....H.....EQVPLVQQ          |    |
|   |                               | 77  |     |                              | 18 |
| 3   | MVRVPVPQ.....H.....           |     | 17  | .....KQVPLVQQ                |    |
|   |                               | 67  |     |                              | 13 |
| 1   | AVRFPVPQ.....L.....           |     | 16  | .....D.....EQVPLVQQ          |    |
|   |                               | 59  |     |                              | 8  |
| 2   | MVRVPVPQ.....                 |     | 13  | .....EQVPLVQQ                |    |
|   |                               | 49  |     |                              | 5  |
| 9   | AVRVPVPQ.....L.....           |     | 22  | .....K.....EQVPLVQQ          |    |
|   |                               | 49  |     |                              | 3  |
| 7   | AVRVPVPQ.....                 |     | 18  | .....L.....EQVPLVQE          |    |
|   |                               | 33  |     |                              | 3  |
| 10  | MVRVPVPQ.....L.....           |     | 19  | .....L.....EQVPLVQE          |    |
|   |                               |     |     |                              |    |
| 4   | MVRVPMPQ.....D.....           | 15  | 20  | P.....P.....GQVPLVQQ         | 0  |
| 12  | AVRVPVPQ.....K.....           | 8   | 21  | P.....P.....RQVPLVQQ         | 0  |
| 11  | AVRVPVPQP.....P.....          | 0   |     |                              |    |
| Core sequence of epitope is underlined. Predicted deamidated sequence is: LQPENPSQEQQPE |                               |     |     |                              |    |

**Table 22: Phylogenetic groupings of wheat (*Triticum aestivum*) gliadins**

| <b>Alpha/beta-gliadins (n=61)</b> |  |        |  |
|-----------------------------------|--|--------|--|
| A1a1                              | AAA96525, EEWTA, P02863  | A1b13  | B22364, P04271                             |
| A1a2                              | CAB76963   | A2a1   | AAAB23109, CAA35238, P18573, S10015        |
| A1a3                              | AAA96276   | A2a2   | CAB76964                                   |
| A1a4                              | CAA26384, S07923   | A2b1   | P04724, T06500, AAA348282                  |
| A1a5                              | AAA34280   | A2b2   | D22364                                     |
| A1a6                              | P04728   | A2b3   | P04722, T06498, AAA34276                   |
| A1b1                              | CAB76962   | A2b4   | C22364                                     |
| A1b2                              | CAB76961   | A2b5   | CAB76956                                   |
| A1b3                              | BAA12318   | A3a1   | AAA34277, CAA26383, P04726, S07361         |
| A1b4                              | CAB76960   | A3a2   | I307187B, A27319, S13333                   |
| A1b5                              | CAB76958   | A3b1   | AAA96522                                   |
| A1b6                              | CAB76959   | A3b2i  | AAA34279, P04727,                          |
| A1b7                              | CAB76955   | A3b2ii | CAA26385, S07924                           |
| A1b8                              | AAA96524   | A3b3   | A22364, AAA34278, AAB23108, C61218, P04725 |
| A1b9                              | C'AA10257  | A4a    | P04723, AAA34283, T06504                   |
| A1b10                             | AAA96523, T06282   | A4b    | E22364                                     |
| A1b11                             | AAA17741, S52124   | A4c    | CAB76957                                   |
| A1b12                             | AAA34281   | A4d    | CAB76954                                   |
| <b>Gamma-gliadins (n=47)</b>      |  |        |  |
| GI1a                              | P08079, AAA34288, PS0094, CAC11079, AAD30556, CAC11057, CAC11065, CAC11056 | GI5a   | AAK84774, AAK84772                         |
| GI1b                              | CAC11089, CAC11064, CAC11080, CAC11078, AAD30440                           | GI5b   | AAK84773                                   |
| GI1c                              | CAC11087   | GI5c   | AAK84776                                   |
| GI1d                              | CAC11088   | GI6a   | JA0153, P21292, AAA34272, I507333A         |
| GI1e                              | CAC11055   | GI6b   | AAK84777                                   |
| GI2a                              | JS0402, P08453, AAA34289   | GI6c   | I802407A, AAK84775, AAK84780               |
| GI2b                              | AAF42989, AAK84779, AAK84779   | GI7    | AAB31090                                   |
| GI3a                              | AAK84778   | GI1a   | AAA34287, P04730, S07398                   |
| GI3b                              | CAB75404   | GI1b   | I209306A                                   |
| GI3c                              | BAA11251   | GI11a  | P04729                                     |
| GI4                               | EEWTC, P06659, AAA34274  | GI11b  | AAA34286                                   |
| <b>Omega-gliadins (n=3)</b>       |  |        |  |
| O1a                               | AAG17702   |        |  |
| O1b                               | P02865   |        |  |
| O1c                               | A59156   |        |  |

**Table 23. Synthetic peptides spanning all known wheat gliadin 12mers (Peptide Numbers 1 through 652 correspond to SEQ ID NO:107 through SEQ ID NO:758, respectively)**

| Protein       | Position* | Sequence                 | No. | Protein        | Position*                 | Sequence | No. |
|---------------|-----------|--------------------------|-----|----------------|---------------------------|----------|-----|
| <b>POOL 1</b> |           |                          |     |                |                           |          |     |
| A1A1          | 20        | AVRF PVPQ LQPQ NPSQ QLPQ | 1   | G1A2 33        | QQQL VPQL QQPL SQQP QQTTF | 331      |     |
| A1A2          | 20        | MVRV PVPQ LQPQ NPSQ QQPQ | 2   | G1A3 33        | QQQP FPQP HQPF SQQP QQTTF | 332      |     |
| A1B1          | 20        | MVRV PVPQ LQPQ NPSQ QHPQ | 3   | G1A4 33        | QQQP FLQP HQPF SQQP QQIF  | 333      |     |
| A1B2          | 20        | MVRV PMPQ LQPQ DPSQ QQPQ | 4   | G1A5 33        | QQQQ PFPQ PQQP FSQP PQQI  | 334      |     |
| A1B7          | 20        | MVRV TVPQ LQPQ NPSQ QQPQ | 5   | G1B3 33        | QQQQ PFPQ PQQP QQPF PQPQ  | 335      |     |
| A1B8          | 20        | AVRV SVQP LQPQ NPSQ QQPQ | 6   | G1C3 33        | QQQP FRQP QQPF YQQP QHTIF | 336      |     |
| A1B8          | 20        | AVRV PVPQ LQPQ NPSQ QQPQ | 7   | G1G4 33        | QQQP FPQP QQPF CQQP QRTI  | 337      |     |
| A1B10         | 20        | AVRW PVPQ LQPQ NPSQ QQPQ | 8   | G1G6 42        | QQQP FPQP QQPF CEQP QRIT  | 338      |     |
| <b>POOL 2</b> |           |                          |     |                |                           |          |     |
| A2B3          | 20        | AVRV PVPQ LQLQ NPSQ QQPQ | 9   | G1I4 42        | HQPF SQQP QQTIF PQPQ QTTF | 339      |     |
| A2B5          | 20        | MVRV PVPQ LQLQ NPSQ QQPQ | 10  | G1A2 42        | QQQL SQQP QQTIF PQPQ QTTF | 340      |     |
| A3A1          | 20        | AVRV PVPQ PQPQ NPSQ PQPQ | 11  | G1H4 42        | HQPF SQQP QQIF PQPQ QTTF  | 341      |     |
| A3B1          | 20        | AVRV PVPQ LQPK NPSQ QQPQ | 12  | G1A4 42        | QQPF SQQP QQIF PQPQ QTTF  | 342      |     |
| A1A1          | 28        | LQPQ NPSQ QQPQ EQVP LVQQ | 13  | G1B3 42        | QQQP QPFP PQQP PQQL FPQQ  | 343      |     |
| A1A2          | 28        | LQPQ NPSQ QQPQ EQVP LVQQ | 14  | G1C3 42        | QQPF YQQP QHTIF PQPQ QTCP | 344      |     |
| A1B1          | 28        | LQPQ NPSQ QHPQ EQVP LVQQ | 15  | G1G4 42        | QQPF CQQP QRIT PQPQ QTTH  | 345      |     |
| A1B2          | 28        | LQPQ DPSQ QQPQ EQVP LVQQ | 16  | G1G6 42        | QQPF CQQP QQTIF PQPQ QTTH | 346      |     |
| <b>POOL 3</b> |           |                          |     |                |                           |          |     |
| A2B1          | 28        | LQPQ NPSQ QQPQ KQVP LVQQ | 17  | G1G6 42        | QQPF CEQP QRIT PQPQ QTTH  | 347      |     |
| A2B3          | 28        | LQLQ NPSQ QQPQ EQVP LVQE | 18  | G1I4 50        | QQTIF PQPQ QTTF HQPQ QQPF | 348      |     |
| A2B5          | 28        | LQLQ NPSQ QQPQ EQVP LVQE | 19  | G1H4 50        | QQIF PQPQ QTTF HQPQ QQPF  | 349      |     |
| A3A1          | 28        | PQPQ NPSQ PQPQ QQVP LVQQ | 20  | G1A5 50        | QQIF PQPQ QTTF HQPQ QAFP  | 350      |     |
| A3A2          | 28        | PQPQ NPSQ PQPQ RQVP LVQQ | 21  | G1G4 50        | QRIT PQPQ QTTF HQPQ QTTF  | 351      |     |
| A3B1          | 28        | LQPK NPSQ QQPQ EQVP LVQQ | 22  | G1A5 58        | QTTF HQPQ QAFP PQQP TTFH  | 352      |     |
| A4A           | 28        | LQPQ NPSQ QQPQ EQVP LMQQ | 23  | G1G4 58        | QTTH HQPQ QTTF PQQP TYPH  | 353      |     |
| A1A1          | 36        | QLPQ EQVP LVQQ QQFL GQQQ | 24  | G1G6 58        | QTTH HQPQ QTTF PQQP TYPH  | 354      |     |
| <b>POOL 4</b> |           |                          |     |                |                           |          |     |
| A1B1          | 36        | QHQP EQVP LVQQ QQFL GQQQ | 25  | G1A5 66        | QAFP PQQP TTFH PQQP QFPQ  | 355      |     |
| A1B2          | 36        | QQPQ EQVP LVQQ QQFL GQQQ | 26  | G1C3 66        | QHIF PQQP QTCP HQPQ QQPF  | 356      |     |
| A1B12         | 36        | QQPQ EQVP LVQQ QQFL GQQQ | 27  | G1G4 66        | QTTF PQQP TYPH PQQP QFPQ  | 357      |     |
| A2A1          | 36        | QQPQ EQVP LVQQ QQFP GQQQ | 28  | G1G6 66        | QTTF PQQP TYPH PQQP QFPQ  | 358      |     |
| A2B1          | 36        | QQPQ KQVP LVQQ QQFP GQQQ | 29  | G1I4 73        | QTTF HQPQ QQPF PQQP PQQP  | 359      |     |
| A2B3          | 36        | QQPQ EQVP LVQE QQFP GQQQ | 30  | G1A2 73        | QTTF HQPQ QQVP PQQP PQQP  | 360      |     |
| A3A1          | 36        | PQPQ GQVP LVQQ QQFP GQQQ | 31  | G1B3 73        | QTTF HQPQ QQFS PQQP PQQP  | 361      |     |
| A3A2          | 36        | PQPQ RQVP LVQQ QQFP GQQQ | 32  | G1C3 73        | QTTF HQPQ QQFP PQQP PQQP  | 362      |     |
| <b>POOL 5</b> |           |                          |     |                |                           |          |     |
| A4A           | 36        | QQPQ EQVP LMQQ QQQF PQQQ | 33  | G1G4 73        | QTTF HQPQ QQFP QTQQ PQQP  | 363      |     |
| A1A1          | 44        | LVQQ QQFL GQQQ PFPQ PQQY | 34  | G1I4 81        | QQFP PQQP QQFP FLQP QQPF  | 364      |     |
| A1B1          | 44        | LVQQ QQFL GQQQ SFPP QQQY | 35  | G1A2 81        | QQVP PQQP PQQP FLQP QQPF  | 365      |     |
| A1B12         | 44        | LVQQ QQFL GQQQ PFPQ PQQY | 36  | G1B3 81        | QQFS PQQP PQQP FIQP QQPF  | 366      |     |
| A2A1          | 44        | LVQQ QQFP GQQQ PFPQ PQQY | 37  | G1H4 81        | QQFP PQQP PQQP FLQP RQPF  | 367      |     |
| A2B3          | 44        | LVQF QQFQ GQQQ PFPQ PQQY | 38  | G1A5 81        | QQFP PQQP PQQP FFPQ PQQP  | 368      |     |
| A3A1          | 44        | LVQQ QQFP GQQQ QFPF PQQY | 39  | G1G4 81        | QQFP QTQQ PQQP FPQP QQTTF | 369      |     |
| A4A           | 44        | LMQQ QQQF PQQQ EQFP PQQP | 40  | G1I4 89        | PQQQ FLQP QQPF PQQP QQPY  | 370      |     |
| <b>POOL 6</b> |           |                          |     |                |                           |          |     |
| A4D           | 44        | LMQQ QQQF PQQQ ERFP PQQP | 41  | G1B3 89        | PQQQ FIQP QQPF PQQP QQTY  | 371      |     |
| A1A1          | 53        | GQQQ PFPQ PQQY PQPQ PFPS | 42  | G1B3 89        | PQQQ FIQP QQPF QTTF QRFP  | 372      |     |
| A1A3          | 53        | GQQQ PFPQ PQQY PQPQ PFPS | 43  | G1H4 89        | PQQQ FLQP RQPF PQQP QQPY  | 373      |     |
| A1B1          | 53        | GQQQ SFPP QQQY PQPQ PFPS | 44  | G1A5 89        | PQQP FPQP PQQP FPQP QQPF  | 374      |     |
| A2B1          | 53        | GQQQ PFPQ PQQY PQQP PFPS | 45  | G1C3 89        | PQQP FPQP PQQP QPFP QQPF  | 375      |     |
| A3A1          | 53        | GQQQ QHFP QQQY PQPQ PFPS | 46  | G1G4 89        | PQQP FPQP QQTIF PQQP QLFP | 376      |     |
| A4A           | 53        | QQQF QFPF PQQY PHQQ PFPS | 47  | <b>POOL 49</b> |                           |          |     |
| A4D           | 53        | QQQF RFPP QQQY PHQQ PFPS | 48  | G1A5 97        | PQQQ FPQP QQPQ QPFP QQPQ  | 377      |     |
| <b>POOL 7</b> |           |                          |     |                |                           |          |     |
| A1A1          | 61        | QQPY PQPQ PFPS QLTP LQLQ | 49  | G1A5 105       | QQPQ QPFP QQPQ QPFP PQQP  | 378      |     |
| A1A3          | 61        | QQPY PQPQ PFPS LPTL QLQP | 50  | G1A5 113       | QQPQ QQFP PQQP PQQP FPQP  | 379      |     |
| A1B1          | 61        | QQPY PQPQ PFPS QQQY LQLQ | 51  | G1A5 121       | QPPQ PQQP FPQP PQQP LFPF  | 380      |     |
| A2B1          | 61        | QQPY PQPQ PFPS QQQY MQLQ | 52  | G1A5 126       | QQPF PQQP QQPF PQQP QQPF  | 381      |     |
| A4A           | 61        | QQPY PHQQ PFPS QQQY PQPQ | 53  | G1A5 126       | QQPF PQQP QQPF PQQP QQPF  | 382      |     |
| A1A1          | 69        | PFPS QLTP LQLQ PFPQ PQLP | 54  | G1A5 126       | QQPF PQQP QQTY PQRP QQPF  | 383      |     |
| A1B1          | 69        | PFPS QQQY LQLQ PFPQ PQLP | 55  | <b>POOL 50</b> |                           |          |     |

A1B10 69 PFPS QQPY LQLQ PFSQ PQLP  
**POOL 8**

A1B11 69 PFPS QQPY LQLQ PFLQ PQLP  
A1B12 69 PFPS QQPY LQLQ PFLQ PQPF  
A2A1 69 PFPS QQPY LQLQ PFQO PQLP  
A2B1 69 PFPS QQPY MQLQ PFQO PQLP  
A2B2 69 PFPS QQPY MQLQ PFQO PQPF  
A2B4 69 PFPS QQPY LQLQ PFQO PQPF  
A2B5 69 PFPS QQPY LQLQ PFPR PQLP  
A4A 69 PFPS QQPY PQFQ PFPP QLPY  
**POOL 9**

A4B 69 PFPS QQPY PQFQ PFQO PQPF  
A1A1 77 LQLQ PFQO PQLP YSQP QPFR  
A1A4 77 LQLQ PFQO PQLP YSQP QQFR  
A1B1 77 LQLQ PFQO PQLP YLQP QPFR  
A1B4 77 LQLQ PFQO PQLS YSQP QPFR  
A1B10 77 LQLQ PFSQ PQLP YSQP QPFR  
A1B11 77 LQLQ PFLQ PQLP YSQP QPFR  
A1B12 77 LQLQ PFLQ PQPF PQLP PQYS

**POOL 10**  
A2A1 77 LQLQ PFQO PQLP YPQF QLPY  
A2B1 77 MQLQ PFQO PQLP YPQF QLPY  
A2B2 77 MQLQ PFQO PQPF PQLP PYFQ  
A2B3 77 LQLQ PFQO PQLP YPQF QPFR  
A2B4 77 LQLQ PFQO PQPF PQLP PYFQ  
A2B5 77 LQLQ PFPR PQLP YPQF QPFR  
A3B1 77 LQLQ PFQO PQPF LPQL PYFQ  
A3B3 77 LQLQ PFQO PQPF PIQL PYFQ  
**POOL 11**

A4A 77 PQFQ PFPP QLPY PQTQ PFPP  
A4B 77 PQFQ PFQO PQPF PQLP PYFQ  
A1A1 85 PQLP YSQP QPFR PQQP YPQF  
A1A6 85 PQLP YSQP QQFR PQQP YPQF  
A1B1 85 PQLP YLQP QPFR PQQP YPQF  
A1B4 85 PQLS YSQP QPFR PQQP YPQF  
A1B6 85 PQLS YSQP QPFR PQQL YPQF  
A1B12 85 PQFQ PPQL PQYS PQFQ RPOQ  
**POOL 12**

A2A1 85 PQLP YPQF QLPY PQFQ LPYP  
A2B1 85 PQLP YPQF QLPY PQFQ PFPR  
A2B2 85 PQFQ PPQL PYFQ PQLP YPQF  
A2B3 85 PQLP YPQF QPFR PQQP YPQF  
A2B4 85 PQFQ PPQL PYFQ PQFQ RPOQ  
A3A1 85 PQFQ PPQL PYFQ PPFF SIQO  
**POOL 13**

A3B1 85 PQFQ LPQL PYFQ PQSF PFQO  
A3B3 85 PQFQ PPQL PYFQ PQSF PFQO  
A4A 85 QLPY PQTQ PFPP PQFQ PQFQ  
A4B 85 PQFQ PPQL PYFQ PQFQ PFQO  
A2A1 106 LPYP PQFQ FRPQ QSYF PQFQ  
A2B1 106 LPYP PQFQ FRPQ QSYF PQFQ  
A3A1 106 LPYP PQFQ RPYF PQFQ  
A3B1 106 LPQL PYFQ PQSF PFQO PYFQ  
**POOL 14**

A4A 106 PPQL PYFQ PQFQ PFQO PYFQ  
A1A1 112 QPFR PQFQ YPQF PQFQ SQPQ  
A1B6 112 QPFR PQQL YPQF PQFQ SQPQ  
A2A1 112 QPFR PQFQ YPQF PQFQ SQPQ  
A2B1 112 QPFR PQQS YPQF PQFQ SQPQ  
A1A1 112 PPHS PQFQ YPQF PQFQ PQFQ  
A2B1 112 QSFQ PQFQ YPQF RPYF LQFQ  
A3B2 112 QSFQ PQFQ YPQF RPYF LQFQ  
**POOL 15**

A3B3 112 QSFQ PQFQ YPQF PQFQ LQFQ  
A4A 112 QPFR PQFQ YPQF PQFQ PQFQ  
A1A1 120 YPQF PQFQ SQPQ QPIS QOQO  
A1B3 120 YPQF PQFQ SQPQ EPIS QOQO  
A2A1 120 YPQS PQFQ SQPQ QPIS QOQO  
A3A1 120 YPQF PQFQ PQFQ QPIS QOQO

108

56 GISA 126 QQPF PQFQ PQQL PFQO PQOQ  
GISC 126 QQPF PQFQ QAQL PFQO PQOQ  
57 GISA 126 QQTF PQFQ QLPF PQOQ QQPF  
58 GISA 134 QQPY PQOQ QQPF PQTQ PQOQ  
59 GISA 134 QQPF PQTQ PQOQ PFQO PQOQ  
60 GISA 134 QQTF PQFQ QQPF PQTQ PQOQ  
61 GISA 134 PQQL PFQO PQOQ PQOQ PFQO  
62 GISC 134 QAQL PFQO PQOQ PQFQ PQOQ  
63 **POOL 51**

64 GISA 134 QLPF PQOQ QQPF PQFQ PQOQ  
GISA 142 PQOQ PFQO PQOQ PFQO TQOQ  
65 GISA 150 PQOQ PFQO TQOQ PQFQ PQOQ  
66 GISA 158 TQOQ QQPF PQOQ QQPF PQTQ  
67 GISA 166 PQOQ QQPF PQTQ PQOQ PFQO  
68 GISA 170 QQPF PQTQ PQOQ LFPQ SQOQ  
69 GISA 170 QQPF PQTQ PQOQ PFQO LQOQ  
70 GISA 170 QQPF PQTQ PQOQ PFQO SQOQ  
71 **POOL 52**

72 GISA 170 QQPF PQTQ PQOQ PFQO SKQF  
GISA 170 QQPF PQFQ PQOQ PFQO LQOQ  
73 GISC 170 QQFL PQOQ PQOQ PFQO SQOQ  
74 GISA 170 QQPF PQOQ PQOQ PFQO SQOQ  
75 GISA 178 QPQO LFPQ SQOQ QQOQ SQOQ  
76 GISA 178 QPQO PFQO LQOQ QQOQ PQOQ  
77 GISA 178 QPQO PFQO SQOQ QQOQ PQOQ  
78 GISA 178 QPQO PFQO SKQF QQPF PQFQ  
79 **POOL 53**

80 GISA 178 QPQO PFQO LQOQ QQFL PQOQ  
GISA 186 SQOQ QQOQ SQOQ QQPF PQOQ  
81 GISA 186 LQOQ QQPF PQFQ QQLP PQOQ  
82 GISA 186 SQOQ QQPF PQFQ QQPF PQOQ  
83 GISA 186 SQOQ QQPF PQFQ PQOQ SFQO  
84 GISA 186 LQOQ QQFL PQFQ PQOQ PFQO  
85 GISC 186 SQOQ QQPF PQOQ PQOQ SFQO  
86 GISA 194 SQOQ QQPF PQOQ PQOQ PFQO  
87 **POOL 54**

88 GISA 194 PQOQ QQLP PQOQ PQQS PFQO  
GISA 194 PQOQ QQPF PQOQ PQQS PFQO  
89 GISA 194 PQOQ PQOQ SFQO QQOQ LQOQ  
90 GISA 194 PQOQ PQOQ PFQO QQOQ LQOQ  
91 GISC 194 PQOQ PQOQ SFQO QQOQ LQOQ  
92 GISA 202 QPQO PQQS FPQO QPFF IQPS  
93 GISA 202 QPQO PQQS FPQO QPFF IQPS  
94 GISA 202 QPQO PQQS FPQO QPFL IQQS  
**POOL 55**

95 GISA 210 FPQO QPFF IQPS LQOQ VNPC  
96 GISA 210 FPQO QPFF IQPS LQOQ LNPC  
97 GISA 210 FPQO QPFL IQPS LQOQ LNPC  
98 GISA 210 FPQO QQFL IQPY LQOQ MNPC  
99 GISA 210 FPQO QQPA IQSF LQOQ MNPC  
100 GISA 218 IQPS LQOQ VNPC KNFL LQOQ  
101 GISA 218 IQPS LQOQ LNPC KNFL LQOQ  
102 GISA 218 IQPS LQOQ LNPC KNFL LQOQ  
**POOL 56**

103 GISA 218 IQPY LQOQ MNPC KNFL LQOQ  
104 GISA 218 IQSF LQOQ MNPC KNFL LQOQ  
105 GISA 226 VNPC KNFL LQOQ KPVS LVSS  
106 GISA 226 LNPC KNFL LQOQ KPVS LVSS  
107 GISA 226 LNPC KNFL LQOQ KPVS LVSS  
108 GISA 226 MNPC KNFL LQOQ NPVS LVSS  
109 GISA 226 MNPC KNFL LQOQ NPVS LVSS  
110 GISA 234 LQOQ KPVS LVSS LWSM IWQO  
**POOL 57**

111 GISA 234 LQOQ KPVS LVSS LWSI IWQO  
112 GISA 234 LQOQ KPVS LVSS LWSM ILPR  
113 GISA 234 LQOQ NPVS LVSS LWSM ILPR  
114 GISA 234 LQOQ NPVS LVSS LWSI ILPR  
115 GISA 242 LVSS LWSM IWQO SDQO VMRO  
116 GISA 242 LVSS LWSI IWQO SDQO VMRO

385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446

|         |     |      |      |      |       |      |      |     |
|---------|-----|------|------|------|-------|------|------|-----|
| A3B1    | 120 | YPQQ | RPKY | LQPO | QPIS  | QQQA | 447  |     |
| A3B2    | 120 | YPQQ | RPMY | LQPO | QPIS  | QQQA | 448  |     |
| POOL 16 |     |      |      |      |       |      |      |     |
| A3B3    | 120 | YPQQ | QPQY | LQPO | QPIS  | QQQA | 449  |     |
| A1A1    | 128 | SQPP | QPIS | QQQQ | QQQQ  | QQQQ | 450  |     |
| A1B3    | 128 | SQPP | EPIS | QQQQ | QQQQ  | QQQI | 451  |     |
| A3A1    | 128 | PQPP | QPIS | QQQA | QQQQ  | QQQQ | 452  |     |
| A1A1    | 138 | QQQQ | QQQQ | QQQQ | QQQQ  | ILQQ | 453  |     |
| A1A6    | 138 | QQQQ | QQQQ | QQQQ | QEQQ  | ILQQ | 454  |     |
| A1B11   | 138 | QQQQ | QQQQ | QQQQ | QQQQ  | ILQQ | 455  |     |
| A2A1    | 138 | QQQQ | QQQQ | QQKQ | QQQQ  | QQQI | 456  |     |
| POOL 17 |     |      |      |      |       |      |      |     |
| A4B     | 139 | AQQQ | QQQQ | QQQQ | QQQQ  | TLQQ | 457  |     |
| A1A1    | 146 | QQQQ | QQQQ | ILQQ | ILQQ  | QLIP | 458  |     |
| A1A6    | 146 | QQQQ | QEQQ | ILQQ | ILQQ  | QLIP | 459  |     |
| A1B6    | 146 | QQQQ | QEQQ | ILQQ | MLQQ  | QLIP | 460  |     |
| A1B10   | 146 | QQQQ | QEQQ | ILQQ | ILQQ  | QLTP | 461  |     |
| A1B11   | 146 | QQQQ | QQQQ | ILQQ | ILQQ  | QLIP | 462  |     |
| A2A1    | 146 | QQQQ | QQQQ | QQQI | LQOI  | LQQQ | 463  |     |
| A3A2    | 146 | QQQQ | QQQQ | ILPQ | ILQQ  | QLIP | 464  |     |
| POOL 18 |     |      |      |      |       |      |      |     |
| A4A     | 146 | QQQQ | QQQQ | TLQQ | ILQQ  | QLIP | 465  |     |
| A1A1    | 163 | ILQQ | ILQQ | QLIP | CMVD  | VLQQ | 466  |     |
| A1B6    | 163 | ILQQ | MLQQ | QLIP | CMVD  | VLQQ | 467  |     |
| A1B10   | 163 | ILQQ | ILQQ | QLTP | CMVD  | VLQQ | 468  |     |
| A2B1    | 163 | ILQQ | ILQQ | QLIP | CRDV  | VLQQ | 469  |     |
| A3A2    | 163 | ILPQ | ILQQ | QLIP | CRDV  | VLQQ | 470  |     |
| A4A     | 163 | TLQQ | ILQQ | QLIP | CRDV  | VLQQ | 471  |     |
| A1A1    | 171 | QLIP | CMVD | VLQQ | HNIA  | HGRS | 472  |     |
| POOL 19 |     |      |      |      |       |      |      |     |
| A1A3    | 171 | QLIP | CMVD | VLQQ | HNIA  | HGRS | 473  |     |
| A1B2    | 171 | QLIP | CMVD | VLQQ | HNIA  | HGRS | 474  |     |
| A1B7    | 171 | QLIP | CMVD | VLQQ | HNIV  | HGRS | 475  |     |
| A1B10   | 171 | QLTP | CMVD | VLQQ | HNIA  | HGRS | 476  |     |
| A1B11   | 171 | QLIP | CMVD | VLQQ | HNIV  | HGRS | 477  |     |
| A2A1    | 171 | QLIP | CRDV | VLQQ | HSIA  | YGSS | 478  |     |
| A2B1    | 171 | QLIP | CRDV | VLQQ | HSIA  | HGSS | 479  |     |
| A2B3    | 171 | QLIP | CRDV | VLQQ | HNIA  | HGSS | 480  |     |
| POOL 20 |     |      |      |      |       |      |      |     |
| A3A1    | 171 | QLIP | CRDV | VLQQ | HNIA  | HARS | 481  |     |
| A3B1    | 171 | QLIP | CRDV | VLQQ | HNIA  | HASS | 482  |     |
| A1A1    | 179 | VLQQ | HNIA | HGRS | QVQL  | QSTY | 483  |     |
| A1A3    | 179 | VLQQ | HNIA | HGRS | QVQL  | QSTY | 484  |     |
| A1B2    | 179 | VLQQ | HNIA | HGRS | QVQL  | QSTY | 485  |     |
| A1B7    | 179 | VLQQ | HNIV | HGRS | QVQL  | QSTY | 486  |     |
| A1B10   | 179 | VLQQ | HNIA | HGRS | QVQL  | QSTY | 487  |     |
| A1B11   | 179 | VLQQ | HNIV | HGRS | QVQL  | QSTY | 488  |     |
| POOL 21 |     |      |      |      |       |      |      |     |
| A2A1    | 179 | VLQQ | HSIA | YGSS | QVQL  | QSTY | 489  |     |
| A2B1    | 179 | VLQQ | HSIA | HGSS | QVQL  | QSTY | 490  |     |
| A2B3    | 179 | VLQQ | HNIA | HGSS | QVQL  | QSTY | 491  |     |
| A3A1    | 179 | VLQQ | HNIA | HARS | QVQL  | QSTY | 492  |     |
| A3B1    | 179 | VLQQ | HNIA | HASS | QVQL  | QSTY | 493  |     |
| A4A     | 179 | VLQQ | HNIA | HASS | QVQL  | QSSY | 494  |     |
| A1A1    | 187 | HGRS | QVQL | QSTY | QLLQ  | ELCC | 495  |     |
| A1A3    | 187 | HGRS | QVQL | QSTY | QLLR  | ELCC | 496  |     |
| POOL 22 |     |      |      |      |       |      |      |     |
| A1B8    | 187 | HGRS | QVQL | QSTY | QLLR  | ELCC | 497  |     |
| A1B11   | 187 | HGRS | QVQL | QSTY | QLLQ  | ELCC | 498  |     |
| A2A1    | 187 | YGSS | QVQL | QSTY | QLVQ  | QLCC | 499  |     |
| A2B1    | 187 | HGSS | QVQL | QSTY | QLVQ  | QFCC | 500  |     |
| A2B3    | 187 | HGSS | QVQL | QSTY | QLVQ  | QLCC | 501  |     |
| A3A1    | 187 | HSIA | QVQL | QSTY | QPLQ  | QLCC | 502  |     |
| A3B1    | 187 | HASS | QVQL | QSTY | QLLQ  | QLCC | 503  |     |
| A4A     | 187 | HASS | QVQL | QSSY | QQLQ  | QLCC | 504  |     |
| POOL 23 |     |      |      |      |       |      |      |     |
| A1A1    | 195 | QSTY | QLLQ | QLCC | QHILW | QIPE | 505  |     |
| A1A3    | 195 | QSTY | QLLR | ELCC | QHILW | QIPE | 506  |     |
| A1B8    | 195 | QSTY | QLLR | ELCC | QHILW | QIPE | 507  |     |
| 417     | G1A | 242  | LVSS | LWSM | ILPR  | SDCQ | VMRQ | 447 |
| 418     | G1A | 242  | LVSS | LWSI | ILPP  | SDCQ | VMRQ | 448 |
| POOL 58 |     |      |      |      |       |      |      |     |
| 419     | G1A | 242  | LVSS | LWSM | ILPR  | SDCK | VMRQ | 449 |
| 420     | G1C | 242  | LVSS | LWSM | ILPR  | SDCQ | VMRQ | 450 |
| 421     | G1A | 242  | LVSS | LWSI | ILPR  | SDCQ | VMRQ | 451 |
| 422     | G1A | 250  | ILWQ | SDCQ | VMRQ  | QCCQ | QLAQ | 452 |
| 423     | G1A | 250  | ILPR | SDCQ | VMRQ  | QCCQ | QLAQ | 453 |
| 424     | G1A | 250  | ILPP | SDCQ | VMRQ  | QCCQ | QLAQ | 454 |
| 425     | G1A | 250  | ILPR | SDCK | VMRQ  | QCCQ | QLAR | 455 |
| 426     | G1C | 250  | ILPR | SDCQ | VMRQ  | QCCQ | QLAQ | 456 |
| POOL 59 |     |      |      |      |       |      |      |     |
| 427     | G1A | 258  | VMRQ | QCCQ | QLAQ  | IPQQ | LQCA | 457 |
| 428     | G1A | 258  | VMRQ | QCCQ | QLAR  | IPQQ | LQCA | 458 |
| 429     | G1C | 258  | VMRQ | QCCQ | QLAQ  | IPRQ | LQCA | 459 |
| 430     | G1A | 258  | VMRQ | QCCQ | QLAQ  | IPQQ | LQCA | 460 |
| 431     | G1A | 266  | QLAQ | IPQQ | LQCA  | AHIT | IVHS | 461 |
| 432     | G1B | 266  | QLAQ | IPQQ | LQCA  | AHIT | IVHS | 462 |
| 433     | G1A | 266  | QLAQ | IPQQ | LQCA  | AHIS | VVHS | 463 |
| 434     | G1A | 266  | QLAQ | IPQQ | LQCA  | AHIS | IVHS | 464 |
| POOL 60 |     |      |      |      |       |      |      |     |
| 435     | G1A | 266  | QLAR | IPQQ | LQCA  | AHIG | IVHS | 465 |
| 436     | G1C | 266  | QLAQ | IPRQ | LQCA  | AHIS | VVHS | 466 |
| 437     | G1A | 266  | QLAQ | IPQQ | LQCA  | AHIS | VAHS | 467 |
| 438     | G1A | 274  | LQCA | AHIT | IVHS  | IMQ  | QEQQ | 468 |
| 439     | G1B | 274  | LQCA | AHIT | IVHS  | IMQ  | QEQQ | 469 |
| 440     | G1A | 274  | LQCA | AHIS | VVHS  | IMQ  | QEQQ | 470 |
| POOL 61 |     |      |      |      |       |      |      |     |
| 441     | G1A | 274  | LQCA | AHIS | IVHS  | IMQ  | QEQQ | 471 |
| 442     | G1A | 274  | LQCA | AHIS | VVHS  | IMQ  | QEQQ | 472 |
| 443     | G1A | 274  | LQCA | AHIG | IVHS  | IMQ  | QEQQ | 473 |
| 444     | G1A | 274  | LQCA | AHIS | VAHS  | IMQ  | QEQQ | 474 |
| 445     | G1A | 282  | IVHS | IMQ  | QEQQ  | EQQQ | GMHI | 475 |
| 446     | G1B | 282  | IVHS | IMQ  | QEQQ  | QGMH | ILP  | 476 |
| 447     | G1A | 282  | IVHS | IMQ  | QEQQ  | QEQQ | QGID | 477 |
| 448     | G1A | 282  | IVHS | IMQ  | QEQQ  | EQRQ | GVQI | 478 |
| POOL 62 |     |      |      |      |       |      |      |     |
| 449     | G1A | 282  | IVHS | IMQ  | QEQQ  | EQQL | GVQI | 479 |
| 450     | G1A | 282  | IVHS | IMQ  | QEQQ  | QEQQ | QEQQ | 480 |
| 451     | G1C | 282  | IVHS | IVMQ | QEQQ  | QEQQ | ILRP | 481 |
| 452     | G1A | 282  | VAHS | IMQ  | QEQQ  | QGVV | ILRP | 482 |
| 453     | G1A | 290  | QEQQ | EQQQ | GMHI  | LLPL | YQQQ | 483 |
| 454     | G1A | 290  | QEQQ | QEQQ | QGMH  | ILPL | LSQH | 484 |
| 455     | G1B | 290  | QEQQ | QEQQ | QGMH  | ILPL | LSQH | 485 |
| 456     | G1A | 290  | QEQQ | EQRQ | GVQI  | LVPL | SQQQ | 486 |
| POOL 63 |     |      |      |      |       |      |      |     |
| 457     | G1A | 290  | QEQQ | EQQL | GVQI  | LVPL | SQQQ | 487 |
| 458     | G1A | 290  | QEQQ | QEQQ | QEQQ  | QEQQ | IQM  | 488 |
| 459     | G1C | 290  | QEQQ | QEQQ | ILRP  | LFQL | VQQG | 489 |
| 460     | G1A | 290  | QEQQ | QGVV | ILRP  | LFQL | QAQL | 490 |
| 461     | G1A | 298  | QEQQ | QEQQ | IQM   | RPLF | QLVQ | 491 |
| 462     | G1A | 305  | GMHI | LLPL | YQQQ  | QVQG | GTLV | 492 |
| 463     | G1A | 305  | GMHI | FLPL | LSQH  | QVQG | GSIV | 493 |
| 464     | G1B | 305  | GMHI | FLPL | SQQQ  | QVQG | GSIV | 494 |
| POOL 64 |     |      |      |      |       |      |      |     |
| 465     | G1A | 305  | GVQI | LVPL | SQQQ  | QVQG | GTLV | 495 |
| 466     | G1A | 305  | GVQI | LVPL | SQQQ  | QVQG | GHLV | 496 |
| 467     | G1A | 305  | GIQI | MRPL | FQLV  | QCGG | IIQP | 497 |
| 468     | G1C | 305  | GIQI | LRPL | FQLV  | QCGG | IIQP | 498 |
| 469     | G1A | 305  | GVPI | LRPL | FQLA  | QGLG | IIQP | 499 |
| 470     | G1A | 313  | YQQQ | QVQG | GTLV  | QCGG | IIQP | 500 |
| 471     | G1A | 313  | LSQH | QVQG | GSIV  | QCGG | IIQP | 501 |
| 472     | G1B | 313  | SQQQ | QVQG | GSIV  | QCGG | IIQP | 502 |
| POOL 65 |     |      |      |      |       |      |      |     |
| 473     | G1A | 313  | SQQQ | QVQG | GTLV  | QCGG | IIQP | 503 |
| 474     | G1A | 313  | SQQQ | QVQG | GHLV  | QCGG | IIQP | 504 |
| 475     | G1A | 321  | GTLV | QCGG | IIQP  | QQPA | QLEA | 505 |
| 476     | G1A | 321  | GSIV | QCGG | IIQP  | QQPA | QLEA | 506 |
| 477     | G1A | 321  | FQLV | QCGG | IIQP  | QQPA | QLEV | 507 |

|                |     |       |       |       |      |       |                |                |      |      |      |      |      |      |     |
|----------------|-----|-------|-------|-------|------|-------|----------------|----------------|------|------|------|------|------|------|-----|
| A2A1           | 195 | QSTY  | QLVQ  | QLQK  | QQW  | QIPE  | 178            | G6A            | 321  | FQLA | QQLG | ILQP | QQPA | QLEQ | 508 |
| A2B1           | 195 | QSTY  | QLVQ  | QFCC  | QQW  | QIPE  | 179            | G1A            | 329  | IIQP | QQPA | QLEV | IRSL | VLQT | 509 |
| A3A1           | 195 | QSTY  | QPLQ  | QLCC  | QQW  | QIPE  | 180            | G3A            | 329  | IIQP | QQPA | QLEV | IRSL | VLQT | 510 |
| A3B1           | 195 | QSTY  | QLLQ  | QLCC  | QQW  | QIPE  | 181            | <b>POOL 66</b> |      |      |      |      |      |      |     |
| A4A            | 195 | QSSY  | QQLQ  | QLCC  | QQW  | QIPE  | 182            | G3C            | 329  | IIQP | QQPA | QLEV | IRSS | VLQT | 511 |
| <b>POOL 24</b> |     |       |       |       |      |       | 183            | G5C            | 329  | IIQP | QQPA | QYEV | IRSL | VLRT | 512 |
| A1A1           | 203 | ELCC  | QHILW | QIPE  | QSRC | QAIIH | 184            | G6A            | 329  | IIQP | QQPA | QLEV | IRSL | VLRT | 513 |
| A1B6           | 203 | ELCC  | QHILW | QIPE  | QSRC | QAIIH | 184            | G1A            | 337  | QLEA | IRSL | VLQT | LPTM | CNVY | 514 |
| A1B10          | 203 | ELCC  | QHILW | QIPE  | KLQC | QAIIH | 185            | G1A            | 337  | QLEA | IRSL | VLQT | LPTM | CNVY | 515 |
| A2A1           | 203 | QLCC  | QQW   | QIPE  | QSRC | QAIIH | 186            | G1A            | 337  | QLEV | IRSL | VLQT | LPTM | CNVY | 516 |
| A2B1           | 203 | QFCC  | QQW   | QIPE  | QSRC | QAIIH | 187            | G3C            | 337  | QLEV | IRSL | VLQT | LPTM | CNVY | 517 |
| A3B1           | 203 | QLCC  | QQW   | QIPE  | QSRC | QAIIH | 188            | G1A            | 337  | QLEV | IRSL | VLQT | LPTM | CNVY | 518 |
| <b>POOL 25</b> |     |       |       |       |      |       | <b>POOL 67</b> |                |      |      |      |      |      |      |     |
| A3B3           | 203 | GLCC  | QQW   | QIPE  | QSRC | QAIIH | 189            | G5C            | 337  | QYEV | IRSL | VLRT | LPNM | CNVY | 519 |
| A4A            | 203 | QLCC  | QQW   | QIPE  | QSRC | QAIIH | 190            | G1A            | 337  | QLEG | IRSL | VLRT | LPNM | CNVY | 520 |
| A1A1           | 211 | QIPE  | QSRC  | QAIIH | NVNH | AIIH  | 191            | G1A            | 345  | VLQT | LPTM | CNVY | VPPE | CSII | 521 |
| A1B3           | 211 | QIPE  | QSRC  | QAIIH | NVNH | AIIH  | 192            | G1A            | 345  | VLQT | LPTM | CNVY | VPPE | CSII | 522 |
| A1A6           | 211 | QIPE  | QSRC  | QAIIH | NVNH | AIIH  | 193            | G1A            | 345  | VLQT | LPTM | CNVY | VPPE | CSII | 523 |
| A1B1           | 211 | QIPE  | QSRC  | QAIIH | NVNH | AIIH  | 194            | G1A            | 345  | VLQT | LPTM | CNVY | VPPE | CSII | 524 |
| A1B10          | 211 | QIPE  | KLQC  | QAIIH | NVNH | AIIH  | 195            | G5C            | 345  | VLRT | LPNM | CNVY | VRPD | CSII | 525 |
| A2A1           | 211 | QIPE  | QSRC  | QAIIH | NVNH | AIIH  | 196            | G6A            | 345  | VLRT | LPNM | CNVY | VRPD | CSII | 526 |
| <b>POOL 26</b> |     |       |       |       |      |       | <b>POOL 68</b> |                |      |      |      |      |      |      |     |
| A3B3           | 211 | QIPE  | QSRC  | QAIIH | NVAH | AIIH  | 197            | G1A            | 353  | CNVY | VPPE | CSII | KAPE | SSVY | 527 |
| A4A            | 211 | QIPE  | QSRC  | QAIIH | NVNH | AIIH  | 198            | G1A            | 353  | CNVY | VPPE | CSII | RAPE | ASIV | 528 |
| A1A1           | 219 | QAIIH | NVNH  | AIIH  | HQQQ | KQQQ  | 199            | G1A            | 353  | CNVY | VPPE | CSII | RAPE | ASIV | 529 |
| A1A6           | 219 | QAIIH | NVNH  | AIIH  | HQQQ | KQQQ  | 200            | G5A            | 353  | CNVY | VPPE | CSII | RAPE | ASIV | 530 |
| A1B3           | 219 | QAIIH | NVNH  | AIIH  | HQQQ | KQQQ  | 201            | G5C            | 353  | CNVY | VRPD | CSII | NAPE | ASIV | 531 |
| A1B9           | 219 | QAIIH | NVNH  | AIIH  | HQQQ | KQQQ  | 202            | G6A            | 353  | CNVY | VPPE | CSII | NAPE | ASIV | 532 |
| A1B13          | 219 | QAIIH | NVNH  | AIIH  | HQQQ | QQQQ  | 203            | G1A            | 361  | CSII | KAPE | SSVY | AGIG | GQ   | 533 |
| A2B3           | 219 | QAIIH | NVNH  | AIIH  | HQQH | HHHQ  | 204            | G1A            | 361  | CSII | RAPE | ASIV | AGIG | GQ   | 534 |
| <b>POOL 27</b> |     |       |       |       |      |       | <b>POOL 69</b> |                |      |      |      |      |      |      |     |
| A3A1           | 219 | QAIIH | NVNH  | AIIH  | HQQQ | RQQQ  | 205            | G3A            | 361  | CSII | RAPE | ASIV | AGIG | GQYR | 535 |
| A3B1           | 219 | QAIIH | NVNH  | AIIH  | HQQQ | QQQQ  | 206            | G1A            | 361  | CSII | RAPE | ASIV | AGIG | GQ   | 536 |
| A3B3           | 219 | QAIIH | NVAH  | AIIH  | HQQQ | QQQQ  | 207            | G5A            | 361  | CSII | RAPE | ASIV | AGIG | GQ   | 537 |
| A4A            | 219 | QAIIH | NVNH  | AIIH  | HHHQ | QQQQ  | 208            | G5C            | 361  | CSII | NAPE | ASIV | AGIG | GQ   | 538 |
| A1A1           | 227 | AIIH  | HQQQ  | KQQQ  | QPSQ | QVSF  | 209            | G6A            | 361  | CSII | NVNY | ADIG | AGIG | GQ   | 539 |
| A1A6           | 227 | AIIH  | HQQQ  | KQQQ  | QPSQ | QVSF  | 210            | G1I            | 1    | POQP | FLPQ | POQS | FLWQ | SOQ  | 540 |
| A1B2           | 227 | AIIH  | HQQQ  | KQQQ  | QPSQ | QVSF  | 211            | G1I            | 9    | POQS | FLWQ | POQP | FLQ  | POQP | 541 |
| A1B10          | 227 | AIIH  | HQQQ  | KQQQ  | QSSQ | VVSF  | 212            | G1I            | 17   | SQOP | FLQ  | POQP | SQOP | POQP | 542 |
| <b>POOL 28</b> |     |       |       |       |      |       | <b>POOL 70</b> |                |      |      |      |      |      |      |     |
| A1B13          | 227 | AIIH  | HQQQ  | QQQQ  | EQQK | QLQ   | 213            | G1I            | 25   | POQP | SPOQ | QVVQ | QIIS | PATP | 543 |
| A2A1           | 227 | AIIH  | HQQQ  | QQQQ  | QQQQ | QPLS  | 214            | G1I            | 33   | QOVV | QIIS | PATP | TTIP | SAGK | 544 |
| A2B3           | 227 | AIIH  | HQQH  | HHHQ  | QQQQ | QQQQ  | 215            | G1I            | 41   | PATP | TTIP | SAGK | PISA | PFQ  | 545 |
| A2B4           | 227 | AIIH  | HQQH  | HHHQ  | QQQQ | QLQ   | 216            | G1I            | 49   | SAGK | PTSA | PFQ  | QQQQ | HQOL | 546 |
| A3A1           | 227 | AIIH  | HQQQ  | RQQQ  | PSSQ | VSLQ  | 217            | G1I            | 57   | PFQ  | QQQQ | HPVQ | HQSL | QPOV | 547 |
| A3B1           | 227 | AIIHM | HQQE  | QQQQ  | QLQ  | QQQQ  | 218            | G1I            | 65   | HQQL | QQQQ | IPVQ | QPSI | QLQ  | 548 |
| A3B2           | 227 | AIIHM | HQQQ  | QQQQ  | EQQK | QLQ   | 219            | G1I            | 73   | VPV  | QPSI | VLQ  | NPCK | VFQ  | 549 |
| A4A            | 227 | AIIH  | HHHQ  | QQQQ  | QPSQ | QVSF  | 220            | G1I            | 81   | VLQ  | NPCK | VFQ  | QQCS | PVAM | 550 |
| <b>POOL 29</b> |     |       |       |       |      |       | <b>POOL 71</b> |                |      |      |      |      |      |      |     |
| A1A1           | 235 | KQQQ  | QPSQ  | QVSF  | QOPL | QOYP  | 221            | G1I            | 89   | VLQ  | QQCS | PVAM | PQRL | ARSQ | 551 |
| A1A6           | 235 | KQQQ  | QPSQ  | QVSF  | QOPL | QOYP  | 222            | G1I            | 97   | PVAM | PQRL | ARSQ | MLQ  | SSCH | 552 |
| A1B2           | 235 | KQQQ  | QLSS  | QVSF  | QOPL | QOYP  | 223            | G1I            | 105  | ARSQ | MLQ  | SSCH | VMMQ | QQCC | 553 |
| A1B10          | 235 | KQQQ  | PSSQ  | VSFQ  | POPQ | QYPL  | 224            | G1I            | 113  | SSCH | VMMQ | QQCC | QLPQ | IPQ  | 554 |
| A2A1           | 235 | QQQQ  | EQQK  | QLQ   | QQQQ | QQQL  | 225            | G1I            | 121  | CCQ  | QLPQ | IPQ  | SRQY | AIQ  | 555 |
| A2B4           | 235 | HHHQ  | EQQK  | QLQ   | QQQQ | QQQL  | 226            | G1I            | 127B | POIP | QQR  | YEA  | RAI  | YSII | 556 |
| A3A1           | 235 | RQQQ  | PSSQ  | VSLQ  | POPQ | QYPS  | 227            | G1I            | 129  | IPQ  | SRQY | AIQ  | IYIS | IIH  | 557 |
| A3B1           | 235 | QQQQ  | LQQQ  | QQQQ  | LQQQ | QQQQ  | 228            | G1I            | 137  | AIQ  | IYIS | IIH  | EQQQ | VQGS | 558 |
| <b>POOL 30</b> |     |       |       |       |      |       | <b>POOL 72</b> |                |      |      |      |      |      |      |     |
| A4A            | 235 | QQQQ  | QPSQ  | QVSQ  | QOYQ | EQYF  | 229            | G1I            | 145  | IIH  | EQQQ | VQGS | IQSQ | QQQ  | 559 |
| A1B13          | 243 | QLQ   | QQQ   | QQQ   | QQQ  | QQQ   | 230            | G1I            | 153  | QVQS | IQSQ | QQQ  | QLQ  | QCVS | 560 |
| A1B13          | 251 | QQQ   | QQQ   | QQQ   | QQQ  | QQS   | 231            | G1I            | 161  | QQQ  | QLQ  | QCVS | QPSQ | QSQ  | 561 |
| A2A1           | 260 | QQQ   | QQQ   | QPS   | VSF  | QOPL  | 232            | G1I            | 169  | VSF  | QOPL | QPSQ | QLQ  | QO   | 562 |
| A2B1           | 260 | QQQ   | QQQ   | QPS   | VSF  | QOPL  | 233            | G1I            | 177  | QQQ  | QLQ  | QPSQ | QLQ  | QO   | 563 |
| A2B3           | 260 | HHHQ  | QQQ   | QQQ   | QPS  | QPS   | 234            | G1I            | 185  | QOQ  | QLQ  | QPSQ | QLQ  | QO   | 564 |
| A3A1           | 260 | QQQQ  | QQQ   | QPSQ  | QVSF | QOPL  | 235            | <b>POOL 73</b> |      |      |      |      |      |      |     |
| A3B1           | 289 | QPS   | QVSF  | QOPL  | QOYP | SGQG  | 236            | G1I            | 193  | QGT  | LEVH | QIAQ | LEVH | TMIC | 565 |
| <b>POOL 31</b> |     |       |       |       |      |       | 237            | G1I            | 201  | QIAQ | LEVH | QIAQ | LEVH | TMIC | 566 |
| A2B1           | 289 | QPS   | QVSF  | QOPL  | QOYP | SGQG  | 238            | G1I            | 209  | QPS  | QVSF | QOPL | QOYP | SGQG | 567 |
| A3A1           | 289 | QPSQ  | QVSF  | QOPL  | QOYP | SSQV  | 239            | G1I            | 217  | TMIC | LEVH | QIAQ | LEVH | TMIC | 568 |

|                |     |                          |     |                |                               |     |
|----------------|-----|--------------------------|-----|----------------|-------------------------------|-----|
|                |     |                          | 111 |                |                               |     |
| A1A1           | 293 | QVSF QOPL QOYP LGQG SFRP | 239 | GII 225        | PLYR TTTS VPHG VGTG VGAY      | 569 |
| A1A6           | 293 | QESF QOPL QOYP LGQG SFRP | 240 | GIII 1A        | 1 TTTR TTPT PTIS SNNN HHFR    | 570 |
| A1B2           | 293 | QVSF QOPL QOYP LGQG SFRP | 241 | GIII 1A        | 9 PTIS SNNN HHFR SNNN HHFH    | 571 |
| A2A1           | 293 | QVSF QOPL QOYP SGQG SFQP | 242 | GIII 1A        | 17 HHFR SNNN HHFH SNNN QFYR   | 572 |
| A2B1           | 293 | QVCF QOQO QOYP SGQG SFQP | 243 | <b>POOL 74</b> |                               |     |
| A2B3           | 293 | QVSF QOPL QOYP SGQG FFQP | 244 | GIII 1A        | 25 HHFH SNNN QFYR NNNN PGHN   | 573 |
| <b>POOL 32</b> |     |                          | 245 | GIII 1A        | 41 PGHN NPLN NNNN SNNN SPNS   | 574 |
| A2B5           | 293 | QVSF QOPL QOYP SGQG FFQP | 246 | GIII 1A        | 49 NNNN SNNN SPNS HHNN SPNN   | 575 |
| A3A1           | 293 | QVSL QOPL QOYP SGQG FFQP | 247 | GIII 1A        | 57 SPNS HHNN SNNN NFQY HTHP   | 576 |
| A3B1           | 293 | QVSF QOPL QOYP SSQV SFQP | 248 | GIII 1A        | 65 SPNN NFQY HTHP SNNH NLPH   | 577 |
| A3B2           | 293 | QVSF QOPL QOYP SSQG SFQP | 249 | GIII 1A        | 73 HTHP SNNH NLPH TNNT QQQQ   | 578 |
| A4A            | 293 | QVSF QOPL QOYP SSQV SFQS | 250 | GIII 1A        | 81 NLPH TNNT QQQQ PPFS QQQQ   | 579 |
| A1A1           | 301 | QOYP LGQG SFRP SQQN PQAQ | 251 | <b>POOL 75</b> |                               |     |
| A1B2           | 301 | QOYP LGQG SFRP SQQN SQAQ | 252 | GIII 1A        | 89 QQQQ PPFS QQQQ PPFS QQQQ   | 581 |
| A2A1           | 301 | QOYP SGQG SFQP SQQN PQAQ | 253 | GIII 1A        | 97 QQQQ PPFS QQQQ PVLP QQSP   | 582 |
| <b>POOL 33</b> |     |                          | 254 | GIII 1A        | 105 QQQQ PVLP QQSP FSQG QQLV  | 583 |
| A2B3           | 301 | QOYP SGQG FFQP SQQN PQAQ | 255 | GIII 1A        | 113 QQQS FSQG QQLV LPQP QQQQ  | 584 |
| A2B5           | 301 | QOYP SGQG FFQP FQQN PQAQ | 256 | GIII 1A        | 121 QQLV LPQP QQQQ QLVQ QQIP  | 585 |
| A3A1           | 301 | QOYP SGQG FFQP SQQN PQAQ | 257 | GIII 1A        | 129 QQQQ QLVQ QQIP IVQP SVLQ  | 586 |
| A3B1           | 301 | QOYP SSQG SFQP SQQN PQAQ | 258 | GIII 1A        | 137 QQIP IVQP SVLQ QLNP CKVF  | 587 |
| A4A            | 301 | EQYP SGQV SFQS SQQN PQAQ | 259 | <b>POOL 76</b> |                               |     |
| A1B1           | 309 | SFRP SQQN PLAQ GSVQ PQQL | 260 | GIII 1A        | 153 CKVF LQQQ CSPV AMPO RLAR  | 589 |
| A1A1           | 309 | SFRP SQQN PQAQ GSVQ PQQL | 261 | GIII 1A        | 161 CSPV AMPO RLAR SQMW QQSS  | 590 |
| <b>POOL 34</b> |     |                          | 262 | GIII 1A        | 169 RLAR SQMW QQSS CHVM QQQC  | 591 |
| A1A3           | 309 | SFRP SQQN PQTG GSVQ PQQL | 263 | GIII 1A        | 177 QQSS CHVM QQQC CQQL QQIP  | 592 |
| A1B2           | 309 | SFRP SQQN SQAQ GSVQ PQQL | 264 | GIII 1A        | 185 QQQC CQQL QQIP EQSR YEAI  | 593 |
| A1B3           | 309 | SFRP SQQN PQDQ GSVQ PQQL | 265 | GIII 1A        | 193 QQIP EQSR YEAI RAIL YSII  | 594 |
| A1B4           | 309 | SFRP SQQN PRAQ GSVQ PQQL | 266 | GIII 1A        | 201 YEAI RAIL YSII LQEQ QQGF  | 595 |
| A2A1           | 309 | SFQP SQQN PQAQ GSVQ PQQL | 267 | GIII 1A        | 209 YSII LQEQ QQGF VQPQ QQQP  | 596 |
| A2B3           | 309 | FFQP SQQN PQAQ GSVQ PQQL | 268 | <b>POOL 77</b> |                               |     |
| A2B5           | 309 | FFQP FQQN PQAQ GSVQ PQQL | 269 | GIII 1A        | 217 QQGF VQPQ QQQP QQSQ QGVN  | 597 |
| A3A1           | 309 | FFQP SQQN PQAQ GSVQ PQQL | 270 | GIII 1A        | 225 QQQP QQSQ QGVN QSQQ QSQQ  | 598 |
| <b>Pool 35</b> |     |                          | 271 | GIII 1A        | 233 QGVN QSQQ QSQQ QLQG CSFQ  | 599 |
| A3B1           | 309 | SFQP SQLN PQAQ GSVQ PQQL | 272 | GIII 1A        | 241 QSQQ QLQG CSFQ PQQP QLQG  | 600 |
| A3B2           | 309 | SFQP SQQN PQAQ GSVQ PQQL | 273 | GIII 1A        | 249 CSFQ PQQP QLQG PQQP QQQQ  | 601 |
| A4A            | 309 | SFQS SQQN PQAQ GSVQ PQQL | 274 | GIII 1A        | 257 QLQG PQQP QQQQ QVLQ GTFL  | 602 |
| A1A1           | 317 | PQAQ GSVQ PQQL PQFE EIRN | 275 | GIII 1A        | 263 QQQQ QVLQ GTFL PQHP IAHIL | 603 |
| A1A3           | 317 | PQTG GSVQ PQQL PQFE EIRN | 276 | GIII 1A        | 271 GTFL PQHP IAHIL EAVT SIAL | 604 |
| A1A6           | 317 | PQAQ GSVQ PQQL PQFE IRNL | 277 | <b>POOL 78</b> |                               |     |
| A1B1           | 317 | PLAQ GSVQ PQQL PQFE EIRN | 278 | GIII 1A        | 279 IAHIL EAVT SIAL RTLP TMCN | 605 |
| <b>POOL 36</b> |     |                          | 279 | GIII 1A        | 287 SIAL RTLP TMCN VNNV LYSA  | 606 |
| A1B3           | 317 | PQDQ GSVQ PQQL PQFE EIRN | 280 | GIII 1A        | 295 TMCN VNNV LYSA TTSV PFGV  | 607 |
| A1B4           | 317 | PRAQ GSVQ PQQL PQFE EIRN | 281 | GIII 1A        | 303 LYSA TTSV PFGV GTGV GAY   | 608 |
| A2B3           | 317 | PQAQ GSVQ PQQL PQFE EIRN | 282 | GIII 1B        | 26 SCIS GLER PWQG QPLP PQQS   | 609 |
| A2B5           | 317 | PQAQ GSVQ PQQL PQFE AIRN | 283 | GIII 1B        | 34 PWQG QPLP PQQS FSQG PPFS   | 610 |
| A3B1           | 317 | PQAQ GSVQ PQQL PQFA EIRN | 284 | GIII 1B        | 42 PQQS FSQG PPFS QQQQ QPLP   | 611 |
| A4A            | 317 | PQAQ GSVQ PQQL PQFE EIRN | 285 | GIII 1B        | 50 PPFS QQQQ QPLP PQQS FSQG   | 612 |
| <b>Pool 37</b> |     |                          | 286 | <b>Pool 79</b> |                               |     |
| A1A1           | 325 | PQQL PQFE EIRN LALQ TLPA | 287 | GIII 1B        | 58 QPLP PQQS FSQG QPFP SQQQ   | 613 |
| A1A6           | 325 | PQQL PQFE IRNL ALQT LPAM | 288 | GIII 1B        | 66 FSQG QPFP SQQQ PILS QQP    | 614 |
| A1B12          | 325 | PQQL PQFE EIRN LARK      | 289 | GIII 1B        | 74 SQQQ PILS QQPFP FSQG QQP   | 615 |
| A2A1           | 325 | PQQL PQFE EIRN LALE TLPA | 290 | O 1A           | 17 ATAA RELN PSNK ELQS PQQS   | 616 |
| A2B5           | 325 | PQQL PQFE AIRN LALQ TLPA | 291 | O 1A           | 25 PSNK ELQS PQQS FSQY QQPF   | 617 |
| A3B1           | 325 | PQQL PQFA EIRN LALQ TLPA | 292 | O 1A           | 33 PQQS FSQY QQPF PQQP YPQO   | 618 |
| A4A            | 325 | PQQL PQFQ EIRN LALQ TLPA | 293 | O 1A           | 41 QQPF PQQP YPQO PYPQ QQPY   | 619 |
| A1A1           | 333 | EIRN LALQ TLPA MCNV YIPP | 294 | O 1A           | 49 YPQO PYPQ QQPY FSQG PFPF   | 620 |
| <b>POOL 38</b> |     |                          | 295 | <b>POOL 80</b> |                               |     |
| A1A3           | 333 | EIRN LALQ TLPS MCNV YIPP | 296 | O 1A           | 57 QQPY FSQG PFPF PQQP FPEQ   | 621 |
| A2A1           | 333 | EIRN LALE TLPA MCNV YIPP | 297 | O 1A           | 65 PFPF PQQP FPEQ SQQP FTQP   | 622 |
| A1A1           | 341 | TLPA MCNV YIPP YCTI APFG | 298 | O 1A           | 73 FPEQ SQQP FTQP QQPF PQQP   | 623 |
| A1A3           | 341 | TLPS MCNV YIPP YCTI APFG | 299 | O 1A           | 81 FTQP QQPF PQQP QQPF PQQP   | 624 |
| A1B1           | 341 | TLPA MCNV YIPP YCTI VPFG | 300 | O 1A           | 89 PQQP QQPF PQQP QQPF QQPF   | 625 |
| A1B4           | 341 | TLPA MCNV YIPP YCAM APFG | 301 | O 1A           | 97 PQQP QQPF QQPF QQPF PFPW   | 626 |
| A1B9           | 341 | TLPA MCNV YIPP YCTI TPFG | 302 | O 1A           | 105 QQPF QQPF PFPW PQQP PFPQ  | 627 |
| <b>Pool 39</b> |     |                          | 303 | O 1A           | 113 PFPW PQQP FFPQ TQQS FPLQ  | 628 |
| A2A1           | 341 | TLPA MCNV YIPP YCTI APVG | 304 | <b>POOL 81</b> |                               |     |
|                |     |                          | 305 | O 1A           | 121 FFPQ TQQS FPLQ PQQP FFPQ  | 629 |

|                                   |                                       |     |
|-----------------------------------|---------------------------------------|-----|
| A2B2 341 TLPA MCNV YIPP YCST TIAP | 300 O 1A 129 FPLQ PQQP FPQQ PQQP FPQP | 630 |
| A3A1 341 TLPR MCNV YIPP YCST TIAP | 301 O 1A 137 FPQQ PQQP FPQP QLPP PQQS | 631 |
| A3A2 341 TLPR MCNV YIPP YCST TTAP | 302 O 1A 145 FPQP QLPP PQQS EQLL PQQL | 632 |
| A3B1 341 TLPA MCNV YIPP HCST TIAP | 303 O 1A 153 PQQS EQLL PQQL QQPF PLQP | 633 |
| A1A1 349 YIPP YCTI APFG IFGT NYR  | 304 O 1A 161 PQQL QQPF PLQP QQPF PQQP | 634 |
| A1B1 349 YIPP YCTI VPFG IFGT NYR  | 305 O 1A 169 PLQP QQPF PQQP QPQP PQPQ | 635 |
| A1B4 349 YIPP YCAM APFG IFGT NYR  | 306 O 1A 177 PQQP QQPF PQQP QPQP VQPQ | 636 |
| <b>Pool 40</b>                    |                                       |     |
| A1B5 349 YIPP YCTM APFG IFGT NYR  | 307 O 1A 185 PQQP QPQP VQPQ QSPQ PQSQ | 637 |
| A1B9 349 YIPP YCTI TPFG IFGT N    | 308 O 1A 193 VQPQ QSPQ PQSQ QSQQ PFAQ | 638 |
| A2A1 349 YIPP YCTI APVG IFGT NYR  | 309 O 1A 201 PQSQ QSQQ PFAQ PQQL FPEL | 639 |
| A2B2 349 YIPP YCST TIAP VGIF GTN  | 310 O 1A 209 PFAQ PQQL FPEL QQPI PQQP | 640 |
| A3A2 349 YIPP YCST TTAP FGIF GTN  | 311 O 1A 217 FPEL QQPI PQQP QQPF PLQP | 641 |
| A3B1 349 YIPP HCST TIAP FGIF GTN  | 312 O 1A 225 PQQP QQPF PLQP QQPF PQQP | 642 |
| A3B3 349 YIPP HCST TIAP FGIS GTN  | 313 O 1A 233 PLQP QQPF PQQP QQPF PQQP | 643 |
| A4D 350 IPPY CSTI IAPF GFHG TNYR  | 314 O 1A 241 PQQP QQPF PQQP PQSQ PQQP | 644 |
| <b>Pool 41</b>                    |                                       |     |
| GI1A 17 GTAN MQVD PSSQ VQWP QQQP  | 315 O 1A 249 PQQP PQSQ PQQP QPQP PQQP | 645 |
| GI2A 17 GTAN IQVD PSGQ VQWL QQQL  | 316 O 1A 257 PQQP QPQP PQQP PYGS SLTS | 646 |
| GI3A 17 ATAN MQVD PSGQ VPWP QQQP  | 317 O 1A 265 PQQP PYGS SLTS IGGQ      | 647 |
| GI3B 19 MN IQVD PSGQ VPWP QQQP LP | 318 O 1B 1 ARQL NPSD QELQ SPQQ LYTPQ  | 648 |
| GI4 17 ATAN MQAD PSGQ VQWP QQQP   | 319 O 1B 9 QELQ SPQQ LYTPQ QPQP QPQP  | 649 |
| GI5A 17 TTAN IQVD PSGQ VQWP QQQQ  | 320 O 1C 1 SRLL SPRG KELH TPQE QFTQ   | 650 |
| GI5C 17 ATAN MQVD PSGQ VQWP QQQP  | 321 O 1C 9 KELH TPQE QFTQ QQQP PQPQ   | 651 |
| GI7 20 QIVF PSGQ VQWP QQQQ PFP    | 322 O 1C 17 QFTQ QQQP PQPQ QFTQ       | 652 |
| <b>Pool 42</b>                    |                                       |     |
| GI1A 25 PSSQ VQWP QQQP VPQP HQPF  | 323                                   |     |
| GI2A 25 PSGQ VQWL QQQL VPQL QQPPL | 324                                   |     |
| GI3A 25 PSGQ VPWP QQQP FPQP HQPF  | 325                                   |     |
| GI4 25 PSGQ VQWP QQQP FLQP HQPF   | 326                                   |     |
| GI5A 25 PSGQ VQWP QQQP PFPQ PQQP  | 327                                   |     |
| GI5C 25 PSGQ VQWP QQQP FRQP QQPF  | 328                                   |     |
| GI6A 25 PSGQ VQWP QQQP FPQP QQPF  | 329                                   |     |
| GI1A 33 QQQP VPQP HQPF SQQP QQTF  | 330                                   |     |

---

\*Position of N-terminal residue in  $\alpha$ -,  $\gamma$ 1-,  $\gamma$ 2-,  $\gamma$ 3-, or  $\omega$  consensus sequence



|                              |       |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|------------------------------|-------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| CELESTAC SUBJECT             | C14   | C15 | C16 | C17 | C18 | C19 | C20 | C21 | C22 | C23 | C24 | C25 | C26 | C27 | C28 |
| H1A-DQ                       | 22    | 22  | 22  | 23  | 22  | 22  | 2X  | 2X  | 2X  | 2X  | 28  | 28  | 2X  | 2X  | 2X  |
| ANTIGEN CHALLENGE            | WHEAT |     |     |     |     |     |     |     |     |     | RYE |     |     |     |     |
| DURATION OF CHALLENGE (DAYS) | 3     | 3   | 6   | 3   | 6   | 3   | 3   | 3   | 3   | 0.5 | 3   | 1   | 3   | 3   | 3   |
| DOMINANT PEPTIDE ELISPOT SFC | 203   | 46  | 96  | 195 | 114 | 136 | 29  | 57  | 129 | 259 | 50  | 18  | 163 | 52  | 229 |
| BLANK ELISPOT SFC            | 1     | 2   | 4   | 0.5 | 3   | 1   | 1   | 4   | 3   | 1   | 2   | 1   | 2   | 1   | 2   |

[illegible]

[illegible]



[illegible]













[illegible]

[illegible]

[illegible]

[illegible]



>3 x 31